A New Auxiliary Model Approach For Fermionic Criticality

Insights on Mottness in Strongly Correlated Systems

ABHIRUP MUKHERJEE

Presubmission Open Seminar

IISER KOLKATA

Emergent Phenomena in Quantum Matter group
IISER Kolkata

EPQM

Acknowledgements

Publications

New J. Phys. 25 113011. **AM**, N S Vidhyadhiraja, A Taraphder, S Lal. (2023)

J. Phys. A: Math. Theor. 57 275401. **AM**, S Patra, S Lal. (2024)

New J. Phys. 28 013503. **AM**, S R. Hassan, A Mukherjee, N S. Vidhyadhiraja, A Taraphder, S Lal. (2026)

in progress. **AM**, S Lal. (2026)

Publications

New J. Phys. 25 113011. **AM**, N S Vidhyadhiraja, A Taraphder, S Lal. (2023)

J. Phys. A: Math. Theor. 57 275401. **AM**, S Patra, S Lal. (2024)

New J. Phys. 28 013503. **AM**, S R. Hassan, A Mukherjee, N S. Vidhyadhiraja, A Taraphder, S Lal. (2026)

in progress. **AM**, S Lal. (2026)

Phys. Rev. B 105, 085119. A Mukherjee, **AM**, N S. Vidhyadhiraja, A Taraphder, S Lal. (2022)

J. Phys.: Condens. Matter 35 315601. S Patra, **AM**, A Mukherjee, N S Vidhyadhiraja, A Taraphder, S Lal. (2023)

Phys. Rev. B 111, L140414. A Kumar N M, S Mukherjee, **AM**, A Punjal, S Purwar, T Setti, S Prabhu S., S Lal, N Kamaraju. (2025)

arXiv:2602.20600. D Debata, **AM**, S Lal. (2026)

Outline of Thesis

Chap 1. Introduction

Chap 2. Methods and Preliminaries

Chap 3. Kondo frustration via charge fluctuations: Mott localisation in $d = \infty$ [**New J. Phys. 25 113011**]

Chap 4. A New Auxiliary Model Mapping For Correlated Lattice Models [**New J. Phys. 28 013503**]

Chap 5. Mott Criticality as the Confinement Transition of a Pseudogap-Mott Metal in $d = 2$ [**New J. Phys. 28 013503**]

Chap 6. Competing tendencies In Heavy Fermion Systems: Kondo screening vs. Mott localisation [**in progress**]

Chap 7. Holographic entanglement renormalisation for fermionic quantum matter [**J. Phys. A: Math. Theor. 57 275401**]

Chap 8. Conclusions

Introduction To Fermionic Criticality and Quantum Materials

What Do Condensed Matter Theorists Do?

What Do Condensed Matter Theorists Do?

Explain Experiments

Build and solve theoretical models that explain observations

Examples: Quantum Hall effect, superconductivity

What Do Condensed Matter Theorists Do?

Explain Experiments

Build and solve theoretical models that explain observations

Examples: Quantum Hall effect, superconductivity

Predict Experiments

Identify interesting theoretical models that predict novel phenomena

Examples: Abrikosov vortex lattice, spin liquids

What Do Condensed Matter Theorists Do?

Explain Experiments

Build and solve theoretical models that explain observations

Examples: Quantum Hall effect, superconductivity

Predict Experiments

Identify interesting theoretical models that predict novel phenomena

Examples: Abrikosov vortex lattice, spin liquids

Classify experiments

Come up with frameworks that explain a large class of phenomena

Examples: Landau theory of phase transition

Quantum Materials

Class of materials that **cannot** be well-approximated by “classical” description

Quantum Materials

Class of materials that **cannot** be well-approximated by “classical” description



topological properties
(*topological insulators*)

Quantum Materials

Class of materials that **cannot** be well-approximated by “classical” description



topological properties
(*topological insulators*)

non-local entanglement
(*spin liquids*)



Quantum Materials

Class of materials that **cannot** be well-approximated by “classical” description



topological properties
(*topological insulators*)

non-local entanglement
(*spin liquids*)

collective excitations
(*strange metals*)



Quantum Materials

Class of materials that **cannot** be well-approximated by “classical” description



topological properties
(*topological insulators*)

non-local entanglement
(*spin liquids*)

collective excitations
(*strange metals*)



Needs dramatically **new ideas** and methods!

- **non-local/collective** order parameters for transition
- ability to work with a large number of **interacting particles**
- ability to work with **emergent** degrees of freedom

Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Mott Insulation



Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Mott Insulation





Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Mott Insulation



Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Mott Insulation



- Not accessible from 1-particle theory
- result of **manybody effects**

Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Mott Insulation



How do we describe the transition from a metal into a Mott insulator?

- Not accessible from 1-particle theory
- result of **manybody effects**

Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Strange Metallicity



Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Strange Metallicity



Conventional metallic theory predicts $R \propto T^2$



Source of $R \propto T^2$?

Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Strange Metallicity



Conventional metallic theory predicts $R \propto T^2$



Source of $R \propto T^2$?

Single particles must propagate finite number of lattice spacing



Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Phenomena: Strange Metallicity



Conventional metallic theory predicts $R \propto T^2$



Source of $R \propto T^2$?

Who is carrying current in strange metals?

Single particles must propagate finite number of lattice spacing



Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Schematic phase diagram of **copper oxide materials**



Strongly Correlated Phenomena

Materials with strong **local** correlation, **low dimensionality**, multiple electronic **orbitals**

Schematic phase diagram of **copper oxide materials**



Theoretical Models of Correlated Materials

Hubbard Model

Theoretical Models of Correlated Materials

Hubbard Model

- Correlated electrons hopping on a lattice
-

Theoretical Models of Correlated Materials

Hubbard Model

- Correlated electrons hopping on a lattice
- Hopping probability is t.

Theoretical Models of Correlated Materials

Hubbard Model

- Correlated electrons hopping on a lattice
- Hopping probability is t.

- Energy cost U if two occupy same site.



Theoretical Models of Correlated Materials

Hubbard Model

- Correlated electrons hopping on a lattice
- Hopping probability is t.

- Energy cost U if two occupy same site.



Hamiltonian: $-t \sum_{i,j,\sigma} (c^{\dagger}_{i,\sigma} c_{j,\sigma} + \text{h.c.}) + \sum_i U n_{i,\uparrow} n_{i,\downarrow}$

Theoretical Models of Correlated Materials

Hubbard Model

- Correlated electrons hopping on a lattice
- Hopping probability is t.

- Energy cost U if two occupy same site.



Hamiltonian: $-t \sum_{i,j,\sigma} (c^{\dagger}_{i,\sigma} c_{j,\sigma} + \text{h.c.}) + \sum_i U n_{i,\uparrow} n_{i,\downarrow}$

Theoretical Models of Correlated Materials

Hubbard Model

- Correlated electrons hopping on a lattice
- Hopping probability is t.

$t \Longrightarrow$ Delocalisation
$U \Longrightarrow$ Localisation

- Energy cost U if two occupy same site.

🔥 Competition
🫂 Correlations

Hamiltonian: $-t \sum_{i,j,\sigma} (c^{\dagger}_{i,\sigma} c_{j,\sigma} + \text{h.c.}) + \sum_i U n_{i,\uparrow} n_{i,\downarrow}$

Quantum Impurity Models as Auxiliary Models



Lattice models of correlated electrons are **hard to solve** directly.

Quantum Impurity Models as Auxiliary Models



Lattice models of correlated electrons are **hard to solve** directly.

Idea: replace full problem with a simpler **auxiliary model** that captures the local correlations.



Quantum Impurity Models as Auxiliary Models

Auxiliary Model for Interacting Quantum Systems?

Quantum Impurity Models as Auxiliary Models

Auxiliary Model for Interacting Quantum Systems?

Auxiliary Model has to be **interacting** in order to capture temporal dynamics!

What is an appropriate auxiliary problem for lattice models with local correlation?



Quantum Impurity Models as Auxiliary Models

Auxiliary Model for Interacting Quantum Systems?

Auxiliary Model has to be **interacting** in order to capture temporal dynamics!

What is an appropriate auxiliary problem for lattice models with local correlation?



Insight from Limit of Infinite Dimensions!

Quantum Impurity Models as Auxiliary Models

Auxiliary Model for Interacting Quantum Systems?

Auxiliary Model has to be **interacting** in order to capture temporal dynamics!

What is an appropriate auxiliary problem for lattice models with local correlation?



Insight from Limit of Infinite Dimensions!

Local physics ⟹ **Quantum Impurity Models**

Core Idea of Dynamical Mean-Field Theory



Quantum Impurity Models as Auxiliary Models

Auxiliary Model for Interacting Quantum Systems?

Auxiliary Model has to be **interacting** in order to capture temporal dynamics!

What is an appropriate auxiliary problem for lattice models with local correlation?



Insight from Limit of Infinite Dimensions!

Local physics ⟹ **Quantum Impurity Models**

Core Idea of Dynamical Mean-Field Theory



- Tractable
- Contains dynamics

(ω –dependence of correlations)

Quantum Impurity Models as Auxiliary Models

Auxiliary Model for Interacting Quantum Systems?

Auxiliary Model has to be **interacting** in order to capture temporal dynamics!

What is an appropriate auxiliary problem for lattice models with local correlation?



Local physics of correlated lattice models can be mapped on to impurity model.

Insight from Limit of Inf

Local physics ⟹ **Quantum Impurity Models**

Core Idea of Dynamical Mean-Field Theory



- Tractable
- Contains dynamics

(ω –dependence of correlations)

Phenomenology of Quantum Impurity Models

Kondo Model

- Impurity spin S_d + conduction bath
- Spin-flip scattering

$$H = \sum_{k,\sigma} \varepsilon_k n_{k,\sigma} + \frac{1}{4} J S_d^z s_0^z + \frac{1}{2} S_d^+ s_0^- + \frac{1}{2} S_d^- s_0^+$$



Phenomenology of Quantum Impurity Models

Kondo Model

- Impurity spin S_d + conduction bath
- Spin-flip scattering

$$H = \sum_{k,\sigma} \varepsilon_k n_{k,\sigma} + \frac{1}{4} J S_d^z s_0^z + \frac{1}{2} S_d^+ s_0^- + \frac{1}{2} S_d^- s_0^+$$



Low-energy ($T \ll 1$) Properties: Kondo Screening

- Kondo coupling generates **entanglement**
- Impurity DOS supports **gapless excitations**
- Impurity electron can **delocalise** via bath



Phenomenology of Quantum Impurity Models

Kondo Model

- Impurity spin S_d + conduction bath
- Spin-flip scattering

$$H = \sum_{k,\sigma} \varepsilon_k n_{k,\sigma} + \frac{1}{4} J S_d^z s_0^z + \frac{1}{2} S_d^+ s_0^- + \frac{1}{2} S_d^- s_0^+$$



Low-energy ($T \ll 1$) Properties: Kondo Screening

- Kondo coupling generates **entanglement**
- Impurity DOS supports **gapless excitations**
- Impurity electron can **delocalise** via bath

$\Longrightarrow$ **metallic state** in auxiliary model language.



Phenomenology of Quantum Impurity Models

Kondo Model

- Impurity spin S_d + conduction bath
- Spin-flip scattering

$$H = \sum_{k,\sigma} \varepsilon_k n_{k,\sigma} + \frac{1}{4} J S_d^z s_0^z + \frac{1}{2} S_d^+ s_0^- + \frac{1}{2} S_d^- s_0^+$$



Metallic State



How Do We Get an Insulating State?

Phenomenology of Quantum Impurity Models

Kondo Model

- Impurity spin S_d + conduction bath
- Spin-flip scattering

$$H = \sum_{k,\sigma} \varepsilon_k n_{k,\sigma} + \frac{1}{4} J S_d^z s_0^z + \frac{1}{2} S_d^+ s_0^- + \frac{1}{2} S_d^- s_0^+$$



Metallic State



How Do We Get an Insulating State?

- Tweak the bath
- Destroy screening
- Local moment emerges



Phenomenology of Quantum Impurity Models

Kondo Model

- Impurity spin S_d + conduction bath
- Spin-flip scattering

$$H = \sum_{k,\sigma} \varepsilon_k n_{k,\sigma} + \frac{1}{4} J S_d^z s_0^z + \frac{1}{2} S_d^+ s_0^- + \frac{1}{2} S_d^- s_0^+$$



Kondo screening ⟹ Metallicity

Local moment ⟹ Mott insulation

Metallic State



- Tweak the bath
- Destroy screening
- Local moment emerges

Lating State?



Outstanding Questions

Appropriate impurity model to describe the Mott transition (locally)?



General Approach To Incorporate Non-Local Fluctuations?

Relation Between Various Correlated Phases?



Minimal Model For Kondo Breakdown and Mott Transition

A Route to Mott Localisation in Infinite Dimensions

New J. Phys. 25 113011 (2023)
AM, N S Vidhyadhiraja, A Taraphder, S Lal

Mott Transition In Infinite Dimensions

DMFT Results on Hubbard Model in $D = \infty$

Dynamical mean-field theory sees a metal-insulator transition by **self-consistent** determination of bath.



Mott Transition In Infinite Dimensions

DMFT Results on Hubbard Model in $D = \infty$

- numerical implementation means precise form of self-consistent impurity model **not known**
- low-energy theory for $T = 0$ **critical point** not known

Dynamical mean-field theory sees a metal-insulator transition by **self-consistent** determination of bath.



Mott Transition In Infinite Dimensions

DMFT Results on Hubbard Model in $D = \infty$

- numerical implementation means precise form of self-consistent impurity model **not known**
- low-energy theory for $T = 0$ **critical point** not known

Our Approach: Add Local Correlation in Bath

$-|U_b|\, n_{0\uparrow} n_{0\downarrow}$



Dynamical mean-field theory sees a metal-insulator transition by **self-consistent** determination of bath.



Phase Diagram and Impurity Phase Transition

Phase Diagram and Impurity Phase Transition





- Impurity has gapless excitations at $U_b = 0$ (**metal**)
- Now, make bath correlation **attractive** (favours 2 electrons)
-

Phase Diagram and Impurity Phase Transition





- Impurity has gapless excitations at $U_b = 0$ (**metal**)
- Now, make bath correlation **attractive** (favours 2 electrons)
- **Destroys** Kondo screening for $|U_b| > J/4$

Local Attractive Bath Correlaton Is The Trick!

Local Attractive Bath Correlaton induces phase transition not present in vanilla impurity model.



Local Attractive Bath Correlaton Is The Trick!

Local Attractive Bath Correlaton induces phase transition not present in vanilla impurity model.







Local Attractive Bath Correlaton Is The Trick!

Local Attractive Bath Correlaton induces phase transition not present in vanilla impurity model.







Local Attractive Bath Correlaton Is The Trick!

Local Attractive Bath Correlaton induces phase transition not present in vanilla impurity model.



State exactly at transition is a **special metal** – more on this later!





Local Attractive Bath Correlaton Is The Trick!

Local Attractive Bath Correlaton induces phase transition not present in vanilla impurity model.



State exactly at transition is a **special metal** – more on this later!

Impurity model with local attractive bath correlation is a good auxiliary system for Mott transition.





Capturing Spatial Fluctuations and Lattice Geometry

Impurity-Lattice Mapping Beyond Mean-Field

New J. Phys. 28 013503 (2026)
AM, S R. Hassan, A Mukherjee, N S. Vidhyadhiraja, A Taraphder, S Lal

Extending This To The More Interesting $D = 2$ Case



Previous demonstration worked because

- **non-local fluctuations vanish** in $d \longrightarrow \infty$ (mean-field is exact)
- Largely insensitive to **lattice details** $\{\varepsilon_k\}$

Extending This To The More Interesting $D = 2$ Case



Previous demonstration worked because

- **non-local fluctuations vanish** in $d \longrightarrow \infty$ (mean-field is exact)
- Largely insensitive to **lattice details** $\{\varepsilon_k\}$

Low-Dimensional Lattices Are More Interesting!

Extending This To The More Interesting $D = 2$ Case



Previous demonstration worked because

- **non-local fluctuations vanish** in $d \longrightarrow \infty$ (mean-field is exact)
- Largely insensitive to **lattice details** $\{\varepsilon_k\}$

Low-Dimensional Lattices Are More Interesting!



Paths extremely constrained

Extending This To The More Interesting $D = 2$ Case



Previous demonstration worked because

- **non-local fluctuations vanish** in $d \longrightarrow \infty$ (mean-field is exact)
- Largely insensitive to **lattice details** $\{\varepsilon_k\}$

Low-Dimensional Lattices Are More Interesting!



Paths extremely constrained



Momentum density-of-states highly anisotropic

Extending This To The More Interesting $D = 2$ Case





- All paths symmetric
- No non-trivial k – dependence

Low-Dimensional Lattices Are More Interesting!



Paths extremely constrained



Momentum density-of-states highly anisotropic

Extending This To The More Interesting $D = 2$ Case



- Spatial fluctuations
- Lattice geometry

- All paths symmetric
- No non-trivial k – dependence

Low-Dimensional Lattices Are More Interesting!



Paths extremely constrained



Momentum density-of-states highly anisotropic

Extending This To The More Interesting $D = 2$ Case



- Spatial fluctuations
- Lattice geometry

- All paths symmetric
- No non-trivial k – dependence

Low-Dimensional Lattices Are More Interesting!



Appropriate auxiliary model for a correlated system on a 2D square lattice?



Momentum
lensity-of-states
highly anisotropic

Why is 2D Important?

- Many quantum materials are layered.
- Weak interlayer coupling $\longrightarrow$ effectively 2D

Canonical Example: Copper Oxide Materials –> Plethora of Mysterious Phases

Why is 2D Important?

- Many quantum materials are layered.
- Weak interlayer coupling → effectively 2D

Canonical Example: Copper Oxide Materials -> Plethora of Mysterious Phases



Why is 2D Important?

- Many quantum materials are layered.
- Weak interlayer coupling ⟶ effectively 2D

Canonical Example: Copper Oxide Materials –> Plethora of Mysterious Phases



- Fermi liquid: conventional metal → described by **Landau Fermi-liquid theory**
- Excitations: long-lived **weakly-interacting** quasiparticles
- Mott insulator: interaction-driven insulator at half-filling (often AF ordered)
- Both endpoints are **well understood**

Why is 2D Important?

- Many quantum materials are layered.
- Weak interlayer coupling → effectively 2D

Canonical Example: Copper Oxide Materials -> Plethora of Mysterious Phases

- Overarching theory for strange metal not available





Why is 2D Important?

- Many quantum materials are layered.
- Weak interlayer coupling → effectively 2D

Canonical Example: Copper Oxide Materials -> Plethora of Mysterious Phases

- Overarching theory for strange metal not available
- Nature of excitations within PG: unclear



ARPES shows **partial gaps** on Fermi surface



Why is 2D Important?

- Many quantum materials are layered.
- Weak interlayer coupling → effectively 2D

Canonical Example: Copper Oxide Materials -> Plethora of Mysterious Phases

- Overarching theory for strange metal not available
- Nature of excitations within PG: unclear
- How are the phases related to each other?



Inter-relations between the phases not clear!

- How do the electrons in Fermi liquid **morph** to form the strange meta/PG phases?
- How do the excitations of the PG phase **bind** to stabilise the Mott insulator?

Embedding An Impurity Into The Lattice

Embed an impurity into a 2D square lattice



- Impurity-Bath spin-interaction (J) can **probe lattice structure**

Embedding An Impurity Into The Lattice

Embed an impurity into a 2D square lattice



Contrast with $D = \infty$ limit



- Impurity-Bath spin-interaction (J) can **probe lattice structure**
- Impurity-Bath interaction is purely local → insensitive to lattice geometry

Embedding An Impurity Into The Lattice

Embed an impurity into a 2D square lattice

Contrast with $D = \infty$ limit



Lattice-embedded impurity model is a reasonable auxiliary model for 2D lattice problems.

- Impurity-Bath spin-interaction (J) can **probe lattice structure**
- Impurity-Bath interaction is purely local → insensitive to lattice geometry

Embedding An Impurity Into The Lattice

Embed an impurity into a 2D square lattice

Contrast with $D = \infty$ limit



Lattice-embedded impurity model is a reasonable auxiliary model for 2D lattice problems.

Lattice geometry encoded in impurity-bath correlation.

- Impurity-Bath spin-interaction (J) can **probe lattice structure**
- Impurity-Bath interaction is purely local → insensitive to lattice geometry

Reconstructing Spatial Fluctuations



$\Longrightarrow$

Spatial Translation symmetry needs to be restored

$\Longrightarrow$



Reconstructing Spatial Fluctuations

Spatial Translation symmetry needs to be restored: <u>Bloch's Theorem!</u>

Reconstructing Spatial Fluctuations

Spatial Translation symmetry needs to be restored: <u>Bloch's Theorem!</u>

Translation of
1-Particle States:
Wannier → Bloch

Reconstructing Spatial Fluctuations

Spatial Translation symmetry needs to be restored: Bloch's Theorem!

Translation of
1-Particle States:
Wannier → Bloch

Many-electron
Translation
Operators:
$\mathbb{T}(a) = \Pi_i T_i(a)$



Reconstructing Spatial Fluctuations

Spatial Translation symmetry needs to be restored: Bloch's Theorem!

Many-electron Translation Operators:
$\mathbb{T}(a) = \Pi_i T_i(a)$



Many-electron translation:
Impurity → Lattice

$$H_{\text{lat}} = \sum_i \mathbb{T}(r_i) H_{\text{aux}} \mathbb{T}^\dagger(r_i)$$

$$-t \sum_{<i,j>} (c_i^\dagger c_j + \text{h.c.}) + U \sum_i n_{i,\uparrow} n_{i,\downarrow} + \sum_{<i,j>} S_i \cdot S_j$$



Reconstructing Spatial Fluctuations

Spatial Translation symmetry needs to be restored: Bloch's Theorem!

Many-electron translation:
Impurity → Lattice

$$H_{\text{lat}} = \sum_i \mathbb{T}(r_i) H_{\text{aux}} \mathbb{T}^\dagger(r_i)$$

$$-t \sum_{<i,j>} (c_i^\dagger c_j + \text{h.c.}) + U \sum_i n_{i,\uparrow} n_{i,\downarrow} + \sum_{<i,j>} S_i \cdot S_j$$



Extended Hubbard Model



Plan: Tiling Procedure + Many-body Bloch's theorem

Connects Hamiltonians, Eigenstates, Correlations

$$\Psi_{\text{lat}} = \sum_i \mathbb{T}(\mathbb{r}_i) \psi_{\text{aux}}$$

$$G_{\text{loc}} = \mathcal{G}(r_d, r_d) + C_0 \mathcal{G}(r_d, r_0) + C_1 \mathcal{G}(r_d, r_1) + \ldots$$

Reconstructing Spatial Fluctuations

Spatial Translation symmetry needs to be restored: Bloch's Theorem!

Many-electron translation:
Impurity → Lattice



Extended Hubbard Model



$$H_{\text{lat}} = \sum_i \mathbb{T}$$

$$-t \sum_{<i,j>} (c_i^\dagger c_j + \text{h.c.}) +$$

- Study lattice-embedded impurity model.
- Use tiling relations to reconstruct low-energy physics of lattice model.

Plan: Tiling Procedure + Many-body Bloch's theorem

$$\Psi_{\text{lat}} = \sum_i \mathbb{T}(\mathbb{r}_i)\psi_{\text{aux}}$$

Connects Hamiltonians, Eigenstates, Correlations

$$G_{\text{loc}} = \mathcal{G}(r_d, r_d) + C_0\mathcal{G}(r_d, r_0) + C_1\mathcal{G}(r_d, r_1) + \ldots$$

Phase Diagram: Lattice-Embedded Impurity Model

We use local spectral function to characterise phases

Competing tendencies In a Multi-Orbital System

Kondo screening vs. Mott localisation in a Heavy-Fermion model

(in progress)

AM, S Lal

Holography of Entanglement in 2D Free Fermions

Emergent Geometry and Fermionic Criticality

J. Phys. A: Math. Theor. 57 275401 (2024)
AM, S Patra, S Lal

What is The Holographic Principle?

QUANTUM FIELD THEORY IN D – DIMENSIONS $\Longrightarrow$ GRAVITY THEORY IN $D+1$ – DIMENSIONS





- Extra dimension arises from **renormalisation group flow** of quantum theory
- **Geometric measures** in +1 dimension arise from **quantum measures** on the boundary

What is The Holographic Principle?

Quantum field theory in D – dimensions $\Longrightarrow$ Gravity theory in $D+1$ – dimensions



Renormalisation Group Flow



- Extra dimension arises from **renormalisation group flow** of quantum theory
- **Geometric measures** in +1 dimension arise from **quantum measures** on the boundary

Questions?

- Can we demonstrate this in the case of a prototypical metal-insulator transition?
- What effect does the critical point have on the emergent dimensions?

Entanglement RG Of 2D Dirac Electrons On a Torus

non-interacting **Dirac electrons** moving on a torus

- **Entanglement entropy** of a subsystem of length l
- RG: Apply **coarse-graining** transformation in k_x-space

Entanglement measure $S_{\mathrm{EE}}(j)$ at every step $\longrightarrow$ **RG Flow of entanglement**






connected spaces
(gapless)
entanglement
between
UV and IR
UV
IR
RG flow


disconnected spaces
(gapped)
entanglement
between
UV and IR
UV
IR
RG flow

Closing Remarks

Summary of Primary Results

Main Conclusions and Takeaways

Future Work