



Quadratic Equation

14.04.2020

Total Ques : 25

MM:100

Origin

A JEE Prep Hub

1. No of the real roots of the equation $|x - 2|^{x^2 - 5x + 6} = 1$ is :
 - a. Exactly two
 - b. Exactly four
 - c. Exactly three
 - d. Can't say

2. If $|x^2 - 2x - 8| + |x^2 + x - 2| = 3|x + 2|$, then the set of all real values of x is
 - a. $[1, 4]$
 - b. $[1, 4] \cup \{-2\}$
 - c. $[-2, 1] \cup [4, \infty)$
 - d. $(-\infty, -2] \cup [1, 4]$

3. Number of values of x satisfying equations $5\{x\} = x + [x]$ and $[x] - \{x\} = \frac{1}{2}$ is :
 - a. 0
 - b. 1
 - c. 2
 - d. 3

4. Let us consider a sequence such that :

$$T_n = \left[\frac{1}{3} + \frac{n}{50} \right], \text{ here } [a] = \text{Greatest Integer Function of } a,$$
 Then calculate value of $\sum_{n=1}^{91} T_n$ is :
 - a. 55
 - b. 57
 - c. 58
 - d. 59

5. How many solutions do the given equation have :

$$\ln x = x + 1$$
 - a. Two distinct solutions
 - b. Only one solution
 - c. Two same solutions
 - d. None of Above

6. A parabola of the form $ax^2 + bx + c$ with $a > 0$ intersects the graph of $f(x) = \frac{1}{x^2 - 4}$.
Number of possible distinct solutions of the graph is
- 0
 - 2
 - 4
 - 3
7. Let $D = a^2 + b^2 + c^2$, a, b being consecutive integers and $c = ab$ then \sqrt{D} is :
- Always an even integer
 - Always odd integer
 - Sometimes an odd integer and sometimes even integer
 - Always irrational
8. The equation $x^2 + ax + b^2 = 0$ has two roots each of which exceeds a number c , then
- $a^2 < 4b^2$
 - $c^2 + ac + b^2 > 0$
 - $\frac{-a}{2} < c$
 - None
9. If the two equations $ax^2 + 2bx + c = 0$ and $ax^2 + 2cx + b = 0$ have a common root, then $a + 4b + 4c =$
- 1
 - 1
 - 2
 - 0
10. If $F(x) = x^4 + 9x^3 + 35x^2 - x + 4$ then $F(-5 + 2\sqrt{-4})$ is equal to
- 160
 - 160
 - 0
 - None
11. No. of solution of the given equation in interval $[0, 4\pi]$ satisfying the equation $6\sin^2 x - \sin x - 1 = 0$
- 2
 - 6

- c. 7
d. 8
12. The equation $a \sin x + b \cos x = c$, where $|c| > \sqrt{a^2 + b^2}$ has
a. A unique solution
b. Infinite no of solutions
c. No solution
d. None
13. For what values of p , will the equation

$$(p^2 - 16)x^2 - (p^2 + 5p + 4)x - p^3 - 4p^2 - 4p - 16 = 0$$
 Has more than two solutions ?
a. No such value exist
b. 4, -5
c. 4
d. 4, -4, -1
14. Find the minimum value of the argument
 $a^2 + b^2 + c^2 - ab - bc - ca$, where $a, b, c \in R$
a. $-\infty$
b. 1
c. ∞
d. 0
15. The value of the expression is

$$\sqrt{6 + \sqrt{6 + \sqrt{6 + \sqrt{6 + \sqrt{6 + \dots \infty}}}}}$$
 a. 1
b. 6
c. 3
d. 9
16. The minimum value of expression $|x| + |x + \frac{1}{2}| + |x - 3| + |x - \frac{5}{2}|$ is :
a. 2
b. 3
c. 6
d. 4
17. The set of values of x for which the inequality $[x]^2 - 5[x] + 6 \leq 0$ (where $[.]$ denotes the

GIF) hold good is :

- a. $2 \leq [x] < 3$
- b. $2 \leq [x] < 4$
- c. $2 \leq x < 3$
- d. $2 \leq [x] \leq 4$

18. The least value of $\frac{x^2-6x+5}{x^2+2x+1}$ is :

- a. -1
- b. $-\frac{1}{2}$
- c. $-\frac{1}{4}$
- d. $-\frac{1}{3}$

19. Suppose $a^2 = 5a - 8$ and $b^2 = 5b - 8$, then equation whose roots are $\frac{a}{b}$ and $\frac{b}{a}$ is

- a. $6x^2 - 5x + 6 = 0$
- b. $8x^2 - 9x + 8 = 0$
- c. $9x^2 - 8x + 9 = 0$
- d. $8x^2 + 9x + 8 = 0$

20. If the set containing the roots of the equation $(a-b)x^2 + (b-c)x + (c-a) = 0$ has infinite no of values then what will be the condition :

- a. $a = b \neq c$
- b. $a \neq b \neq c$
- c. $a = b = c$
- d. None

21. If $P(x) = ax^2 + bx + c$ and $Q(x) = -ax^2 + dx + c$, where $ac \neq 0$, then $P(x)Q(x) = 0$ has

- a. No real root
- b. Exactly two roots
- c. At least two distinct roots
- d. None of these

22. The sum of all the real roots of the equation $|x-2|^2 + |x-2| - 2 = 0$ is :

- a. 7
- b. 4
- c. 1

- d. None
23. If α, β are the roots of $x^2 + 3x - 2 = 0$, then $\frac{\alpha+1}{\beta} + \frac{\beta+1}{\alpha} =$
- $\frac{-15}{2}$
 - $\frac{6}{5}$
 - $\frac{-5}{2}$
 - 5
24. If one root of the equation $ax^2 + bx + c = 0$, $a \in R^+$, are two consecutive odd positive integers. Then
- $|b| \leq 4a$
 - $|b| \geq 4a$
 - $|b| \leq 2a$
 - $|b| \leq a$
25. If the roots of the equation $a(b-c)x^2 + b(c-a)x + c(a-b) = 0$ are equal then a,b,c are in :
- A.P.
 - G.P.
 - H.P.
 - None