

Confidential Medical Report

Medical Report

Generated from Uploaded Patient Documents

Date Generated: October 28, 2025

Report ID: bbffea31

This is an automatically generated report. Always verify findings with original source documents.

Confidential Medical Report

Report Contents

- Introduction
- Clinical Findings
- Patient Tables
- Graphs

pISSN 2287-2728 eISSN 2287-285X <https://doi.org/10.3350/cmh.2022.0365> Review Clinical and Molecular Hepatology 2023;29(Suppl):S32-S42 Global incidence and prevalence of nonalcoholic fatty liver disease Margaret LP Teng^{1,2,*}, Cheng Han Ng^{2,*}, Daniel Q. Huang^{1,2,3}, Kai En Chan², Darren JH Tan², Wen Hui Lim², Ju Dong Yang⁴, Eunice Tan^{1,2,3}, and Mark D. Muthiah^{1,2,3} 1Division of Gastroenterology and Hepatology, Department of Medicine, National University Hospital, Singapore; 2Yong Loo Lin School of Medicine, National University of Singapore, Singapore; 3National University Centre for Organ Transplantation, National University Hospital, Singapore; 4Karsh Division of Gastroenterology and Hepatology, Cedars-Sinai Medical Center, Los Angeles, CA, USA Nonalcoholic fatty liver disease (NAFLD) is a leading cause of liver disease worldwide. The estimated global incidence of NAFLD is 47 cases per 1,000 population and is higher among males than females. The estimated global prevalence of NAFLD among adults is 32% and is higher among males (40%) compared to females (26%). The global prevalence of NAFLD has increased over time, from 26% in studies from 2005 or earlier to 38% in studies from 2016 or beyond. The prevalence of NAFLD varies substantially by world region, contributed by differing rates of obesity, and genetic and socioeconomic factors. The prevalence of NAFLD exceeds 40% in the Americas and South-East Asia. The prevalence of NAFLD is projected to increase significantly in multiple world regions by 2030 if current trends are left unchecked. In this review, we discuss trends in the global incidence and prevalence of NAFLD and discuss future projections. (Clin Mol Hepatol 2023;29(Suppl):S32-S42) Keywords: Nonalcoholic fatty liver disease; Incidence; Prevalence; Epidemiology INTRODUCTION hepatitis (NASH), which can progress to liver fibrosis and cirrhosis.² The global prevalence of NAFLD has been increasing Nonalcoholic fatty liver disease (NAFLD) is a leading cause over time, with a recent meta-analysis estimating that 32% of of chronic liver disease globally.¹ It encompasses a spectrum the adult population is afflicted by NAFLD.³ This has occurred ranging from simple hepatic steatosis to nonalcoholic steato- in tandem with the global obesity and diabetes epidemics.^{4,5} Corresponding author : Daniel Q. Huang Division of Gastroenterology and Hepatology, Department of Medicine, National University Health System, Tower Block Level 10, 1E Kent Ridge Road, Singapore 119228, Singapore Tel: +65 6772 4354, Fax: +65 6775 1518, E-mail: daniel_huang@nus.edu.sg <https://orcid.org/0000-0002-5165-5061> Mark D. Muthiah Division of Gastroenterology and Hepatology, Department of Medicine, National University Health System, Tower Block Level 10, 1E Kent Ridge Road, Singapore 119228, Singapore Tel: +65 6772 4354, Fax: +65 6775 1518, E-mail: mdcmdm@nus.edu.sg <https://orcid.org/0000-0002-9724-4743> *Margaret LP Teng and Cheng Han Ng contributed equally as co-first authors. Editor: Sang Gyune Kim, Soonchunhyang University Hospital Bucheon, Korea Received : Nov. 2, 2022 / Revised : Dec. 6, 2022 / Accepted : Dec. 12, 2022 Copyright © 2023 by Korean Association for the Study of the Liver This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which

Confidential Medical Report
permits unrestricted non-commercial use, distribution, and reproduction in any medium,
provided the original work is properly cited.

pISSN 2287-2728 eISSN 2287-285X <https://doi.org/10.3350/cmh.2022.0365> Review Clinical and Molecular Hepatology 2023;29(Suppl):S32-S42 Global incidence and prevalence of nonalcoholic fatty liver disease Margaret LP Teng^{1,2,*}, Cheng Han Ng^{2,*}, Daniel Q. Huang^{1,2,3}, Kai En Chan², Darren JH Tan², Wen Hui Lim², Ju Dong Yang⁴, Eunice Tan^{1,2,3}, and Mark D. Muthiah^{1,2,3} ¹Division of Gastroenterology and Hepatology, Department of Medicine, National University Hospital, Singapore; ²Yong Loo Lin School of Medicine, National University of Singapore, Singapore; ³National University Centre for Organ Transplantation, National University Hospital, Singapore; ⁴Karsh Division of Gastroenterology and Hepatology, Cedars-Sinai Medical Center, Los Angeles, CA, USA Nonalcoholic fatty liver disease (NAFLD) is a leading cause of liver disease worldwide. The estimated global incidence of NAFLD is 47 cases per 1,000 population and is higher among males than females. The estimated global prevalence of NAFLD among adults is 32% and is higher among males (40%) compared to females (26%). The global prevalence of NAFLD has increased over time, from 26% in studies from 2005 or earlier to 38% in studies from 2016 or beyond. The prevalence of NAFLD varies substantially by world region, contributed by differing rates of obesity, and genetic and socioeconomic factors. The prevalence of NAFLD exceeds 40% in the Americas and South-East Asia. The prevalence of NAFLD is projected to increase significantly in multiple world regions by 2030 if current trends are left unchecked. In this review, we discuss trends in the global incidence and prevalence of NAFLD and discuss future projections. (Clin Mol Hepatol 2023;29(Suppl):S32-S42) Keywords: Nonalcoholic fatty liver disease; Incidence; Prevalence; Epidemiology INTRODUCTION hepatitis (NASH), which can progress to liver fibrosis and cirrhosis.² The global prevalence of NAFLD has been increasing Nonalcoholic fatty liver disease (NAFLD) is a leading cause over time, with a recent meta-analysis estimating that 32% of of chronic liver disease globally.¹ It encompasses a spectrum the adult population is afflicted by NAFLD.³ This has occurred ranging from simple hepatic steatosis to nonalcoholic steato- in tandem with the global obesity and diabetes epidemics.^{4,5} Corresponding author : Daniel Q. Huang Division of Gastroenterology and Hepatology, Department of Medicine, National University Health System, Tower Block Level 10, 1E Kent Ridge Road, Singapore 119228, Singapore Tel: +65 6772 4354, Fax: +65 6775 1518, E-mail: daniel_huang@nus.edu.sg <https://orcid.org/0000-0002-5165-5061> Mark D. Muthiah Division of Gastroenterology and Hepatology, Department of Medicine, National University Health System, Tower Block Level 10, 1E Kent Ridge Road, Singapore 119228, Singapore Tel: +65 6772 4354, Fax: +65 6775 1518, E-mail: mdcmdm@nus.edu.sg <https://orcid.org/0000-0002-9724-4743> *Margaret LP Teng and Cheng Han Ng contributed equally as co-first authors. Editor: Sang Gyune Kim, Soonchunhyang University Hospital Bucheon, Korea Received : Nov. 2, 2022 / Revised : Dec. 6, 2022 / Accepted : Dec. 12, 2022 Copyright © 2023 by Korean Association for the Study of the Liver This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which

permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. Clinical and Molecular Hepatology Volume_29 Supplement February 2023 also pooled data from 245 studies (2,699,627 individuals) and estimated the global prevalence of NAFLD at 29.8%, which is consistent with Riazi's findings. Likewise, in this study, there was limited or no data from Africa, Oceania, and North and South America. Asia The prevalence of NAFLD varies substantially by region (Fig. 2). The overall prevalence of NAFLD in Asia is approximately 30%. A meta-analysis by Le et al.¹² conducted a literature search in 2019 (182 studies with 2,385,999 individuals) and estimated NAFLD prevalence in Asia at 30.5%. A recent meta- analysis by Riazi et al.³ which included 63 studies (1,000,681 individuals) found that NAFLD prevalence in Asia was 31.6%. This is consistent with a previous meta-analysis by Li et al.¹⁰ which reported NAFLD prevalence in Asia to be 29.62%. The prevalence of NAFLD within Asia is highly variable as it encompasses countries with a wide spectrum of ethnicities and socioeconomic factors. Among Asian subregions, South- east Asia had the highest NAFLD prevalence of 42%. Among Asian countries with more than 3 studies available, Li et al.¹⁰ determined that the highest pooled NAFLD prevalence was in Iran (38.07%), and the lowest pooled NAFLD prevalence was in Japan (22.28%). Riazi et al.³ determined similar results and found that Iran had the highest NAFLD prevalence (40.8%), followed by Taiwan (36.1%), South Korea (34.6%), and China (32.5%). On the other hand, Japan had a strikingly low NAFLD prevalence of 22.3%, which may be related to a low prevalence of obesity. In China, a meta-analysis by Wu et al.¹³ estimated a NAFLD prevalence of 29.88%, and another study by Zhou et al.¹⁴ estimated that NAFLD prevalence was 29.2%. NAFLD prevalence in South Korea is also approximately 30%—a meta-analysis by Im et al.¹⁵ reported a NAFLD prevalence of 30.3%, and Li et al.¹⁰ reported a similar prevalence of 32.9%.¹¹ A large cross- sectional study of 571,872 Korean males in their early 20s found that even among young adult males, NAFLD prevalence was 13.47%, with an increase from 10.66% in 2015 to 16.44% in 2021. There was a higher prevalence of metabolic risk factors such as hypertension, hypercholesterolemia, and hyperglycemia during the same period.¹⁶ Another study utilizing data from Korea National Health and Nutrition Examination Survey found that NAFLD prevalence increased from 18.6% in 1998–2001 to 21.5% in 2016–2017, and there was a S34

<https://doi.org/10.3350/cmh.2022.0365> <http://www.e-cmh.org> esaesid revil yttaf cilohoclanon fo ecnedicni eht rof atad gnidivorp sesylana-atem detceleS .1 elbaT stnemmoC sgnidnfi niaM dedulcni seirtnuoC/snoigeR sraey yduts slaudividni fo .oN seiduts fo .oN yduts dedulcni fo ytirojam ehT 000,1 rep sesac 9.64 ecnedicni ,aeroK htuoS ,napaJ ,anihC(aisA 8102–4991 567,183 61)2202(3.la te izaiR aisA morf erew seiduts ecnedicni eht ;sraey-nosrep)learsl ,gnoK gnoH eb ton yam atad ecneh 000,1 rep sesac 8.07(nem ni elbazilareneg rehgiH saw)sraey-nosrep ytiengoreteh laitnatsbuS sesac 6.92(nemow ni naht)sraey-nosrep 000,1 rep aisA morf erew seiduts IIA 000,1 rep sesac 9.05 ecnedicni ,aeroK htuoS ,napaJ ,anihC(aisA 7102–2002 889,614 81)9102(01.la te iL eb ton yam atad ecneh ecnedicni ;sraey-nosrep)gnoK gnoH elbazilareneg rep 36(anihC ni tsehgiH tsewol ,)sraey-nosrep 000,1 000,1 rep 92(napaJ ni)sraey-nosrep dedulcni fo rebmun detimil A rep sesac 43.25 ecnedicni)learsl ,napaJ ,anihC(aisA 3102–7991 598,4 5)6102(9.la te issonuoY seiduts anihC(sraey-nosrep 000,1

SA 3 morr erew seiduts IIA rep sesac 10.82 ,)napaJ dna seirtnuoc)learsI(sraey-nosrep
 000,1 Clinical and Molecular Hepatology Volume_29 Supplement February 2023 higher
 prevalence of obesity and diabetes over the same pe- riod.¹⁷ These suggest that the
 increasing NAFLD prevalence may be driven by an increase in metabolic risk factors. Ito et
 al.¹⁸ reported a comparatively lower NAFLD prevalence of 25.5% in Japan, in line with the
 findings by Li and Riazi. This could be attributed to a lower prevalence of obesity and dia-
 betes in Japan compared to other countries,^{19,20} and may be related to a diet that is
 traditionally lower in fat and red meat.²¹ In South Asia, India had a NAFLD prevalence of
 25.7– 32.74%, Bangladesh had a NAFLD prevalence of 26.2–33.86%, and Sri Lanka had a
 NAFLD prevalence of 24.74%.^{3,10} In South- east Asia, Li et al.¹⁰ reported that NAFLD
 prevalence was 38.5% in Malaysia, 40.43% in Singapore, and 51.04% in Indo- nesia. Data
 from Central Asia is lacking, but the Global Bur- den of Disease Study (GBD) 2019 reported
 NAFLD prevalence in Central Asia increased from 12.4% in 1990 to 19.7% in 2019, although
 these estimates require cautious interpretation as the Global Burden of Disease Study relied
 on complex mod- eling and past trends when data was limited.²² A distinct feature of the
 NAFLD epidemic in Asia is the high prevalence of lean NAFLD (body mass index [BMI] <23)
 and non-obese NAFLD (BMI <25).²³ Up to 19% of non-obese Asians have NAFLD,²⁴ which
 may be contributed to a higher percentage of visceral adiposity in Asians compared to other
 ethnicities.²⁵ Visceral adiposity plays an important role in ath- erogenic dyslipidemia and
 insulin resistance. It is a major risk factor for type 2 diabetes and has been implicated in the de-
 velopment and progression of NAFLD.²⁶ Asians also tend to develop diabetes at a
 younger age and lower BMI level, re- sulting in a longer duration of disease and increased
 likeli- hood of complications.^{27–29} Worryingly, emerging data sug- gest that individuals with
 lean NAFLD may be at a higher risk of progressive liver disease, but this hypothesis requires
 vali- dation.³⁰ Europe Meta-analyses by Le and Riazi had similar estimates of the prevalence
 of NAFLD in Europe at 30.9% (11 studies with 15,062 individuals)¹² and 32.6% (7 studies
 with 14,111 individ- uals),³ respectively. Another meta-analysis by Cholongitas et al.³¹
 pooled data from 17 studies (85,203 individuals) and esti- mated NAFLD prevalence in
 Europe to be 26.9%. Cholongitas also found that NAFLD prevalence in Mediterranean coun-
 S36 <https://doi.org/10.3350/cmh.2022.0365> <http://www.e-cmh.org> deunitnoC .2 elbaT
 stnemmoC sgnidnfi niaM dedulcni seirtnuoC/snoigeR sraey yduts slauidivdni fo .oN seiduts fo
 .oN yduts ot ylno elbacilppa era ataD ecnelaverp llarevo detamitsE aisA 7102–4991
 815,440,31 732)9102(01.la te iL aisA %6.92 emit revo desaercni ecnelaverP %5.82
 ,)5002–5991(%3.52 – –2102(%9.33 ,)1102–6002()7102 .esaesid revil yttaf cilohoclanon
 ,DLFAN Clinical and Molecular Hepatology Volume_29 Supplement February 2023 tries at
 23.9% compared to non-Mediterranean countries at included in this meta-analysis were
 hospital-based studies 28.5%, although the difference was not statistically signifi- and
 included patients with risk factors for NAFLD, hence the cant. results may not have been fully
 representative of the general Within Europe, Turkey had the highest NAFLD prevalence
 population. Le et al.¹² pooled data from 3 studies (5,716 indi- at 48.4%, followed by Italy at
 38.2%. Germany, Portugal, and viduals) and determined that South America had the greatest
 Spain had NAFLD prevalence between 25–27%.³ A cross-sec- estimated NAFLD prevalence

among the continents at 35.7%. tional study utilizing data from a large population-based co- This may be due to a combination of genetic susceptibility hort in France found that NAFLD prevalence in France was and a greater prevalence of metabolic risk factors.⁴⁴ There is a 18.2%.³² A study involving individuals from population-based high prevalence of PNPLA3 genetic polymorphism in the studies in Russia reported that NAFLD prevalence was 40% in general population, especially among individuals with Native the Ural Eye and Medical Study (UEMS) (5,852 individuals), American ancestry.⁴⁵⁻⁴⁷ Furthermore, obesity is extremely and 69.8% in the Ural Very Old Study (UVOS) (1,130 individu- common in the region—a cross-sectional study across 4 als).³³ However, it should be noted that in the UVOS, individu- geographical regions found that central obesity was highest als were older with minimum age of 85 years, and methods in South America.⁴⁸ Type 2 diabetes has also been rising in for diagnosis of NAFLD differed between studies as well. prevalence in South America.⁴⁹ Data from the meta-analysis by Le showed that compared to other regions, NAFLD indi- North America viduals in South America had a higher likelihood of having diabetes and higher mean cholesterol levels.¹² In addition, Based on subgroup data from 4 studies (18,356 individuals), physical activity is often inadequate—Latin America was Le et al.¹² estimated that the prevalence of NAFLD in North ranked as the top region for physical inactivity, with a third of America was 35.3%. More recently, NAFLD prevalence was the population experiencing a lack of physical activity.⁵⁰ reported at 47.8% in the meta-analysis by Riazi which includ- ed 2 large studies with 15,178 individuals from the USA.³ This Africa is driven by a high prevalence of obesity in the USA. In North America, Hispanics have the highest NAFLD prevalence, fol- There is a paucity of data from Africa on the epidemiology lowed by non-Hispanic Whites and non-Hispanic blacks.³⁴⁻³⁹ of NAFLD. A meta-analysis estimated the prevalence of Based on data from National Health and Nutrition Examina- NAFLD in Africa at 13.5%, ranging from 9% in Nigeria to 20% tion Surveys (NHANES) 2017–2018, NAFLD prevalence was es- in Sudan.⁵¹ More recently, NAFLD prevalence was reported at timated at 63.7% in Hispanics, 56.8% in non-Hispanic whites, 28.2% in the meta-analysis by Le, and 56.8% in the meta- and 46.2% in non-Hispanic blacks.³⁷ This could be attributed analysis by Riazi.^{3,12} Of note, the meta-analysis by Riazi only to genetic factors like the patatin-like phospholipase do- included 1 study from Egypt. The wide variation in estimates main-containing protein 3 (PNPLA3) mutation, which is asso- of NAFLD prevalence is likely related to a lack of reliable data ciated with elevated risk for hepatic steatosis and NASH, oc- from Africa.^{52,53} ccurring more frequently in Hispanics.^{40,41} This could also be related to metabolic factors like the higher prevalence of Oceania central adiposity and insulin resistance in Hispanics com- pared to non-Hispanic whites.^{34,42} Lower serum triglyceride Likewise, there is scarce data from Oceania on the inci- levels in African-Americans may also contribute to reduced dence and prevalence of NAFLD. Population-based studies NAFLD prevalence.³⁸ using fatty liver index have demonstrated NAFLD prevalence of 35.7–38% in Australia.^{54,55} There are no population-based South America studies on NAFLD prevalence using imaging modalities such as ultrasound.⁵⁶ A meta-analysis comprising 19 studies (5,626 individuals) by Rojas et al. estimated the prevalence of NAFLD in South America at up to 59%.⁴³ Notably, the majority of the studies S38 <https://doi.org/10.3350/cmh.2022.0365> <http://www.e-cmh.org>

Clinical and Molecular Hepatology Volume_29 Supplement February 2023 REFERENCES

ogy of NAFLD in Mainland China with analysis by adjusted gross regional domestic product: a meta-analysis. *Hepatol Int* 1. Cheemerla S, Balakrishnan M. Global epidemiology of chronic liver disease. *Clin Liver Dis (Hoboken)* 2021;17:365-370. 14. Zhou F, Zhou J, Wang W, Zhang XJ, Ji YX, Zhang P, et al. Unex- 2. Chalasani N, Younossi Z, Lavine JE, Charlton M, Cusi K, Rinella M, pected rapid increase in the burden of NAFLD in China from et al. The diagnosis and management of nonalcoholic fatty liver 2008 to 2018: a systematic review and meta-analysis. *Hepatol- disease: practice guidance from the American Association for ogy* 2019;70:1119-1133. the Study of Liver Diseases. *Hepatology* 2018;67:328-357. 15. Im HJ, Ahn YC, Wang JH, Lee MM, Son CG. Systematic review 3. Riazi K, Azhari H, Charette JH, Underwood FE, King JA, Afshar on the prevalence of nonalcoholic fatty liver disease in South EE, et al. The prevalence and incidence of NAFLD worldwide: Korea. *Clin Res Hepatol Gastroenterol* 2021;45:101526. a systematic review and meta-analysis. *Lancet Gastroenterol* 16. Lee J, Kim T, Yang H, Bae SH. Prevalence trends of non-alcoholic *Hepatol* 2022;7:851-861. fatty liver disease among young men in Korea: a Korean mili- 4. GBD 2015 Obesity Collaborators. Health effects of overweight tary population-based cross-sectional study. *Clin Mol Hepatol and obesity in 195 countries over 25 years. N Engl J Med* 2022;28:196-206. 2017;377:13-27. 17. Park SH, Plank LD, Suk KT, Park YE, Lee J, Choi JH, et al. Trends 5. Stefan N, Cusi K. A global view of the interplay between non- in the prevalence of chronic liver disease in the Korean adult alcoholic fatty liver disease and diabetes. *Lancet Diabetes En- population, 1998-2017. Clin Mol Hepatol* 2020;26:209-215. *ocrinol* 2022;10:284-296. 18. Ito T, Ishigami M, Zou B, Tanaka T, Takahashi H, Kurosaki M, et al. 6. Tan DJH, Setiawan VW, Ng CH, Lim WH, Muthiah MD, Tan EX, et The epidemiology of NAFLD and lean NAFLD in Japan: a meta- al. Global burden of liver cancer in males and females: chang- analysis with individual and forecasting analysis, 1995-2040. ing etiological basis and the growing contribution of NASH. *Hepatol Int* 2021;15:366-379. *Hepatology* 2022 Aug 29. doi: 10.1002/hep.32758. 19. Organisation for Economic Co-operation and Development 7. Huang DQ, Singal AG, Kono Y, Tan DJ, El-Serag HB, Loomba R. (OECD). Obesity update 2017. OECD web site, <<https://www.oecd.org/health/health-systems/Obesity-Update-2017.pdf>>. 2019: NASH is the fastest growing cause of liver cancer. *Cell* Accessed 22 Oct 2022. *Metab* 2022;34:969-977.e2. 20. International Diabetes Federation (IDF). IDF Diabetes Atlas 8. Younossi ZM, Stepanova M, Ong J, Trimble G, AlQahtani S, You- Ninth edition 2019. IDF web site, <<https://www.diabetesatlas.org/>>. 2019: NASH is the most rapidly increasing indication for liver transplantation in the United States. *Clin Gastroenterol Hepatol* 2021;19:580-589.e5. 21. Ikeda N, Saito E, Kondo N, Inoue M, Ikeda S, Satoh T, et al. 9. Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. What has made the population of Japan healthy? *Lancet Global epidemiology of nonalcoholic fatty liver disease-meta- 2011;378:1094-1105. analytic assessment of prevalence, incidence, and outcomes. 22. Global Burden of Disease Collaborative Network. Global Burden Hepatology* 2016;64:73-84. of Disease Study 2019 (GBD 2019) results. *Global Health Data* 10. Li J, Zou

B, Yeo YH, Feng Y, Xie X, Lee DH, et al. Prevalence, Exchange web site, <<http://ghdx.healthdata.org/gbd-results>> incidence, and outcome of non-alcoholic fatty liver disease in tool>. Accessed 22 Oct 2022. Asia, 1999-2019: a systematic review and meta-analysis. *Lancet* 23. Yoo JJ, Kim W, Kim MY, Jun DW, Kim SG, Yeon JE, et al. Recent Gastroenterol Hepatol 2019;4:389-398. research trends and updates on nonalcoholic fatty liver disease. 11. Kang SH, Lee HW, Yoo JJ, Cho Y, Kim SU, Lee TH, et al. KASL clini- Clin Mol Hepatol 2019;25:1-11. cal practice guidelines: management of nonalcoholic fatty liver 24. Fan JG, Kim SU, Wong VW. New trends on obesity and NAFLD in disease. Clin Mol Hepatol 2021;27:363-401. Asia. J Hepatol 2017;67:862-873. 12. Le MH, Yeo YH, Li X, Li J, Zou B, Wu Y, et al. 2019 Global NAFLD 25. Nazare JA, Smith JD, Borel AL, Haffner SM, Balkau B, Ross R, et al. prevalence: a systematic review and meta-analysis. Clin Gastro- Ethnic influences on the relations between abdominal subcuta- enterol Hepatol 2022;20:2809-2817.e28. neous and visceral adiposity, liver fat, and cardiometabolic risk 13. Wu Y, Zheng Q, Zou B, Yeo YH, Li X, Li J, et al. The epidemiol- profile: the International Study of Prediction of Intra-Abdominal S40 <https://doi.org/10.3350/cmh.2022.0365> <http://www.e-cmh.org> Margaret LP Teng, et al. Global incidence and prevalence of NAFLD Adiposity and Its Relationship With Cardiometabolic Risk/Intra- transient elastography among US adults: results from NHANES Abdominal Adiposity. Am J Clin Nutr 2012;96:714-726. 2017-2018. PLoS One 2021;16:e0252164. 26. Hanlon CL, Yuan L. Nonalcoholic fatty liver disease: the role of 38. Sherif ZA, Saeed A, Ghavimi S, Nouraie SM, Laiyemo AO, Brim visceral adipose tissue. Clin Liver Dis (Hoboken) 2022;19:106- H, et al. Global epidemiology of nonalcoholic fatty liver dis- 110. ease and perspectives on US minority populations. Dig Dis Sci 27. Yoon KH, Lee JH, Kim JW, Cho JH, Choi YH, Ko SH, et al. Epidemic 2016;61:1214-1225. obesity and type 2 diabetes in Asia. Lancet 2006;368:1681-1688. 39. Arshad T, Golabi P, Henry L, Younossi ZM. Epidemiology of non- 28. Huxley R, James WP, Barzi F, Patel JV, Lear SA, Suriyawongpaisal alcoholic fatty liver disease in North America. Curr Pharm Des P, et al. Ethnic comparisons of the cross-sectional relationships 2020;26:993-997. between measures of body size with diabetes and hyperten- 40. Romeo S, Kozlitina J, Xing C, Pertsemlidis A, Cox D, Pennacchio sion. Obes Rev 2008;9 Suppl 1:53-61. LA, et al. Genetic variation in PNPLA3 confers susceptibility to 29. Chan JC, Malik V, Jia W, Kadowaki T, Yajnik CS, Yoon KH, et al. nonalcoholic fatty liver disease. Nat Genet 2008;40:1461-1465. Diabetes in Asia: epidemiology, risk factors, and pathophysiol- 41. Wagenknecht LE, Palmer ND, Bowden DW, Rotter JI, Norris JM, ogy. JAMA 2009;301:2129-2140. Ziegler J, et al. Association of PNPLA3 with non-alcoholic fatty 30. Hagström H, Nasr P, Ekstedt M, Hammar U, Ståhl P, Hultcrantz R, liver disease in a minority cohort: the Insulin Resistance Athero- et al. Risk for development of severe liver disease in lean pa- sclerosis Family Study. Liver Int 2011;31:412-416. tients with nonalcoholic fatty liver disease: a long-term follow- 42. Guerrero R, Vega GL, Grundy SM, Browning JD. Ethnic differ- up study. Hepatol Commun 2018;2:48-57. ences in hepatic steatosis: an insulin resistance paradox? Hepa- 31. Cholongitas E, Pavlopoulou I, Papatheodoridi M, Markakis GE, tology 2009;49:791-801. Bouras E, Haidich AB, et al. Epidemiology of nonalcoholic fatty 43. Rojas YAO, Cuellar CLV, Barrón KMA, Arab JP, Miranda AL. Non- liver disease in Europe: a systematic review and meta-analysis. alcoholic

fatty liver disease prevalence in Latin America: a systematic review and meta-analysis. *Ann Gastroenterol* 2021;34:404-414.

32. Nabi O, Lacombe K, Boursier J, Mathurin P, Zins M, Serfaty L. 44. Pinto Marques Souza de Oliveira C, Pinchemel Cotrim H, Arrese Prevalence and risk factors of nonalcoholic fatty liver disease M.

Nonalcoholic fatty liver disease risk factors in Latin American and advanced fibrosis in general population: the French Na- populations: current scenario and perspectives. *Clin Liver Dis*

tionwide NASH-CO Study. *Gastroenterology* 2020;159:791-793. (Hoboken) 2019;13:39-42.

e2. 45. Arrese M, Arab JP, Riquelme A, Benítez CE, Barrera F, Soza A, 33. Bikbov MM, Gilmanshin TR, Zainullin RM, Kazakbaeva GM, Iaku- et al. High prevalence of PNPLA3 rs738409 (I148M) polymor- pova EM, Fakhretdinova AA, et al. Prevalence of non-alcoholic phism in Chilean latino population and its association to non- fatty liver disease in the Russian Ural Eye and Medical Study alcoholic fatty liver disease risk and histological disease sever- and the Ural Very Old Study. *Sci Rep* 2022;12:7842. ity. *Hepatology* 2015;62 Suppl 1:1285A.

34. Browning JD, Szczepaniak LS, Dobbins R, Nuremberg P, Horton 46. Pontoriero AC, Trinks J, Hulaniuk ML, Caputo M, Fortuny L, Pratx JD, Cohen JC, et al. Prevalence of hepatic steatosis in an urban LB, et al. Influence of ethnicity on the distribution of genetic population in the United States: impact of ethnicity. *Hepatol- polymorphisms associated with risk of chronic liver disease in ogy* 2004;40:1387-1395. South American populations. *BMC Genet* 2015;16:93. 35. Lazo M, Hernaez R, Eberhardt MS, Bonekamp S, Kamel I, Gual- 47. Chinchilla-López P, Ramírez-Pérez O, Cruz-Ramón V, Canizales- lar E, et al. Prevalence of nonalcoholic fatty liver disease in the Quinteros S, Domínguez-López A, Ponciano-Rodríguez G, et al. United States: the Third National Health and Nutrition Examina- More evidence for the genetic susceptibility of Mexican popu- tion Survey, 1988-1994. *Am J Epidemiol* 2013;178:38-45. lation to nonalcoholic fatty liver disease through PNPLA3. *Ann* 36. Kim D, Kim W, Adejumo AC, Cholanteril G, Tighe SP, Wong RJ, *Hepatol* 2018;17:250-255. et al. Race/ethnicity-based temporal changes in prevalence of 48. Patel SA, Ali MK, Alam D, Yan LL, Levitt NS, Bernabe-Ortiz A, et NAFLD-related advanced fibrosis in the United States, 2005- al. Obesity and its relation with diabetes and hypertension: a 2016. *Hepatol Int* 2019;13:205-213. cross-sectional study across 4 geographical regions. *Glob Heart* 37. Zhang X, Heredia NI, Balakrishnan M, Thrift AP. Prevalence and 2016;11:71-79.e4. factors associated with NAFLD detected by vibration controlled 49. Gallardo-Rincón H, Cantoral A, Arrieta A, Espinal C, Magnus MH, S41 <http://www.e-cmh.org> <https://doi.org/10.3350/cmh.2022.0365> Clinical and Molecular Hepatology Volume_29 Supplement February 2023 Palacios C, et al. Review: type 2 diabetes in Latin America and disease using hierarchical bayesian approach. *Clin Mol Hepatol the Caribbean: regional and country comparison on prevalence*, 2022;28:841-850. trends, costs and expanded prevention. *Prim Care Diabetes* 58. Estes C, Anstee QM, Arias-Loste MT, Bantel H, Bellentani S, 2021;15:352-359. Caballeria J, et al. Modeling NAFLD disease burden in China, 50. Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends France, Germany, Italy, Japan, Spain, United Kingdom, and in insufficient physical activity from 2001 to 2016: a pooled United States for the period 2016-2030. *J Hepatol* 2018;69:896- analysis of 358 population-based surveys with 1.9 million par- 904. ticipants. *Lancet Glob Health* 2018;6:e1077-e1086. Erratum

in: 59. Estes C, Chan HL, Chien RN, Chuang WL, Fung J, Goh GB, Lancet Glob Health 2019;7:e36. et al. Modelling NAFLD disease burden in four Asian re- 51. Younossi Z, Anstee QM, Marietti M, Hardy T, Henry L, Eslam M, gions-2019-2030. Aliment Pharmacol Ther 2020;51:801-811. et al. Global burden of NAFLD and NASH: trends, predictions, 60. Alswat K, Aljumah AA, Sanai FM, Abaalkhail F, Alghamdi M, Al risk factors and prevention. Nat Rev Gastroenterol Hepatol Hamoudi WK, et al. Nonalcoholic fatty liver disease burden - 2018;15:11-20. Saudi Arabia and United Arab Emirates, 2017-2030. Saudi J Gas- 52. Spearman CW, Afihene M, Betiku O, Bobat B, Cunha L, Kassian- troenterol 2018;24:211-219. Erratum in: Saudi J Gastroenterol ides C, et al. Epidemiology, risk factors, social determinants of 2018;24:255. health, and current management for non-alcoholic fatty liver 61. Estes C, Razavi H, Loomba R, Younossi Z, Sanyal AJ. Modeling disease in sub-Saharan Africa. Lancet Gastroenterol Hepatol the epidemic of nonalcoholic fatty liver disease demonstrates 2021;6:1036-1046. an exponential increase in burden of disease. Hepatology 53. Ge X, Zheng L, Wang M, Du Y, Jiang J. Prevalence trends in non- 2018;67:123-133. alcoholic fatty liver disease at the global, regional and national 62. Swain MG, Ramji A, Patel K, Sebastiani G, Shaheen AA, Tam E, et levels, 1990-2017: a population-based observational study. BMJ al. Burden of nonalcoholic fatty liver disease in Canada, 2019- Open 2020;10:e036663. 2030: a modelling study. CMAJ Open 2020;8:E429-E436. 54. Roberts SK, Majeed A, Glenister K, Magliano D, Lubel JS, Bourke 63. Adams LA, Roberts SK, Strasser SI, Mahady SE, Powell E, Estes C, L, et al. Prevalence of non-alcoholic fatty liver disease in region- et al. Nonalcoholic fatty liver disease burden: Australia, 2019- al Victoria: a prospective population-based study. Med J Aust 2030. J Gastroenterol Hepatol 2020;35:1628-1635. 2021;215:77-82. 64. Lazarus JV, Mark HE, Anstee QM, Arab JP, Batterham RL, Cas- 55. Farrell AM, Magliano DJ, Shaw JE, Thompson AJ, Croagh C, Ryan tera L, et al. Advancing the global public health agenda for MC, et al. A problem of proportions: estimates of metabolic as- NAFLD: a consensus statement. Nat Rev Gastroenterol Hepatol sociated fatty liver disease and liver fibrosis in Australian adults 2022;19:60-78. in the nationwide 2012 AusDiab Study. Sci Rep 2022;12:1956. 65. Karlsen TH, Sheron N, Zelber-Sagi S, Carrieri P, Dusheiko G, Bu- 56. Mahady SE, Adams LA. Burden of non-alcoholic fatty liver dis- gianesi E, et al. The EASL-Lancet Liver Commission: protecting ease in Australia. J Gastroenterol Hepatol 2018;33 Suppl 1:1-11. the next generation of Europeans against liver disease compli- 57. Le MH, Yeo YH, Zou B, Barnet S, Henry L, Cheung R, et al. cations and premature mortality. Lancet 2022;399:61-116. Forecasted 2040 global prevalence of nonalcoholic fatty liver S42 <https://doi.org/10.3350/cmh.2022.0365> <http://www.e-cmh.org>

The following tables have been extracted from the source documents.

Tables from: Introduction

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS
slaudividni fo .oN
seiduts fo .oN
ydutS

aisA eb morf ton yam erew elbazilareneg atad seiduts ecneh IIA
000,1 tsewol ecnedicni rep rep 000,1 36(,)sraey-nosrep sesac anihC rep ;sraey-nosrep)sraey-nosrep 9.05 92(ni ecnedicni tsehgih napaJ 000,1 ni

aisA eb morf ton yam erew elbazilareneg atad seiduts ecneh IIA
,aeroK htuoS ,napaJ)gnoK ,anihC(gnoH aisA
7102–2002
889,614
81
)9102(01.la te iL

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS
slaudividni fo .oN
seiduts fo .oN
ydutS

,aciremA htroN morf htuoS atad acirfA ,aciremA detimiL dna
9102 ecnelaverp htuoS htroN morf ni %3.73)%7.0 ni dna desaercni tsehgih labolg)%7.53()%3.53(ot esaercni 1991 ecnelaverP ecnelaverP detamitsE aciremA aciremA ni %8.92 %9.12 yltraey(
11(3(4()seiduts eporuE aciremA aciremA 2(,)seiduts htroN htuoS acirfA ,)seiduts ,)seiduts ,)seiduts 281(aisA
8102–1991
452,993,5
542
)1202(21.la te eL

htuoS)849,2(dna gnitisiv ot ot seitilicaf ylno ytienegoreteh elbitpecsus noitroporp stneitap saib elbacilppa erachtlaeh noitceles aciremA egral erew ecneh hgiH ataD A
%42 ecnelaverp lareneg noitalupop ni llarevo ecnelaverp 'evitpac' detamitsE ;%95 dna
)ureP ,lizarB(,anitnegrA ylno aciremA ,elihC ,ocixeM htuoS
526,5
91
)2202(34.la te sajoR

stnemmoC	ot ylno elbacilppa era aisA ataD
sgnidnfi niaM	emit ecnelaverp %5.82 –2102(revo ,)5002–5991(desaercni %9.33 llarevo ,)1102–6002(ecnelaverP detamitsE %3.52 %6.92)7102 –
dedulcni seirtnuoC/snoigeR	aisA
sraey ydutS	7102–4991
slaudividni fo .oN	815,440,31
seiduts fo .oN	732
ydutS	)9102(01.la te iL

Tables from: Clinical Findings

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS

stnemmoC
slaudividni fo .oN
seiduts fo .oN
ydutS

aisA eb morf ton yam erew elbazilareneg atad seiduts ecneh IIA
000,1 tsewol ecnedicni rep rep 000,1 36(,)sraey-nosrep sesac anihC rep ;sraey-nosrep)sraey-nosrep 9.05 92(ni ecnedicnl tsehgiH napaJ 000,1 ni
,aeroK htuoS ,napaJ)gnoK ,anihC(gnoH aisA
7102–2002
889,614
81
)9102(01.la te iL

Confidential Medical Report

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS
slaudividni fo .oN
seiduts fo .oN
ydutS

,aciremA htroN morf htuoS atad acirfA ,aciremA detimiL dna
9102 ecnelaverp htuoS htroN morf ni %3.73)%7.0 ni dna desaercni tsehgiH labolg)%7.53()%3.53(ot esaercni 1991 ecnelaverP ecnelaverP detamitsE aciremA aciremA ni %8.92 %9.12 ylaey(

,aciremA htroN morf htuoS atad acirfA ,aciremA detimiL dna
11(3(4()seiduts eporuE aciremA aciremA 2(,)seiduts htroN htuoS acirfA ,)seiduts ,)seiduts ,)seiduts 281(aisA
8102–1991
452,993,5
542
)1202(21.la te eL

htuoS)849,2(dna gnitisiv ot ot seitilicaf ylno ytienegoreteh elbitpecsus noitroporp stneitap saib elbacilppa erachtlaeh noitceles aciremA egral erew ecneh hgiH ataD A
%42 ecnelaverp lareneg noitalupop ni llarevo ecnelaverp 'evitpac' detamitsE ;%95 dna
)ureP ,lizarB(,anitnegrA ylno aciremA ,elihC ,ocixeM htuoS
526,5
91
)2202(34.la te sajoR

stnemmoC	ot ylno elbacilppa era aisA ataD
sgnidnfi niaM	emit ecnelaverp %5.82 –2102(revo ,)5002–5991(desaercni %9.33 llarevo ,)1102–6002(ecnelaverP detamitsE %3.52 %6.92)7102 –
dedulcni seirtnuoC/snoigeR	aisA
sraey ydutS	7102–4991
slaudividni fo .oN	815,440,31
seiduts fo .oN	732
ydutS	)9102(01.la te iL

Tables from: Patient Tables

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS

stnemmoC
slaudividni fo .oN
seiduts fo .oN
ydutS

aisA eb morf ton yam erew elbazilareneg atad seiduts ecneh IIA
000,1 tsewol ecnedicni rep rep 000,1 36(,)sraey-nosrep sesac anihC rep ;sraey-nosrep)sraey-nosrep 9.05 92(ni ecnedicnl tsehgiH napaJ 000,1 ni
,aeroK htuoS ,napaJ)gnoK ,anihC(gnoH aisA
7102–2002
889,614
81
)9102(01.la te iL

Confidential Medical Report

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS
slaudividni fo .oN
seiduts fo .oN
ydutS

,aciremA htroN morf htuoS atad acirfA ,aciremA detimiL dna
9102 ecnelaverp htuoS htroN morf ni %3.73)%7.0 ni dna desaercni tsehgiH labolg)%7.53()%3.53(ot esaercni 1991 ecnelaverP ecnelaverP detamitsE aciremA aciremA ni %8.92 %9.12 ylaey(

,aciremA htroN morf htuoS atad acirfA ,aciremA detimiL dna
11(3(4()seiduts eporuE aciremA aciremA 2(,)seiduts htroN htuoS acirfA ,)seiduts ,)seiduts ,)seiduts 281(aisA
8102–1991
452,993,5
542
)1202(21.la te eL

htuoS)849,2(dna gnitisiv ot ot seitilicaf ylnO ytienegoreteh elbitpecsus noitroporp stneitap saib elbacilppa erachtlaeh noitceles aciremA egral erew ecneh hgiH ataD A
%42 ecnelaverp lareneg noitalupop ni llarevo ecnelaverp 'evitpac' detamitsE ;%95 dna
)ureP ,lizarB(,anitnegrA ylnO aciremA ,elihC ,ocixeM htuoS
526,5
91
)2202(34.la te sajoR

stnemmoC	ot ylno elbacilppa era aisA ataD
sgnidnfi niaM	emit ecnelaverp %5.82 –2102(revo ,)5002–5991(desaercni %9.33 llarevo ,)1102–6002(ecnelaverP detamitsE %3.52 %6.92)7102 –
dedulcni seirtnuoC/snoigeR	aisA
sraey ydutS	7102–4991
slaudividni fo .oN	815,440,31
seiduts fo .oN	732
ydutS	)9102(01.la te iL

Tables from: Graphs

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS

stnemmoC
slaudividni fo .oN
seiduts fo .oN
ydutS

aisA eb morf ton yam erew elbazilareneg atad seiduts ecneh IIA
000,1 tsewol ecnedicni rep rep 000,1 36(,)sraey-nosrep sesac anihC rep ;sraey-nosrep)sraey-nosrep 9.05 92(ni ecnedicnl tsehgiH napaJ 000,1 ni
,aeroK htuoS ,napaJ)gnoK ,anihC(gnoH aisA
7102–2002
889,614
81
)9102(01.la te iL

Confidential Medical Report

stnemmoC
sgnidnfi niaM
dedulcni seirtnuoC/snoigeR
sraey ydutS
slaudividni fo .oN
seiduts fo .oN
ydutS

,aciremA htroN morf htuoS atad acirfA ,aciremA detimiL dna
9102 ecnelaverp htuoS htroN morf ni %3.73)%7.0 ni dna desaercni tsehgih labolg)%7.53()%3.53(ot esaercni 1991 ecnelaverP ecnelaverP detamitsE aciremA aciremA ni %8.92 %9.12 ylaey(

,aciremA htroN morf htuoS atad acirfA ,aciremA detimiL dna
11(3(4()seiduts eporuE aciremA aciremA 2(,)seiduts htroN htuoS acirfA ,)seiduts ,)seiduts ,)seiduts 281(aisA
8102–1991
452,993,5
542
)1202(21.la te eL

htuoS)849,2(dna gnitisiv ot ot seitilicaf ylno ytienegoreteh elbitpecsus noitroporp stneitap saib elbacilppa erachtlaeh noitceles aciremA egral erew ecneh hgiH ataD A
%42 ecnelaverp lareneg noitalupop ni llarevo ecnelaverp 'evitpac' detamitsE ;%95 dna
)ureP ,lizarB(,anitnegrA ylno aciremA ,elihC ,ocixeM htuoS
526,5
91
)2202(34.la te sajoR

stnemmoC	ot ylno elbacilppa era aisA ataD
sgnidnfi niaM	emit ecnelaverp %5.82 –2102(revo ,)5002–5991(desaercni %9.33 llarevo ,)1102–6002(ecnelaverP detamitsE %3.52 %6.92)7102 –
dedulcni seirtnuoC/snoigeR	aisA
sraey ydutS	7102–4991
slaudividni fo .oN	815,440,31
seiduts fo .oN	732
ydutS	)9102(01.la te iL

Margaret LP Teng, et al. Global incidence and prevalence of NAFLD NASH is now the fastest-rising cause of hepatocellular carcinoma person-years in Japan (Fig. 1).¹⁰ The NAFLD incidence in South Korea worldwide^{6,7} and is also the fastest-rising indication for Korea was around 45 cases per 1,000 person-years.^{10,11} Taken liver transplantation in the United States.⁸ together, the estimates for NAFLD incidence in Asia remain consistent across several meta-analyses (Table 1). INCIDENCE OF NAFLD PREVALENCE OF NAFLD A recent meta-analysis by Riazi et al.³ estimated the incidence of NAFLD at 46.9 cases per 1,000 person-years. The in- Riazi et al.³ pooled data from 72 studies (1,030,160 individuals) of NAFLD was higher in males (70.8 cases per 1,000 males) and estimated that the global prevalence of NAFLD in person-years) vs. females (26.9 cases per 1,000 person-years, adults was 32% (Table 2). The prevalence was higher in males $P < 0.0001$. However, all included studies were conducted in males (40% vs. 26%, $P < 0.0001$). The prevalence of Asia, hence it is unclear whether these data are generalizable NAFLD increased from 26% in studies from 2005 or earlier to to other parts of the world. A previous meta-analysis published 38% in studies from 2016 or beyond. However, data from this published in 2016 had estimated the NAFLD incidence at 52.34 study by Riazi et al.³ requires cautious interpretation, as data per 1,000 person-years in Asia and 28.01 per 1,000 person-years were available from only 17 countries, hence it is unclear if years in Israel.⁹ Another meta-analysis focused on NAFLD in the estimates from this study are a true reflection of 'global' Asia reported an incidence of 50.9 per 1,000 person-years, prevalence. The relative lack of studies emphasizes the need with the highest incidence of 63 per 1,000 person-years in to improve data collection from regions such as Africa, Oceania, mainland China and the lowest incidence of 29 per 1,000 person-years, and South America, where data was lacking. Le et al.¹² Estimated incidence of nonalcoholic fatty liver disease ≥ 40 cases per 1000 person-years 30-40 cases per 1000 person-years < 30 cases per 1000 person-years Created with mapchart.net Figure 1. Estimated incidence of nonalcoholic fatty liver disease. Data for China, Hong Kong, Japan, and South Korea was obtained from Li et al.¹⁰ and Riazi et al.³. Data for Israel was obtained from Riazi et al.³ and Younossi et al.⁹. NAFLD, nonalcoholic fatty liver disease. Abbreviations: NAFLD, nonalcoholic fatty liver disease; NASH, nonalcoholic steatohepatitis; HCC, hepatocellular carcinoma; BMI, body mass index; NHANES, National Health and Nutrition Examination Surveys; PNPLA3, patatin-like phospholipase domain-containing protein 3; FLI, fatty liver index S33 <http://www.e-cmh.org> <https://doi.org/10.3350/cmh.2022.0365> Margaret LP Teng, et al. Global incidence and prevalence of NAFLD Estimated prevalence of nonalcoholic fatty liver disease $\geq 40\%$ 35-39.9% 30-34.9% 25-29.9% 20-25% Created with mapchart.net Figure 2. Estimated prevalence of nonalcoholic fatty liver disease (NAFLD). Data for Iran, China, Taiwan, South Korea, Europe (Turkey, Italy, Germany, Portugal, Spain), North America (USA), and Egypt was obtained from Riazi et al.³. Data for South America was obtained from Le et al.¹². Data for South Asia (India, Bangladesh, Sri Lanka) and Southeast Asia (Indonesia, Malaysia,

Singapore) was obtained from Li et al.¹⁰. Data for Japan was obtained from Li et al.¹⁰ and Riazi et al.³. Data for Sudan was obtained from Younossi et al.⁹. Data for Australia was obtained from population-based studies by Farrell et al.⁵⁵ and Roberts et al.⁵⁴. Projections for prevalence of NAFLD 35 30 25 20 15 10 5 0 China South Korea Singapore Saudi Arabia Japan Germany France Italy Spain UK US Canada Australia 2015-2019 2030 Figure 3. Estimated projections for the prevalence of nonalcoholic fatty liver disease (NAFLD). Data was obtained from Estes et al.^{58,59,61}. Data for Saudi Arabia was obtained from Alswat et al.⁶⁰, data for Canada was obtained from Swain et al.⁶², and data for Australia was obtained from Adams et al.⁶³. S37 <http://www.e-cmh.org> <https://doi.org/10.3350/cmh.2022.0365> Clinical and Molecular Hepatology Volume_29 Supplement February 2023 tries at 23.9% compared to non-Mediterranean countries at included in this meta-analysis were hospital-based studies 28.5%, although the difference was not statistically signifi- and included patients with risk factors for NAFLD, hence the cant. results may not have been fully representative of the general Within Europe, Turkey had the highest NAFLD prevalence population. Le et al.¹² pooled data from 3 studies (5,716 indi- at 48.4%, followed by Italy at 38.2%. Germany, Portugal, and viduals) and determined that South America had the greatest Spain had NAFLD prevalence between 25–27%.³ A cross-sec- estimated NAFLD prevalence among the continents at 35.7%. tional study utilizing data from a large population-based co- This may be due to a combination of genetic susceptibility hort in France found that NAFLD prevalence in France was and a greater prevalence of metabolic risk factors.⁴⁴ There is a 18.2%.³² A study involving individuals from population-based high prevalence of PNPLA3 genetic polymorphism in the studies in Russia reported that NAFLD prevalence was 40% in general population, especially among individuals with Native the Ural Eye and Medical Study (UEMS) (5,852 individuals), American ancestry.⁴⁵⁻⁴⁷ Furthermore, obesity is extremely and 69.8% in the Ural Very Old Study (UVOS) (1,130 individu- common in the region—a cross-sectional study across 4 als).³³ However, it should be noted that in the UVOS, individu- geographical regions found that central obesity was highest als were older with minimum age of 85 years, and methods in South America.⁴⁸ Type 2 diabetes has also been rising in for diagnosis of NAFLD differed between studies as well. prevalence in South America.⁴⁹ Data from the meta-analysis by Le showed that compared to other regions, NAFLD indi- North America viduals in South America had a higher likelihood of having diabetes and higher mean cholesterol levels.¹² In addition, Based on subgroup data from 4 studies (18,356 individuals), physical activity is often inadequate—Latin America was Le et al.¹² estimated that the prevalence of NAFLD in North ranked as the top region for physical inactivity, with a third of America was 35.3%. More recently, NAFLD prevalence was the population experiencing a lack of physical activity.⁵⁰ reported at 47.8% in the meta-analysis by Riazi which includ- ed 2 large studies with 15,178 individuals from the USA.³ This Africa is driven by a high prevalence of obesity in the USA. In North America, Hispanics have the highest NAFLD prevalence, fol- There is a paucity of data from Africa on the epidemiology lowed by non-Hispanic Whites and non-Hispanic blacks.³⁴⁻³⁹ of NAFLD. A meta-analysis estimated the prevalence of Based on data from National Health and Nutrition Examina- NAFLD in Africa at 13.5%, ranging from 9% in Nigeria to 20% tion Surveys (NHANES) 2017–2018, NAFLD prevalence was es- in

Sudan.⁵¹ More recently, NAFLD prevalence was reported at 63.7% in Hispanics, 56.8% in non-Hispanic whites, 28.2% in the meta-analysis by Le, and 56.8% in the meta- and 46.2% in non-Hispanic blacks.³⁷ This could be attributed analysis by Riazi.^{3,12} Of note, the meta-analysis by Riazi only to genetic factors like the patatin-like phospholipase do- included 1 study from Egypt. The wide variation in estimates main-containing protein 3 (PNPLA3) mutation, which is asso- of NAFLD prevalence is likely related to a lack of reliable data ciated with elevated risk for hepatic steatosis and NASH, oc- from Africa.^{52,53} ccurring more frequently in Hispanics.^{40,41} This could also be related to metabolic factors like the higher prevalence of Oceania central adiposity and insulin resistance in Hispanics com- pared to non-Hispanic whites.^{34,42} Lower serum triglyceride Likewise, there is scarce data from Oceania on the inci- levels in African-Americans may also contribute to reduced dence and prevalence of NAFLD. Population-based studies NAFLD prevalence.³⁸ using fatty liver index have demonstrated NAFLD prevalence of 35.7–38% in Australia.^{54,55} There are no population-based South America studies on NAFLD prevalence using imaging modalities such as ultrasound.⁵⁶ A meta-analysis comprising 19 studies (5,626 individuals) by Rojas et al. estimated the prevalence of NAFLD in South America at up to 59%.⁴³ Notably, the majority of the studies S38 <https://doi.org/10.3350/cmh.2022.0365> <http://www.e-cmh.org> Margaret LP Teng, et al. Global incidence and prevalence of NAFLD Adiposity and Its Relationship With Cardiometabolic Risk/Intra- transient elastography among US adults: results from NHANES Abdominal Adiposity. *Am J Clin Nutr* 2012;96:714-726. 2017-2018. *PLoS One* 2021;16:e0252164. 26. Hanlon CL, Yuan L. Nonalcoholic fatty liver disease: the role of 38. Sherif ZA, Saeed A, Ghavimi S, Nouraie SM, Laiyemo AO, Brim visceral adipose tissue. *Clin Liver Dis (Hoboken)* 2022;19:106- H, et al. Global epidemiology of nonalcoholic fatty liver dis- 110. ease and perspectives on US minority populations. *Dig Dis Sci* 27. Yoon KH, Lee JH, Kim JW, Cho JH, Choi YH, Ko SH, et al. Epidemic 2016;61:1214-1225. obesity and type 2 diabetes in Asia. *Lancet* 2006;368:1681-1688. 39. Arshad T, Golabi P, Henry L, Younossi ZM. Epidemiology of non- 28. Huxley R, James WP, Barzi F, Patel JV, Lear SA, Suriyawongpaisal alcoholic fatty liver disease in North America. *Curr Pharm Des P*, et al. Ethnic comparisons of the cross-sectional relationships 2020;26:993-997. between measures of body size with diabetes and hyperten- 40. Romeo S, Kozlitina J, Xing C, Pertsemlidis A, Cox D, Pennacchio sion. *Obes Rev* 2008;9 Suppl 1:53-61. LA, et al. Genetic variation in PNPLA3 confers susceptibility to 29. Chan JC, Malik V, Jia W, Kadowaki T, Yajnik CS, Yoon KH, et al. nonalcoholic fatty liver disease. *Nat Genet* 2008;40:1461-1465. Diabetes in Asia: epidemiology, risk factors, and pathophysiol- 41. Wagenknecht LE, Palmer ND, Bowden DW, Rotter JI, Norris JM, ogy. *JAMA* 2009;301:2129-2140. Ziegler J, et al. Association of PNPLA3 with non-alcoholic fatty 30. Hagström H, Nasr P, Ekstedt M, Hammar U, Stål P, Hultcrantz R, liver disease in a minority cohort: the Insulin Resistance Athero- et al. Risk for development of severe liver disease in lean pa- scleriosis Family Study. *Liver Int* 2011;31:412-416. tients with nonalcoholic fatty liver disease: a long-term follow- 42. Guerrero R, Vega GL, Grundy SM, Browning JD. Ethnic differ- up study. *Hepatol Commun* 2018;2:48-57. ences in hepatic steatosis: an insulin resistance paradox? Hepa- 31. Cholongitas E, Pavlopoulou I, Papatheodoridi M, Markakis GE, tology 2009;49:791-801. Bouras E, Haidich AB, et al.

Epidemiology of nonalcoholic fatty 43. Rojas YAO, Cuellar CLV, Barrón KMA, Arab JP, Miranda AL. Non- liver disease in Europe: a systematic review and meta-analysis. alcoholic fatty liver disease prevalence in Latin America: a sys- Ann Gastroenterol 2021;34:404-414. tematic review and meta-analysis. Ann Hepatol 2022;27:100706. 32. Nabi O, Lacombe K, Boursier J, Mathurin P, Zins M, Serfaty L. 44. Pinto Marques Souza de Oliveira C, Pinchemel Cotrim H, Arrese Prevalence and risk factors of nonalcoholic fatty liver disease M. Nonalcoholic fatty liver disease risk factors in Latin American and advanced fibrosis in general population: the French Na- populations: current scenario and perspectives. Clin Liver Dis tionwide NASH-CO Study. Gastroenterology 2020;159:791-793. (Hoboken) 2019;13:39-42. e2. 45. Arrese M, Arab JP, Riquelme A, Benítez CE, Barrera F, Soza A, 33. Bikbov MM, Gilmanshin TR, Zainullin RM, Kazakbaeva GM, Iaku- et al. High prevalence of PNPLA3 rs738409 (I148M) polymor- pova EM, Fakhretdinova AA, et al. Prevalence of non-alcoholic phism in Chilean latino population and its association to non- fatty liver disease in the Russian Ural Eye and Medical Study alcoholic fatty liver disease risk and histological disease sever- and the Ural Very Old Study. Sci Rep 2022;12:7842. ity. Hepatology 2015;62 Suppl 1:1285A. 34. Browning JD, Szczepaniak LS, Dobbins R, Nuremberg P, Horton 46. Pontoriero AC, Trinks J, Hulaniuk ML, Caputo M, Fortuny L, Pratx JD, Cohen JC, et al. Prevalence of hepatic steatosis in an urban LB, et al. Influence of ethnicity on the distribution of genetic population in the United States: impact of ethnicity. Hepatol- polymorphisms associated with risk of chronic liver disease in ogy 2004;40:1387-1395. South American populations. BMC Genet 2015;16:93. 35. Lazo M, Hernaez R, Eberhardt MS, Bonekamp S, Kamel I, Gual- 47. Chinchilla-López P, Ramírez-Pérez O, Cruz-Ramón V, Canizales- lar E, et al. Prevalence of nonalcoholic fatty liver disease in the Quinteros S, Domínguez-López A, Ponciano-Rodríguez G, et al. United States: the Third National Health and Nutrition Examina- More evidence for the genetic susceptibility of Mexican popu- tion Survey, 1988-1994. Am J Epidemiol 2013;178:38-45. lation to nonalcoholic fatty liver disease through PNPLA3. Ann 36. Kim D, Kim W, Adejumo AC, Cholanteril G, Tighe SP, Wong RJ, Hepatol 2018;17:250-255. et al. Race/ethnicity-based temporal changes in prevalence of 48. Patel SA, Ali MK, Alam D, Yan LL, Levitt NS, Bernabe-Ortiz A, et NAFLD-related advanced fibrosis in the United States, 2005- al. Obesity and its relation with diabetes and hypertension: a 2016. Hepatol Int 2019;13:205-213. cross-sectional study across 4 geographical regions. Glob Heart 37. Zhang X, Heredia NI, Balakrishnan M, Thrift AP. Prevalence and 2016;11:71-79.e4. factors associated with NAFLD detected by vibration controlled 49. Gallardo-Rincón H, Cantoral A, Arrieta A, Espinal C, Magnus MH, S41 <http://www.e-cmh.org> <https://doi.org/10.3350/cmh.2022.0365>