

MCA Sem. I Core

Mathematical Techniques for Computer Applications(MCAC 103) L 5

Expectation and Variance of Random Variables

3 Feb 2022

Expectation

Moments

Statistical Inequalities

Covariance

Conditional
Expectation

Vasudha Bhatnagar
Department of Computer Science, University of Delhi
Delhi, India.

Outline

1 Expectation

2 Moments

3 Statistical Inequalities

4 Covariance

5 Conditional Expectation

Expectation

Moments

Statistical Inequalities

Covariance

Conditional
Expectation

Expectation of a random variable X is its mean (weighted average)

Expectation

Moments

Statistical Inequalities

Covariance

Conditional
Expectation

Discrete case - RV X

$$\mu = E[X] = \sum_x xP(x)$$

Continuous case - RV X

$$\mu = E[X] = \int_x xf(x)dx$$

Expectation of a random variable X is its mean (weighted average)

Expectation

Moments

Statistical Inequalities

Covariance

Conditional
Expectation

Discrete case - RV X

$$\mu = E[X] = \sum_x xP(x)$$

Continuous case - RV X

$$\mu = E[X] = \int_x xf(x)dx$$

Practice Exercise - Consider RV X

- ① If $P(0) = P(1) = \frac{1}{2}$, then $E[X] = \frac{1}{2}$
- ② If $P(0) = \frac{1}{4}$, $P(1) = \frac{3}{4}$, then $E[X] = \frac{3}{4}$
- ③ If X is the outcome when we roll a fair die, then $E[X] = \frac{7}{2}$

Practice Exercise

- ① Find the expected value of a discrete random variable X having the probability distribution

$$f(x) = \frac{|x - 2|}{7} \text{ for } x = -1, 0, 1, 3$$

(1/7)

- ② Find the expected value of a continuous random variable X having the probability density

$$f(x) = \begin{cases} \frac{1}{8}(x + 1) & \text{for } 2 < x < 4 \\ 0 & \text{elsewhere} \end{cases}$$

(37/12)

- ③ If the probability density is given by

$$f(x) = \begin{cases} x/2 & \text{for } 0 < x \leq 1 \\ 1/2 & \text{for } 1 < x \leq 2 \\ (3 - x)/2 & \text{for } 2 < x < 3 \\ 0 & \text{elsewhere} \end{cases}$$

find $E[X]$.

Expectation of function of X - $g(x)$

Discrete case - RV X

$$E[g(x)] = \sum_x g(x)P(x)$$

Continuous case - RV X

$$E[g(x)] = \int_x g(x)f(x)dx$$

Expectation

Moments

Statistical Inequalities

Covariance

Conditional
Expectation

Expectation of function of X - $g(x)$

Discrete case - RV X

$$E[g(x)] = \sum_x g(x)P(x)$$

Continuous case - RV X

$$E[g(x)] = \int_x g(x)f(x)dx$$

Other Properties of $E[X]$

- ① $E[c] = c$, where c is a constant
- ② $E[cX + b] = cE[X] + b$, where c, b are constants
- ③ $E[X + Y] = E[X] + E[Y]$
- ④ If c_1, c_2, \dots, c_n are constants, then
$$E[\sum_{i=1}^n c_i g_i(X)] = [\sum_{i=1}^n c_i E[g_i(X)]]$$

Expectation

Moments

Statistical Inequalities

Covariance

Conditional
Expectation

Practice Exercise

- 1 Find $E[g(X)]$, where $g(X) = X^2 - 5X + 3$, for Q3 on slide 4.

Practice Exercise

- 1 Find $E[g(X)]$, where $g(X) = X^2 - 5X + 3$, for Q3 on slide 4. $(-11/6)$
- 2 If X is the number obtained after rolling a dice, find $E[2X^2 + 1]$

Practice Exercise

- 1 Find $E[g(X)]$, where $g(X) = X^2 - 5X + 3$, for Q3 on slide 4. $(-11/6)$
- 2 If X is the number obtained after rolling a dice, find $E[2X^2 + 1]$ $(94/3)$

The r^{th} moment about the origin of the random variable X , denoted by μ'_r is expected value of X^r

$$\mu'_r = E[X^r] = \sum_x x^r P(x)$$

when X is discrete

$$\mu'_r = E[X^r] = \int_x x^r f(x) dx$$

When X is continuous rv

[Expectation](#)[Moments](#)[Statistical Inequalities](#)[Covariance](#)[Conditional
Expectation](#)

The r^{th} moment about the origin of the random variable X , denoted by μ'_r is expected value of X^r

$$\mu'_r = E[X^r] = \sum_x x^r P(x)$$

when X is discrete

$$\mu'_r = E[X^r] = \int_x x^r f(x) dx$$

When X is continuous rv

The r^{th} moment about mean of the random variable X , denoted by μ_r is expected value of $(X - \mu)^r$

$$\mu_r = E[(X - \mu)^r] = \sum_x (x - \mu)^r P(x)$$

when X is discrete

$$\mu_r = E[(X - \mu)^r] = \int_x (x - \mu)^r f(x) dx$$

When X is continuous rv

Note that

- ① μ'_1 is mean of the distribution of X (denoted by μ)
- ② μ_2 is the variance of the distribution denoted by σ^2
- ③ Positive square root of μ_2 is the standard deviation denoted by σ
- ④ $\sigma^2 = \mu'_2 - \mu^2$
- ⑤ μ_3 describes skewness
- ⑥ If X has the variance σ^2 , then $Var(aX + b) = a^2\sigma^2$

Practice Exercises

- ① If X denotes the number on the dice, calculate variance of X .

Practice Exercises

- 1 If X denotes the number on the dice, calculate variance of X . (35/12)
- 2 A construction firm has recently sent in bids for 3 jobs worth (in profits) 10, 20, and 40 (thousand) INR. If its probabilities of winning the jobs are respectively .2, .8, and .3, what is the firm's expected total profit?

Practice Exercises

- 1 If X denotes the number on the dice, calculate variance of X . (35/12)
- 2 A construction firm has recently sent in bids for 3 jobs worth (in profits) 10, 20, and 40 (thousand) INR. If its probabilities of winning the jobs are respectively .2, .8, and .3, what is the firm's expected total profit? (INR 30,000)
- 3 A group of n cars enters an intersection from the south. Through prior observations, it is estimated that each car has the probability p of turning east, probability q of turning west, and probability r of going straight on ($p + q + r = 1$). Assume that drivers behave independently and let X be the number of cars turning west. Find $E[X]$.

Practice Exercises

- 1 If X denotes the number on the dice, calculate variance of X . (35/12)
- 2 A construction firm has recently sent in bids for 3 jobs worth (in profits) 10, 20, and 40 (thousand) INR. If its probabilities of winning the jobs are respectively .2, .8, and .3, what is the firm's expected total profit? (INR 30,000)
- 3 A group of n cars enters an intersection from the south. Through prior observations, it is estimated that each car has the probability p of turning east, probability q of turning west, and probability r of going straight on ($p + q + r = 1$). Assume that drivers behave independently and let X be the number of cars turning west. Find $E[X]$. (nq)
- 4 The waiting time X (in minutes) of a customer waiting to be served at a ticket counter has the density function

$$f_X(x) = \begin{cases} 2e^{-2x} & \text{for } x > 0 \\ 0 & \text{elsewhere} \end{cases}$$

find $E[X]$.

Practice Exercises

- 1 If X denotes the number on the dice, calculate variance of X . (35/12)
- 2 A construction firm has recently sent in bids for 3 jobs worth (in profits) 10, 20, and 40 (thousand) INR. If its probabilities of winning the jobs are respectively .2, .8, and .3, what is the firm's expected total profit? (INR 30,000)
- 3 A group of n cars enters an intersection from the south. Through prior observations, it is estimated that each car has the probability p of turning east, probability q of turning west, and probability r of going straight on ($p + q + r = 1$). Assume that drivers behave independently and let X be the number of cars turning west. Find $E[X]$. (nq)
- 4 The waiting time X (in minutes) of a customer waiting to be served at a ticket counter has the density function

$$f_X(x) = \begin{cases} 2e^{-2x} & \text{for } x > 0 \\ 0 & \text{elsewhere} \end{cases}$$

find $E[X]$. (1/2 minutes)

Markov Inequality: $P(X \geq a) \leq \frac{E(X)}{a}$, for $X, a \geq 0$

- 1 Gives probability bound for extreme values of a rv i.e. $P(X \geq a)$, where $X \geq 0$
- 2 Useful when the exact distribution of X is unknown, but $E(X)$ is known
- 3 Intuitive Proof,

Markov Inequality: $P(X \geq a) \leq \frac{E(X)}{a}$, for $X, a \geq 0$

- ① Gives probability bound for extreme values of a rv i.e. $P(X \geq a)$, where $X \geq 0$
- ② Useful when the exact distribution of X is unknown, but $E(X)$ is known
- ③ Intuitive Proof,

$$E(X) = \sum_{x=0}^{\infty} xP(x) = \sum_{x=0}^{x=a} xP(x) + \sum_{x=a}^{\infty} xP(x)$$

$$E(X) \geq \sum_{x=a}^{\infty} xP(x) \geq a \sum_{x=a}^{\infty} P(x) = aP(X \geq a)$$

- ④ Loose bound - not useful sometimes

Alternate Proof of Markov Inequality

Define a rv Y as follows:

$$Y = \begin{cases} 0 & \text{for } x < a \\ a & \text{for } x \geq a \end{cases}$$

Thus $Y \leq X$

$$E(Y) \leq E(X)$$

$$0 * P(y = 0) + a * P(y = a) \leq E(X)$$


$$a \sum_{x=a}^{x=\infty} P(x) \leq E(X)$$

$$a(P(X \geq a)) \leq E(X)$$

$$P(X \geq a) \leq \frac{E(X)}{a}$$

$$\text{Chebychev's Inequality: } P(|X - \mu| < k\sigma) \geq 1 - \frac{1}{k^2},$$

for rv X with mean and Var as μ, σ and constant $k \geq 0$



- ① Gives lower bound of the probability that rv will take value within k standard deviations
- ② Useful when we do not have exact probability distribution for rv X

$$\text{Chebychev's Inequality: } P(|X - \mu| \geq k) \leq \frac{\sigma^2}{k^2},$$

for rv X with mean and Var as μ, σ and constant $k \geq 0$

Proof using Markov's Inequality

Since $(X - \mu)^2$ is a nonnegative random variable, we can apply Markov's inequality (with $a = k^2$) to obtain

$$P\{(X - \mu)^2 \geq k^2\} \leq \frac{E[(X - \mu)^2]}{k^2}$$

But since $(X - \mu)^2 \geq k^2$ if and only if $|X - \mu| \geq k$, Equation 4.9.1 is equivalent to

$$P\{|X - \mu| \geq k\} \leq \frac{E[(X - \mu)^2]}{k^2} = \frac{\sigma^2}{k^2}$$

and the proof is complete. \square

Alternate Proof

$$\begin{aligned}
 \sigma^2 &= E(X - E(X))^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx \\
 &= \int_{-\infty}^{\mu - k\sigma} (x - \mu)^2 f(x) dx + \int_{\mu - k\sigma}^{\mu + k\sigma} (x - \mu)^2 f(x) dx + \\
 &\quad \int_{\mu + k\sigma}^{\infty} (x - \mu)^2 f(x) dx \\
 \sigma^2 &\geq \int_{-\infty}^{\mu - k\sigma} (x - \mu)^2 f(x) dx + \int_{\mu + k\sigma}^{\infty} (x - \mu)^2 f(x) dx
 \end{aligned}$$

But, $(x - \mu)^2 \geq k^2 \sigma^2$ for $x \leq (\mu - k\sigma)$ or $x \geq (\mu + k\sigma)$, we can write

$$\begin{aligned}
 \sigma^2 &\geq \int_{-\infty}^{\mu - k\sigma} k^2 \sigma^2 f(x) dx + \int_{\mu + k\sigma}^{\infty} k^2 \sigma^2 f(x) dx \\
 \frac{1}{k^2} &\geq \int_{-\infty}^{\mu - k\sigma} f(x) dx + \int_{\mu + k\sigma}^{\infty} f(x) dx \\
 \frac{1}{k^2} &\geq P(|x - \mu| \geq k\sigma) \\
 \frac{1}{k^2} &\geq 1 - P(|x - \mu| < k\sigma) \\
 P(|x - \mu| < k\sigma) &\geq 1 - \frac{1}{k^2}
 \end{aligned}$$

Expectation

Moments

Statistical Inequalities

Covariance

Conditional
Expectation

Covariance of two random variables X and Y, denoted by $Cov(X, Y)$ is denoted by

$$Cov(X, Y) = E[(X - E(X))(Y - E(Y))] = E(XY) - E(X)E(Y)$$

If X and Y are independent random variables, the
 $E(XY) = E(X)E(Y)$

Properties of Covariance for any random variables X, Y, Z and constant k

- ① $Cov(X, X) = Var(X)$
- ② $Cov(X, Y) = Cov(Y, X)$
- ③ $Cov(kX, Y) = kCov(X, Y)$
- ④ $Cov(X, Y+Z) = Cov(X, Y) + Cov(X, Z)$

Practice Exercises

Suppose X_1, X_2, \dots, X_n are independently and identically distributed rvs with $E(X_i) = \mu$ and $Var(X_i) = \sigma^2$. Consider the sample mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ as a derived random variable. Then show that :

- ① $E(\bar{X}) = \mu$
- ② $Var(\bar{X}) = \sigma^2/n$
- ③ $Cov(\bar{X}, X_i - \bar{X}) = 0$

Conditional Expectation

- ① Conditional expectation of a rv is defined in terms of conditional probability distribution.
- ② $E(X|Y)$ is a function of a rv Y , and hence rv itself.

If X is a discrete rv and $p(x|y)$ is the conditional probability distribution of X given $Y=y$ at x , then the conditional expectation $E(X|Y = y)$ is

$$\mu_{X|y} = \sum_x x * p(x|y)$$

Correspondingly, conditional variance is defined as:

$$\sigma_{X|y}^2 = E[(X - \mu_{X|y})^2|y]$$
$$E(X^2|y) - \mu_{X|y}^2$$