

MCA Sem. I Core

# Mathematical Techniques for Computer Applications(MCAC 103) L 2

Conditional Probability and Bayes' Theorem

27 Dec 2021

Vasudha Bhatnagar  
Department of Computer Science, University of Delhi  
Delhi, India.

# Outline

Independent Events

Conditional Probability

Bayes' Theorem

**1 Independent Events**

**2 Conditional Probability**

**3 Bayes' Theorem**

**Multiple events can be associated with an experiment and its sample space**

- ❶ Two successive rolls of a fair dice

**Multiple events can be associated with an experiment and its sample space**

❶ Two successive rolls of a fair dice

❶ E1: Sum is even

**Multiple events can be associated with an experiment and its sample space**

**① Two successive rolls of a fair dice**

**① E1: Sum is even**

**② E2: Both numbers even**

**Multiple events can be associated with an experiment and its sample space**

**① Two successive rolls of a fair dice**

**① E1: Sum is even**

**② E2: Both numbers even**

**③ E3: Both numbers same**

**Multiple events can be associated with an experiment and its sample space**

- ❶ Two successive rolls of a fair dice
  - ❶ E1: Sum is even
  - ❷ E2: Both numbers even
  - ❸ E3: Both numbers same
  - ❹ E4: Both numbers greater than 4

**Multiple events can be associated with an experiment and its sample space**

- ① Two successive rolls of a fair dice**
  - ① E1: Sum is even**
  - ② E2: Both numbers even**
  - ③ E3: Both numbers same**
  - ④ E4: Both numbers greater than 4**
  - ⑤ E5: One number twice the other**



**Multiple events can be associated with an experiment and its sample space**

- ① Two successive rolls of a fair dice**
  - ① E1: Sum is even**
  - ② E2: Both numbers even**
  - ③ E3: Both numbers same**
  - ④ E4: Both numbers greater than 4**
  - ⑤ E5: One number twice the other**
- ② Choosing three cards from a deck of 52 cards**

## Multiple events can be associated with an experiment and its sample space

- ① Two successive rolls of a fair dice
  - ① E1: Sum is even
  - ② E2: Both numbers even
  - ③ E3: Both numbers same
  - ④ E4: Both numbers greater than 4
  - ⑤ E5: One number twice the other
- ② Choosing three cards from a deck of 52 cards
- ③ Time taken by a train to travel between stations A and B on two different days

**Mutual Independence of events  $E_1, E_2, \dots, E_k$** 

**iff** the probability of intersection of any 2, 3, ..., k events is the product of their respective probabilities

**Pairwise independence does not generally lead to mutual independence**

Independent Events

Conditional Probability

Bayes' Theorem

### Mutual Independence of events $E_1, E_2, \dots, E_k$

**iff** the probability of intersection of any 2, 3, ..., k events is the product of their respective probabilities

### Pairwise independence does not generally lead to mutual independence

Consider three events  $A_1, A_2$ , and  $A_3$  defined by  $A_1 = B_1 \cup B_2, A_2 = B_1 \cup B_3, A_3 = B_2 \cup B_3$ , where  $B_1, B_2, B_3$  are mutually exclusive events, each occurring with probability  $1/4$ . Then

$$P(A_1) =$$

Independent Events

Conditional Probability

Bayes' Theorem

### Mutual Independence of events $E_1, E_2, \dots, E_k$

**iff** the probability of intersection of any 2, 3, ..., k events is the product of their respective probabilities

### Pairwise independence does not generally lead to mutual independence

Consider three events  $A_1, A_2$ , and  $A_3$  defined by  $A_1 = B_1 \cup B_2, A_2 = B_1 \cup B_3, A_3 = B_2 \cup B_3$ , where  $B_1, B_2, B_3$  are mutually exclusive events, each occurring with probability  $1/4$ . Then

$$P(A_1) = 1/2 = P(A_2) = P(A_3)$$

$$P(A_1 A_2) =$$

Independent Events

Conditional Probability

Bayes' Theorem

### Mutual Independence of events $E_1, E_2, \dots, E_k$

**iff** the probability of intersection of any 2, 3, ..., k events is the product of their respective probabilities

### Pairwise independence does not generally lead to mutual independence

Consider three events  $A_1, A_2$ , and  $A_3$  defined by  $A_1 = B_1 \cup B_2$ ,  $A_2 = B_1 \cup B_3$ ,  $A_3 = B_2 \cup B_3$ , where  $B_1, B_2, B_3$  are mutually exclusive events, each occurring with probability  $1/4$ . Then

$$P(A_1) = 1/2 = P(A_2) = P(A_3)$$

$$P(A_1 A_2) = 1/4 = P(A_1 A_3) = P(A_2 A_3)$$

$$P(A_1 A_2 A_3) =$$

Independent Events

Conditional Probability

Bayes' Theorem

## Mutual Independence of events $E_1, E_2, \dots, E_k$

**iff** the probability of intersection of any 2, 3, ..., k events is the product of their respective probabilities

## Pairwise independence does not generally lead to mutual independence

Consider three events  $A_1, A_2$ , and  $A_3$  defined by  $A_1 = B_1 \cup B_2$ ,  $A_2 = B_1 \cup B_3$ ,  $A_3 = B_2 \cup B_3$ , where  $B_1, B_2, B_3$  are mutually exclusive events, each occurring with probability  $1/4$ . Then

$$P(A_1) = 1/2 = P(A_2) = P(A_3)$$

$$P(A_1 A_2) = 1/4 = P(A_1 A_3) = P(A_2 A_3)$$

$$P(A_1 A_2 A_3) = 0$$

Events  $A_1, A_2, A_3$  are pairwise, but not mutually independent

## Applications in Reliability

- 1 There is a 1% probability for a hard drive to crash. Therefore, it has two backups, each having a 2% probability to crash, and all three components are mutually independent of each other. The stored information is lost only in an unfortunate situation when all three devices crash. What is the probability that the information is saved?

[Independent Events](#)[Conditional Probability](#)[Bayes' Theorem](#)



## Applications in Reliability

- ① There is a 1% probability for a hard drive to crash. Therefore, it has two backups, each having a 2% probability to crash, and all three components are mutually independent of each other. The stored information is lost only in an unfortunate situation when all three devices crash. What is the probability that the information is saved? (0.999996)
- ② Suppose that a shuttle's launch depends on three key devices that operate independently of each other and malfunction with probabilities 0.01, 0.02, and 0.02, respectively. If any of the key devices malfunctions, the launch will be postponed. Compute the probability for the shuttle to be launched on time, according to its schedule.

Independent Events

Conditional Probability

Bayes' Theorem

## Applications in Reliability

- ① There is a 1% probability for a hard drive to crash. Therefore, it has two backups, each having a 2% probability to crash, and all three components are mutually independent of each other. The stored information is lost only in an unfortunate situation when all three devices crash. What is the probability that the information is saved? (0.999996)
- ② Suppose that a shuttle's launch depends on three key devices that operate independently of each other and malfunction with probabilities 0.01, 0.02, and 0.02, respectively. If any of the key devices malfunctions, the launch will be postponed. Compute the probability for the shuttle to be launched on time, according to its schedule. (0.9508)

Independent Events

Conditional Probability

Bayes' Theorem


## Systems in series and parallel

A system composed of  $n$  separate components is said to be a parallel system if it functions when at least one of the components functions. If the system functions when each one of the components functions, then it is in series.

[Independent Events](#)[Conditional Probability](#)[Bayes' Theorem](#)

## Systems in series and parallel

A system composed of  $n$  separate components is said to be a parallel system if it functions when at least one of the components functions. If the system functions when each one of the components functions, then it is in series.




System functions if either (A & B function) or (C and (either or both (D or E)) function) or both modules function.

Reliability of the system if each component is operable with probability 0.92 independently of the other components -

## Systems in series and parallel

A system composed of  $n$  separate components is said to be a parallel system if it functions when at least one of the components functions. If the system functions when each one of the components functions, then it is in series.



System functions if either (A & B function) or (C and (either or both (D or E)) function) or both modules function.

Reliability of the system if each component is operable with probability 0.92 independently of the other components - (0.9868)

What is  $((A \cap B) \cup (C \cap (D \cup E)))'$ ?


**Conditional probability of event  $A$  given event  $B$ ,  $(P(A|B))$ , is the probability that  $A$  occurs when  $B$  has occurred**

**Conditional probability of event  $A$  given event  $B$ ,  $(P(A|B))$ , is the probability that  $A$  occurs when  $B$  has occurred**

**Useful when (i) initially partial information concerning the result of the experiment is available,**

**Conditional probability of event A given event B, ( $P(A|B)$ ), is the probability that A occurs when B has occurred**

**Useful when (i) initially partial information concerning the result of the experiment is available, (ii) some additional information is available**



$$P(A|B) = P(A \cap B) / P(B), \text{ where } P(B) \text{ is not zero.}$$



	Good service	Poor service
In business $>10$ yrs	16	4
In business $\leq 10$ yrs	10	20

	Good service	Poor service
In business $>10$ yrs	16	4
In business $\leq 10$ yrs	10	20

Find —  $P(\text{Good Service})$ ?  $P(\leq 10 \text{ yrs in business})$ ?  $P(\text{Good Service} \ \& \ \leq 10 \text{ yrs in business})$ ?  $P(\text{Poor Service} \mid \leq 10 \text{ yrs in business})$ ?

- 1 A family has two children. What is the probability that both are boys given that at least one of them is a boy?

	Good service	Poor service
In business >10 yrs	16	4
In business $\leq$ 10 yrs	10	20

**Find —  $P(\text{Good Service})$ ?  $P(\leq 10 \text{ yrs in business})$ ?  $P(\text{Good Service} \ \& \ \leq 10 \text{ yrs in business})$ ?  $P(\text{Poor Service} \mid \leq 10 \text{ yrs in business})$ ?**

- 1 A family has two children. What is the probability that both are boys given that at least one of them is a boy? ( $1/3$ )
- 2 Suppose cards numbered one through ten are placed in a hat, mixed up, and then one of the cards is drawn. If we are told that the number on the drawn card is at least five, then what is the conditional probability that it is ten?

	Good service	Poor service
In business >10 yrs	16	4
In business $\leq$ 10 yrs	10	20

**Find —  $P(\text{Good Service})$ ?  $P(\leq 10 \text{ yrs in business})$ ?  $P(\text{Good Service} \ \& \ \leq 10 \text{ yrs in business})$ ?  $P(\text{Poor Service} \mid \leq 10 \text{ yrs in business})$ ?**

- 1 A family has two children. What is the probability that both are boys given that at least one of them is a boy? ( $1/3$ )
- 2 Suppose cards numbered one through ten are placed in a hat, mixed up, and then one of the cards is drawn. If we are told that the number on the drawn card is at least five, then what is the conditional probability that it is ten? ( $1/6$ )

## Total Probability Theorem: relates the unconditional probability of an event $A$ with its conditional probabilities

Consider a partition of the sample space  $S$  with mutually exclusive and exhaustive events  $B_1, \dots, B_k$ . If these events also partition the event  $A$ , then  $A = (A \cap B_1) \cup \dots \cup (A \cap B_k)$  is also a union of  $k$  mutually exclusive events, and  $P(A) = \sum_{i=1}^k P(A \cap B_i)$ .


## Total Probability Theorem: relates the unconditional probability of an event A with its conditional probabilities

Consider a partition of the sample space  $S$  with mutually exclusive and exhaustive events  $B_1, \dots, B_k$ . If these events also partition the event  $A$ , then  $A = (A \cap B_1) \cup \dots \cup (A \cap B_k)$  is also a union of  $k$  mutually exclusive events, and

$$P(A) = \sum_{i=1}^k P(A \cap B_i).$$

$$P(A) = \sum_{i=1}^k P(A/B_i)P(B_i)$$

Intuitive meaning is that  $P(A)$  is equal to a weighted average of  $P(A/B_i)$ , each term being weighted by the probability of the event on which it is conditioned.



## Example Problems

- 1 In answering a question on a multiple-choice test a student either knows the answer or guesses. Let  $p$  be the probability that she knows the correct answer and  $(1 - p)$  the probability that she guesses. Assume that a student who guesses at the answer will be correct with probability  $1/m$ , where  $m$  is the number of multiple-choice alternatives. What is the conditional probability that a student knew the answer to a question given that she answered it correctly?
- 2 A laboratory blood test is 95 percent effective in detecting a certain disease when it is, in fact, present. However, the test also yields a “false positive” result for 1 percent of the healthy persons tested. (That is, if a healthy person is tested, then, with probability 0.01, the test result will imply he has the disease.) If 0.5 percent of the population actually has the disease, what is the probability a person has the disease given that his test result is positive?

**Bayes' Theorem: Let A and B be two arbitrary events with**

**$P(A) \neq 0$  and  $P(B) \neq 0$ . Then  $P(B|A) = \frac{P(A|B)P(B)}{P(A)}$**

Generalization: if  $B_1, \dots, B_k$  are exhaustive, mutually exclusive events in a sample space  $S$  and  $A$  is another event in the same sample space, then for event  $B_i$

$$P(B_i|A) = \frac{P(AB_i)}{P(A)}$$

$$P(B_i|A) = P(A|B_i) * P(B_i) / \sum_{j=1}^k P(A|B_j)P(B_j)$$

## Applications of Bayes' Theorem: Machine Learning

- 1 Naive Bayes Classifier: Find posterior probabilities of Classes
- 2 Bayesian Belief Networks : Reasoning with uncertain knowledge (formalizes subjective interpretations)




## Evaluation of posteriori probability in terms of a priori information

A simple binary communication channel carries messages by using only two signals, say 0 and 1. We assume that, for a given binary channel, 40% of the time a 1 is transmitted; the probability that a transmitted 0 is correctly received is 0.90, and the probability that a transmitted 1 is correctly received is 0.95. Determine (a) the probability of a 1 being received, and (b) given a 1 is received, the probability that 1 was transmitted.


## Evaluation of posteriori probability in terms of a priori information

A simple binary communication channel carries messages by using only two signals, say 0 and 1. We assume that, for a given binary channel, 40% of the time a 1 is transmitted; the probability that a transmitted 0 is correctly received is 0.90, and the probability that a transmitted 1 is correctly received is 0.95. Determine (a) the probability of a 1 being received, and (b) given a 1 is received, the probability that 1 was transmitted.



## Evaluation of posteriori probability in terms of a priori information

A simple binary communication channel carries messages by using only two signals, say 0 and 1. We assume that, for a given binary channel, 40% of the time a 1 is transmitted; the probability that a transmitted 0 is correctly received is 0.90, and the probability that a transmitted 1 is correctly received is 0.95. Determine (a) the probability of a 1 being received, and (b) given a 1 is received, the probability that 1 was transmitted.



An insurance company believes that people can be divided into two classes — those that are accident prone and those that are not. Their statistics show that an accident-prone person will have an accident at some time within a fixed 1-year period with probability .4, whereas this probability decreases to .2 for a non-accident-prone person. If we assume that 30 percent of the population is accident prone, what is the probability that a new policy holder will have an accident within a year of purchasing a policy?

An insurance company believes that people can be divided into two classes — those that are accident prone and those that are not. Their statistics show that an accident-prone person will have an accident at some time within a fixed 1-year period with probability .4, whereas this probability decreases to .2 for a non-accident-prone person. If we assume that 30 percent of the population is accident prone, what is the probability that a new policy holder will have an accident within a year of purchasing a policy? (.26)

Additional information: Suppose a new policy holder has an accident within a year of purchasing his policy. What is the probability that he is accident prone?

An insurance company believes that people can be divided into two classes — those that are accident prone and those that are not. Their statistics show that an accident-prone person will have an accident at some time within a fixed 1-year period with probability .4, whereas this probability decreases to .2 for a non-accident-prone person. If we assume that 30 percent of the population is accident prone, what is the probability that a new policy holder will have an accident within a year of purchasing a policy? (.26)

Additional information: Suppose a new policy holder has an accident within a year of purchasing his policy. What is the probability that he is accident prone? (.4615)

## Practice Questions

- 1 A new computer program consists of two modules. The first module contains an error with probability 0.2. The second module is more complex; it has a probability of 0.4 to contain an error, independently of the first module. An error in the first module alone causes the program to crash with probability 0.5. For the second module, this probability is 0.8. If there are errors in both modules, the program crashes with probability 0.9. Suppose the program crashed. What is the probability of errors in both modules?

## Practice Questions

- ① A new computer program consists of two modules. The first module contains an error with probability 0.2. The second module is more complex; it has a probability of 0.4 to contain an error, independently of the first module. An error in the first module alone causes the program to crash with probability 0.5. For the second module, this probability is 0.8. If there are errors in both modules, the program crashes with probability 0.9. Suppose the program crashed. What is the probability of errors in both modules? (0.1856)
- ② Of the one million animals in a Zoo, most are good-natured. But one hundred of them are dangerous. Zoo scientists develop an “Dangerous animal Alarm” which is to be tested for reliability by conducting trials. When presented with a dangerous animal, the alarm goes off 99% of the time. When presented with a good-natured animal, the alarm goes off 1% of the time. If an animal sets off the alarm, what is the probability that it is dangerous? Should the Zoo deploy the system?



## Practice Questions

- ① A new computer program consists of two modules. The first module contains an error with probability 0.2. The second module is more complex; it has a probability of 0.4 to contain an error, independently of the first module. An error in the first module alone causes the program to crash with probability 0.5. For the second module, this probability is 0.8. If there are errors in both modules, the program crashes with probability 0.9. Suppose the program crashed. What is the probability of errors in both modules? (0.1856)
- ② Of the one million animals in a Zoo, most are good-natured. But one hundred of them are dangerous. Zoo scientists develop an "Dangerous animal Alarm" which is to be tested for reliability by conducting trials. When presented with a dangerous animal, the alarm goes off 99% of the time. When presented with a good-natured animal, the alarm goes off 1% of the time. If an animal sets off the alarm, what is the probability that it is dangerous? Should the Zoo deploy the system? (0.01, No)