

Tutorial 3

AID-521: Mathematics for Data Science

Abhishek Samantray, Vimal Kumar

August 20, 2021

Qn. 1.

A process for refining sugar yields up to 1 ton of pure sugar per day, but the actual amount produced, Y , is a random variable because of machine breakdowns and other slowdowns. Suppose that Y has density function given by

$$f(y) = \begin{cases} 2y, & 0 \leq y \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

The company is paid at the rate of Rs. 300 per ton for the refined sugar, but it also has a fixed overhead cost of Rs. 100 per day. Thus the daily profit, in hundreds of rupees, is $U = 3Y - 1$.

- (a) Find the probability density function for U , using method of distribution functions.
- (b) Find the probability density function for U , using the transformation method.
- (c) Find the density function of $V = -4Y + 3$.

Ans/Sol.

Qn. 2.

Y_1 and Y_2 have a joint density function

$$f(y_1, y_2) = \begin{cases} e^{-(y_1+y_2)}, & 0 \leq y_1, 0 \leq y_2, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Find the density function for $Y_1 + Y_2$.

Ans/Sol.

Qn. 3.

The fracture strength of tempered glass averages 14 (measured in thousands of pounds per square inch) and has standard deviation 2.

- (a) What is the probability that the average fracture strength of 100 randomly selected pieces of this glass exceeds 14.5?
- (b) Find an interval that includes, with probability 0.95, the average fracture strength of 100 randomly selected pieces of this glass.

Ans/Sol.