Find all the local maxima, local minima, and saddle points of following functions:

1. $f(x, y) = x^2 + xy + y^2 + 3x - 3y + 4$
2. $f(x, y) = x^2 + 3xy + 3y^2 - 6x + 3y - 6$
3. $f(x, y) = 2xy - 5x^2 - 2y^2 + 4x + 4y - 4$
4. $f(x, y) = 2xy - 5x^2 - 2y^2 + 4x - 4$
5. $f(x, y) = x^2 + xy + 3x + 2y + 5$
6. $f(x, y) = y^2 + xy - 2x - 2y + 2$
7. $f(x, y) = 5xy - 7x^2 + 3x - 6y + 2$
8. $f(x, y) = 2xy - x^2 - 2y^2 + 3x + 4$
9. $f(x, y) = x^2 - 4xy + y^2 + 6y + 2$
10. $f(x, y) = 3x^2 + 6xy + 7y^2 - 2x + 4y$
11. $f(x, y) = 2x^2 + 3xy + 4y^2 - 5x + 2y$
12. $f(x, y) = 4x^2 - 6xy + 5y^2 - 20x + 26y$
13. $f(x, y) = x^2 - y^2 - 2x + 4y + 6$
14. $f(x, y) = x^2 - 2xy + 2y^2 - 2x + 2y + 1$
15. $f(x, y) = x^2 + 2xy$
16. $f(x, y) = 3 + 2x + 2y - 2x^2 - 2xy - y^2$
17. $f(x, y) = x^3 - y^3 - 2xy + 6$
18. $f(x, y) = x^3 + 3xy + y^3$
19. $f(x, y) = 6x^2 - 2x^3 + 3y^2 + 6xy$
20. $f(x, y) = 3y^2 - 2y^3 - 3x^2 + 6xy$
21. $f(x, y) = 9x^3 + y^3/3 - 4xy$
22. $f(x, y) = 8x^3 + y^3 + 6xy$
23. $f(x, y) = x^3 + y^3 + 3x^2 - 3y^2 - 8$
24. $f(x, y) = 2x^3 + 2y^3 - 9x^2 + 3y^2 - 12y$
25. $f(x, y) = 4xy - x^4 - y^4$
26. $f(x, y) = x^4 + y^4 + 4xy$
27. $f(x, y) = \dfrac{1}{x^2 + y^2 - 1}$
28. $f(x, y) = \dfrac{1}{x} + xy + \dfrac{1}{y}$
29. $f(x, y) = y \sin x$
30. $f(x, y) = e^{2x} \cos y$

Find the absolute maxima and minima of following functions on the given domains:

31. $f(x, y) = 2x^2 - 4x + y^2 - 4y + 1$ on the closed triangular plate bounded by the lines $x = 0, y = 2, y = 2x$ in the first quadrant

32. $D(x, y) = x^2 - xy + y^2 + 1$ on the closed triangular plate in the first quadrant bounded by the lines $x = 0, y = 4, y = x$

33. $f(x, y) = x^2 + y^2$ on the closed triangular plate bounded by the lines $x = 0, y = 0, y + 2x = 2$ in the first quadrant

34. $T(x, y) = x^2 + xy + y^2 - 6x$ on the rectangular plate $0 \le x \le 5, -3 \le y \le 3$

35. $T(x, y) = x^2 + xy + y^2 - 6x + 2$ on the rectangular plate $0 \le x \le 5, -3 \le y \le 0$

36. $f(x, y) = 48xy - 32x^3 - 24y^2$ on the rectangular plate $0 \le x \le 1, 0 \le y \le 1$

Taylor's series expansions:

Finding Quadratic and Cubic Approximations

In Exercises , use Taylor's formula for $f(x, y)$ at the origin to find quadratic and cubic approximations of f near the origin.

1. $f(x, y) = xe^y$ **2.** $f(x, y) = e^x \cos y$

3. $f(x, y) = y \sin x$ **4.** $f(x, y) = \sin x \cos y$

5. $f(x, y) = e^x \ln(1 + y)$ **6.** $f(x, y) = \ln(2x + y + 1)$

7. $f(x, y) = \sin(x^2 + y^2)$ **8.** $f(x, y) = \cos(x^2 + y^2)$

Chain Rule: One Independent Variable

In Exercises 1–6, **(a)** express dw/dt as a function of t, both by using the Chain Rule and by expressing w in terms of t and differentiating directly with respect to t. Then **(b)** evaluate dw/dt at the given value of t.

1. $w = x^2 + y^2, \quad x = \cos t, \quad y = \sin t; \quad t = \pi$

2. $w = x^2 + y^2, \quad x = \cos t + \sin t, \quad y = \cos t - \sin t; \quad t = 0$

3. $w = \frac{x}{z} + \frac{y}{z}, \quad x = \cos^2 t, \quad y = \sin^2 t, \quad z = 1/t; \quad t = 3$

4. $w = \ln(x^2 + y^2 + z^2), \quad x = \cos t, \quad y = \sin t, \quad z = 4\sqrt{t};$ $t = 3$

5. $w = 2ye^x - \ln z, \quad x = \ln(t^2 + 1), \quad y = \tan^{-1} t, \quad z = e^t;$ $t = 1$

6. $w = z - \sin xy, \quad x = t, \quad y = \ln t, \quad z = e^{t-1}; \quad t = 1$

Method of Lagrange Multiplier

Three Independent Variables with One Constraint

17. **Minimum distance to a point** Find the point on the plane $x + 2y + 3z = 13$ closest to the point $(1, 1, 1)$.

18. **Maximum distance to a point** Find the point on the sphere $x^2 + y^2 + z^2 = 4$ farthest from the point $(1, -1, 1)$.

19. **Minimum distance to the origin** Find the minimum distance from the surface $x^2 + y^2 - z^2 = 1$ to the origin.

20. **Minimum distance to the origin** Find the point on the surface $z = xy + 1$ nearest the origin.

21. **Minimum distance to the origin** Find the points on the surface $z^2 = xy + 4$ closest to the origin.

22. **Minimum distance to the origin** Find the point(s) on the surface $xyz = 1$ closest to the origin.

23. **Extrema on a sphere** Find the maximum and minimum values of

$$f(x, y, z) = x - 2y + 5z$$

on the sphere $x^2 + y^2 + z^2 = 30$.

24. **Extrema on a sphere** Find the points on the sphere $x^2 + y^2 + z^2 = 25$ where $f(x, y, z) = x + 2y + 3z$ has its maximum and minimum values.

25. **Minimizing a sum of squares** Find three real numbers whose sum is 9 and the sum of whose squares is as small as possible.

26. **Maximizing a product** Find the largest product the positive numbers x, y, and z can have if $x + y + z^2 = 16$.

Chain Rule: Two and Three Independent Variables

In Exercises 7 and 8, **(a)** express $\partial z/\partial u$ and $\partial z/\partial v$ as functions of u and v both by using the Chain Rule and by expressing z directly in terms of u and v before differentiating. Then **(b)** evaluate $\partial z/\partial u$ and $\partial z/\partial v$ at the given point (u, v).

7. $z = 4e^x \ln y, \quad x = \ln(u \cos v), \quad y = u \sin v;$
   $(u, v) = (2, \pi/4)$

8. $z = \tan^{-1}(x/y), \quad x = u \cos v, \quad y = u \sin v;$
   $(u, v) = (1.3, \pi/6)$

In Exercises 9 and 10, **(a)** express $\partial w/\partial u$ and $\partial w/\partial v$ as functions of u and v both by using the Chain Rule and by expressing w directly in terms of u and v before differentiating. Then **(b)** evaluate $\partial w/\partial u$ and $\partial w/\partial v$ at the given point (u, v).

9. $w = xy + yz + xz, \quad x = u + v, \quad y = u - v, \quad z = uv;$ $(u, v) = (1/2, 1)$

10. $w = \ln(x^2 + y^2 + z^2), \quad x = ue^v \sin u, \quad y = ue^v \cos u,$ $z = ue^v; \quad (u, v) = (-2, 0)$

In Exercises 11 and 12, **(a)** express $\partial u/\partial x$, $\partial u/\partial y$, and $\partial u/\partial z$ as functions of x, y, and z both by using the Chain Rule and by expressing u directly in terms of x, y, and z before differentiating. Then **(b)** evaluate $\partial u/\partial x$, $\partial u/\partial y$, and $\partial u/\partial z$ at the given point (x, y, z).

11. $u = \dfrac{p - q}{q - r}, \quad p = x + y + z, \quad q = x - y + z,$ $r = x + y - z; \quad (x, y, z) = \left(\sqrt{3}, 2, 1\right)$

12. $u = e^{qr} \sin^{-1} p, \quad p = \sin x, \quad q = z^2 \ln y, \quad r = 1/z;$ $(x, y, z) = (\pi/4, 1/2, -1/2)$

Finding Specified Partial Derivatives

33. Find $\partial w/\partial r$ when $r = 1, s = -1$ if $w = (x + y + z)^2$, $x = r - s, y = \cos(r + s), z = \sin(r + s)$.

34. Find $\partial w/\partial v$ when $u = -1, v = 2$ if $w = xy + \ln z$, $x = v^2/u, y = u + v, z = \cos u$.

35. Find $\partial w/\partial v$ when $u = 0, v = 0$ if $w = x^2 + (y/x)$, $x = u - 2v + 1, y = 2u + v - 2$.

36. Find $\partial z/\partial u$ when $u = 0, v = 1$ if $z = \sin xy + x \sin y$, $x = u^2 + v^2, y = uv$.

37. Find $\partial z/\partial u$ and $\partial z/\partial v$ when $u = \ln 2, v = 1$ if $z = 5 \tan^{-1} x$ and $x = e^u + \ln v$.

38. Find $\partial z/\partial u$ and $\partial z/\partial v$ when $u = 1$ and $v = -2$ if $z = \ln q$ and $q = \sqrt{v + 3} \tan^{-1} u$.

Assuming that the equations in Exercises 25–28 define y as a differentiable function of x, use Theorem 8 to find the value of dy/dx at the given point.

25. $x^3 - 2y^2 + xy = 0, \quad (1, 1)$

26. $xy + y^2 - 3x - 3 = 0, \quad (-1, 1)$

27. $x^2 + xy + y^2 - 7 = 0, \quad (1, 2)$

28. $xe^y + \sin xy + y - \ln 2 = 0, \quad (0, \ln 2)$

Limits with Two Variables

Find the limits in Exercises 1–12.

1. $\lim_{(x,y)\to(0,0)} \frac{3x^2 - y^2 + 5}{x^2 + y^2 + 2}$

2. $\lim_{(x,y)\to(0,4)} \frac{x}{\sqrt{y}}$

3. $\lim_{(x,y)\to(3,4)} \sqrt{x^2 + y^2 - 1}$

4. $\lim_{(x,y)\to(2,-3)} \left(\frac{1}{x} + \frac{1}{y}\right)^2$

5. $\lim_{(x,y)\to(0,\pi/4)} \sec x \tan y$

6. $\lim_{(x,y)\to(0,0)} \cos \frac{x^2 + y^3}{x + y + 1}$

7. $\lim_{(x,y)\to(0,\ln 2)} e^{x-y}$

8. $\lim_{(x,y)\to(1,1)} \ln |1 + x^2 y^2|$

9. $\lim_{(x,y)\to(0,0)} \frac{e^y \sin x}{x}$

10. $\lim_{(x,y)\to(1,1)} \cos \sqrt[3]{|xy| - 1}$

11. $\lim_{(x,y)\to(1,0)} \frac{x \sin y}{x^2 + 1}$

12. $\lim_{(x,y)\to(\pi/2,0)} \frac{\cos y + 1}{y - \sin x}$

Limits of Quotients

Find the limits in Exercises 13–20 by rewriting the fractions first.

13. $\displaystyle\lim_{\substack{(x,y)\to(1,1)\\ x\neq y}} \frac{x^2 - 2xy + y^2}{x - y}$

14. $\displaystyle\lim_{\substack{(x,y)\to(1,1)\\ x\neq y}} \frac{x^2 - y^2}{x - y}$

15. $\displaystyle\lim_{\substack{(x,y)\to(1,1)\\ x\neq 1}} \frac{xy - y - 2x + 2}{x - 1}$

16. $\displaystyle\lim_{\substack{(x,y)\to(2,-4)\\ y\neq -4,\, x\neq x^2}} \frac{y + 4}{x^2y - xy + 4x^2 - 4x}$

17. $\displaystyle\lim_{\substack{(x,y)\to(0,0)\\ x\neq y}} \frac{x - y + 2\sqrt{x} - 2\sqrt{y}}{\sqrt{x} - \sqrt{y}}$

18. $\displaystyle\lim_{\substack{(x,y)\to(2,2)\\ x+y\neq 4}} \frac{x + y - 4}{\sqrt{x + y} - 2}$

19. $\displaystyle\lim_{\substack{(x,y)\to(2,0)\\ 2x-y\neq 4}} \frac{\sqrt{2x - y} - 2}{2x - y - 4}$

20. $\displaystyle\lim_{\substack{(x,y)\to(4,3)\\ x\neq y+1}} \frac{\sqrt{x} - \sqrt{y + 1}}{x - y - 1}$

Continuity in the Plane

At what points (x, y) in the plane are the functions in Exercises 27–30 continuous?

27. **a.** $f(x, y) = \sin(x + y)$ **b.** $f(x, y) = \ln(x^2 + y^2)$

28. **a.** $f(x, y) = \dfrac{x + y}{x - y}$ **b.** $f(x, y) = \dfrac{y}{x^2 + 1}$

29. **a.** $g(x, y) = \sin \dfrac{1}{xy}$ **b.** $g(x, y) = \dfrac{x + y}{2 + \cos x}$

30. **a.** $g(x, y) = \dfrac{x^2 + y^2}{x^2 - 3x + 2}$ **b.** $g(x, y) = \dfrac{1}{x^2 - y}$