SWASTHYA DIAGNOSTICS
DEFINING HEALTH
A Unit of Kidney Care Centre, Panchkula

Dr. Kavita Goyal
MBBS, MD Pathology
swasthyadiagnostics.sd@gmail.com

UHID No	:-	UD-24250009812	**Reg No**	:-	24250009812
Patient Name	:-	Mr VIPIN KUMAR GARG	**Sample Gnt At**	:-	27 Aug 2025 08:59 AM
Age/Sex	:-	67 Y/M	**Sample Rvd At**	:-	27 Aug 2025 08:59 AM
Address	:-	323, SECTOR 9 PANCHKULA , Panchkula	**Report Gnt At**	:-	27 Aug 2025 04:25 PM
Mobile No	:-	9815311118	**Consultant**	:-	Dr AJAY GOYAL (DM(NEPHROLOGY))

Final Report

Investigations	Observations	Biological Ref. Interval	Unit
BIOCHEMISTRY			
IRON STUDIES	-		
SERUM IRON	60.0	60.00 - 145.00	ug / dl
UIBC (UNSATURATED IRON BINDING CAPACITY)	223.8	120.00 - 345.00	ug/dl
TIBC (TOTAL IRON BINDING CAPACITY)	283.80	250.00 - 450.00	ug/dl
TRANSFERRIN SATURATION	21.14	14.00 - 50.00	%

Interpretation :

1. Measurements of serum iron, TIBC and the percentage of iron saturation of transferrin are useful screening tests for iron deficiency anaemia.

2. However, serum iron exhibits significant diurnal variation and may transiently rise or reach reference values after dietary or iron supplements & post blood transfusion.

3. The diagnostic specificity of a low serum iron for iron deficiency is lost in the presence of acute & chronic inflammatory processes as the concentrations of iron and transferrin in the serum are significantly affected, and fall rapidly as part of the acute phase response irrespective of the iron stores status in the body.

4. Hence, Concurrent measurement of the markers mentioned in the below interpretative table alongwith serum iron studies improves the diagnostic specificity for iron deficiency anaemia & also provides a reliable work up for microcytic hypochromic anaemia.

FERRITIN, SERUM	33.39	20.00 - 350.00	ng/mL

Interpretation : 1. Increased ferritin is seen in iron overload as in multiple blood transfusions, hemochromatosis and anemia of chronic disorders . 2. Decreased ferritin levels are seen in iron deficiency anemia, early stage before iron deficiency manifests as anemia. 3. Increased ferritin is also seen in liver disease,alcoholism, inflammatory conditions, leukemia, Hodgkins disease and some malignancies. 4. Levels of ferritin are used for monitoring of iron levels during pregnancy, dialysis and during iron therapy

SWASTHYA DIAGNOSTICS
DEFINING HEALTH
A Unit of kidney Care Centre, Panchkula

Dr. Kavita Goyal
MBBS, MD Pathology
swasthyadiagnostics.sd@gmail.com

UHID No	**:-**	UD-24250009812	**Reg No**	**:-**	24250009812
Patient Name	**:-**	Mr VIPIN KUMAR GARG	**Sample Gnt At**	:-	27 Aug 2025 08:59 AM
Age/Sex	**:-**	67 Y/M	**Sample Rvd At**	:-	27 Aug 2025 08:59 AM
Address	**:-**	323, SECTOR 9 PANCHKULA , Panchkula	**Report Gnt At**	:-	27 Aug 2025 04:25 PM
Mobile No	**:-**	9815311118	**Consultant**	**:-**	Dr AJAY GOYAL (DM(NEPHROLOGY))

Remarks :-

* END OF REPORT *

(Technologist)

Dr. Kavita Goyal
MD(Pathology)
HMC Reg. No. 14871

Note :- * If the result of the test is alarming or unexpected,the patient is contact the laboratory immediately
* The lab report is not the diagnosis,it represents only an opinion. Kindly Contact your doctor for interpretation diagnosis and treatment. Not Valid for Medico legal purpose *Report is Electronically validated.