

1 Gradient Drift Case

$$d\mathbf{x} = -\nabla\psi(\mathbf{x})dt + \sqrt{2\beta^{-1}}dw \quad (1)$$

$$\frac{\partial\rho}{\partial t} = \nabla \cdot (\rho\nabla\psi) + \beta^{-1}\Delta\rho \quad (2)$$

We claim that

$$\begin{aligned} F(\rho) &= \mathbb{E}_\rho[\psi + \beta^{-1}\log\rho] \\ &= \int \psi(\mathbf{x})\rho(\mathbf{x}) d\mathbf{x} + \beta^{-1} \int \rho(\mathbf{x}) \log\rho(\mathbf{x}) d\mathbf{x} \\ &= \beta^{-1} \int \rho \log\rho - \rho \log(\exp(-\beta\psi)) d\mathbf{x} \\ &= \beta^{-1} \int \rho \log\left(\frac{\rho}{\exp(-\beta\psi)}\right) d\mathbf{x} \\ &= \beta^{-1} \int \rho \log\left(\frac{\rho}{\frac{1}{Z}\exp(-\beta\psi)}\right) d\mathbf{x} + C \end{aligned}$$

is a Lyapunov functional along the trajectories of (??) and $Z = \int \exp(-\beta\psi)d\mathbf{x}$ is a normalization constant. If you have

$$\dot{\mathbf{x}} = \phi(\mathbf{x}) \quad (3)$$

$$\frac{dV}{dt} = \langle \nabla V, \dot{\mathbf{x}} \rangle = \langle \nabla V, \phi(\mathbf{x}) \rangle \leq 0 \quad (4)$$

$$\begin{aligned} \frac{dF}{dt} &= \left\langle \frac{\delta F}{\delta\rho}, \frac{\partial\rho}{\partial t} \right\rangle \\ &= \left\langle \frac{\delta F}{\delta\rho}, \nabla \cdot (\rho\nabla\psi) + \beta^{-1}\Delta\rho \right\rangle \\ &= \int \frac{\delta F}{\delta\rho} (\nabla \cdot (\rho\nabla\psi) + \beta^{-1}\Delta\rho) d\mathbf{x} \\ &= \int (\psi(\mathbf{x}) + \beta^{-1}(\log\rho(\mathbf{x}) + 1)) (\nabla \cdot (\rho\nabla\psi) + \beta^{-1}\Delta\rho) d\mathbf{x} \\ &= \int (\psi(\mathbf{x}) + \beta^{-1}(\log\rho(\mathbf{x}) + 1)) (\nabla \cdot (\rho(\nabla\psi + \beta^{-1}\nabla\log\rho))) d\mathbf{x} \\ &= - \int \langle \nabla(\psi(\mathbf{x}) + \beta^{-1}\log\rho(\mathbf{x})), \nabla\psi(\mathbf{x}) + \beta^{-1}\nabla\log\rho \rangle \rho d\mathbf{x} \\ &= - \int \|\nabla(\psi(\mathbf{x}) + \beta^{-1}\log\rho)\|^2 \rho(\mathbf{x}) d\mathbf{x} \\ &= -\mathbb{E}_\rho[\|\nabla\zeta\|^2] \leq 0 \end{aligned} \quad (5)$$

where $\zeta = \psi + \log \rho$.

We consider the system of 2nd-order SDEs given by

$$m_i \ddot{\theta}_i + \gamma_i \dot{\theta}_i = P_i - \sum_{j=1}^n K_{ij} \sin(\theta_i - \theta_j) + \sigma_i \times \text{stochastic forcing } i = 1, \dots, n.$$

We can transform the system into

$$\begin{aligned} d\boldsymbol{\theta} &= \boldsymbol{\omega} dt \\ \mathbf{M} d\boldsymbol{\omega} &= (-\Gamma \boldsymbol{\omega} - \nabla_{\boldsymbol{\theta}} V(\boldsymbol{\theta})) dt + \boldsymbol{\Sigma} d\mathbf{w} \end{aligned} \quad (6)$$

where $\boldsymbol{\theta} = (\theta_1, \theta_2, \dots, \theta_n)$, $\boldsymbol{\omega} = (\dot{\theta}_1, \dot{\theta}_2, \dots, \dot{\theta}_n)$, $\mathbf{M} = \text{diag}(m_1, m_2, \dots, m_n)$, $\Gamma = \text{diag}(\gamma_1, \gamma_2, \dots, \gamma_n)$, $\boldsymbol{\Sigma} = \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_n)$ and the potential function is given by

$$V(\boldsymbol{\theta}) := \sum_{i=1}^n P_i \theta_i + \sum_{(i,j) \in \mathcal{E}} k_{ij} (1 - \cos(\theta_i - \theta_j)) \quad (7)$$

we can rewrite as

$$d \underbrace{\begin{bmatrix} \boldsymbol{\theta} \\ \boldsymbol{\omega} \end{bmatrix}}_x = \underbrace{\begin{bmatrix} \boldsymbol{\omega} \\ -\mathbf{M}^{-1} \Gamma \boldsymbol{\omega} - \mathbf{M}^{-1} \nabla_{\boldsymbol{\theta}} V(\boldsymbol{\theta}) \end{bmatrix}}_f dt + \underbrace{\begin{bmatrix} \mathbf{0}_{n \times n} & \mathbf{0}_{n \times n} \\ \mathbf{0}_{n \times n} & \mathbf{M}^{-1} \boldsymbol{\Sigma} \end{bmatrix}}_g d\mathbf{w} \quad (8)$$

1.1 Fokker Planck

$$\begin{aligned} \frac{\partial \rho}{\partial t} &= -\nabla_{\boldsymbol{\theta}, \boldsymbol{\omega}} \cdot \left(\rho \begin{bmatrix} \boldsymbol{\omega} \\ -\mathbf{M}^{-1} \Gamma \boldsymbol{\omega} - \mathbf{M}^{-1} \nabla_{\boldsymbol{\theta}} V(\boldsymbol{\theta}) \end{bmatrix} \right) + \frac{1}{2} \sum_{i,j=1}^n \frac{\partial^2}{\partial \omega_i \partial \omega_j} (\rho \mathbf{M}^{-1} \boldsymbol{\Sigma} \boldsymbol{\Sigma}^\top \mathbf{M}^{-1})_{i,j} \\ &= -\nabla_{\boldsymbol{\theta}} \cdot (\rho \boldsymbol{\omega}) + \nabla_{\boldsymbol{\omega}} \cdot (\rho (\mathbf{M}^{-1} \Gamma \boldsymbol{\omega} + \mathbf{M}^{-1} \nabla_{\boldsymbol{\theta}} V(\boldsymbol{\theta}))) + \nabla_{\boldsymbol{\omega}} \cdot (\rho \mathbf{D} \nabla_{\boldsymbol{\omega}} \log \rho) \\ &= -\langle \boldsymbol{\omega}, \nabla_{\boldsymbol{\theta}} \rho \rangle + \nabla_{\boldsymbol{\omega}} \cdot (\rho (\mathbf{M}^{-1} \Gamma \boldsymbol{\omega} + \mathbf{M}^{-1} \nabla_{\boldsymbol{\theta}} V(\boldsymbol{\theta}) + \mathbf{D} \nabla_{\boldsymbol{\omega}} \log \rho)) \end{aligned} \quad (9)$$

where $\mathbf{D} = \frac{1}{2} \mathbf{M}^{-1} \boldsymbol{\Sigma} \boldsymbol{\Sigma}^\top \mathbf{M}^{-1}$.

Theorem 1. (*Ito's Lemma for Multi-Dimensional Processes*) Let

$$d\mathbf{X}_t = \mathbf{f}(\mathbf{X}_t, t) dt + \mathbf{g}(\mathbf{X}_t, t) d\mathbf{w}_t \quad (10)$$

with $\mathbf{X}_t, \mathbf{w}_t, \mathbf{f}(\cdot, \cdot) \in \mathbb{R}^n$ and $\mathbf{g}(\cdot) \in \mathbb{R}^{n \times n}$. Then for a given transformation

$$\mathbf{Z}_t = \boldsymbol{\psi}(\mathbf{X}_t, t) = [\psi_1(\mathbf{X}_t, t), \dots, \psi_n(\mathbf{X}_t, t)] \quad (11)$$

where ψ is a function from $\mathbb{R}^n \times [0, \infty]$ into \mathbb{R}^n , then \mathbf{Z}_t is again an Ito process given by

$$\begin{aligned} dZ_{k,t} &= \frac{\partial \psi_k}{\partial t}(\mathbf{X}_t, t) dt + \sum_{i=1}^n \frac{\partial \psi_k}{\partial x_i}(\mathbf{X}_t, t) dX_{i,t} \\ &\quad + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2 \psi_k}{\partial x_i \partial x_j}(\mathbf{X}_t, t) dX_{j,t} dX_{i,t} \end{aligned} \quad (12)$$

Theorem 2. Let $n = 2d$ and consider the transformation defined by

$$\mathbf{Z}_t = \psi(\mathbf{X}_t) = \begin{bmatrix} \boldsymbol{\xi} \\ \boldsymbol{\eta} \end{bmatrix} := \begin{bmatrix} \text{diag}(\mathbf{m} \oslash \boldsymbol{\sigma}) & 0 \\ 0 & \text{diag}(\mathbf{m} \oslash \boldsymbol{\sigma}) \end{bmatrix} \begin{bmatrix} \boldsymbol{\theta} \\ \boldsymbol{\omega} \end{bmatrix} \quad (13)$$

i.e.

$$\begin{aligned} \mathbf{Z}_t &= [\psi_1(X_{1,t}), \dots, \psi_d(X_{d,t}), \psi_{d+1}(X_{d+1,t}), \dots, \psi_{2d}(X_{2d,t})]^\top \\ &= \left[\frac{m_1}{\sigma_1} \theta_{1,t}, \dots, \frac{m_d}{\sigma_d} \theta_{d,t}, \dots, \frac{m_1}{\sigma_1} \omega_{1,t}, \dots, \frac{m_d}{\sigma_d} \omega_{d,t} \right]^\top \end{aligned} \quad (14)$$

Proof. Since the transformation (??) is a linear transformation and doesn't depend on t , the first and third term in (??) vanish so we are left with

$$\begin{aligned} dZ_{k,t} &= \sum_{i=1}^{2d} \frac{\partial \psi_k}{\partial x_i}(\mathbf{X}_t) dX_{i,t} \\ &= \sum_{i=1}^{2d} \frac{\partial \psi_k}{\partial x_i}(X_{k,t}) dX_{i,t} \\ &= \frac{\partial \psi_k}{\partial x_k} dX_{k,t} \\ &= \begin{cases} \frac{m_k}{\sigma_k} d\theta_{k,t} & 1 \leq k \leq d \\ \frac{\sigma_k}{m_k} d\omega_{k,t} & n+1 \leq k \leq 2d \end{cases} \\ &= \begin{cases} \frac{m_k}{\sigma_k} \omega_{k,t} dt & 1 \leq k \leq d \\ \frac{m_k}{\sigma_k} \left(-\frac{\gamma_k}{m_k} \omega_{k,t} - \frac{P_k}{\sigma_k} - \sum_{j=1}^d \frac{K_{k,j}}{m_k} \sin(\theta_{k,t} - \theta_{j,t}) + \frac{\sigma_k}{m_k} dw_{k,t} \right) & n+1 \leq k \leq 2d \end{cases} \\ &= \begin{cases} \eta_{k,t} dt & 1 \leq k \leq d \\ -\frac{\gamma_k}{m_k} \eta_{k,t} - \frac{P_k}{\sigma_k} - \sum_{j=1}^d \frac{K_{k,j}}{\sigma_k} \sin\left(\frac{\sigma_k}{m_k} \xi_{k,t} - \frac{\sigma_j}{m_j} \xi_{j,t}\right) + dw_{k,t} & n+1 \leq k \leq 2d \end{cases} \end{aligned}$$

Writing in vector form, we get

$$\begin{bmatrix} d\boldsymbol{\xi} \\ d\boldsymbol{\eta} \end{bmatrix} = \begin{bmatrix} \boldsymbol{\eta} \\ -\nabla_{\boldsymbol{\eta}} F(\boldsymbol{\eta}) - \nabla_{\boldsymbol{\xi}} U(\boldsymbol{\xi}) \end{bmatrix} dt + \begin{bmatrix} \mathbf{0}_{d \times d} & \mathbf{0}_{d \times d} \\ \mathbf{0}_{d \times d} & \mathbf{I}_{d \times d} \end{bmatrix} d\mathbf{w} \quad (15)$$

where $F = \frac{1}{2}\langle \boldsymbol{\eta}, \text{diag}(\boldsymbol{\gamma} \odot \boldsymbol{m})\boldsymbol{\eta} \rangle$ and the potential function in $\boldsymbol{\xi}$ is given by

$$U(\boldsymbol{\xi}) = \sum_{i=1}^{2d} \frac{P_i}{\sigma_i} \xi_i + \sum_{(i,j) \in \mathcal{E}} K_{i,j} \frac{m_i}{\sigma_i^2} \left(1 - \cos \left(\frac{\sigma_i}{m_i} \xi_i - \frac{\sigma_j}{m_j} \xi_j \right) \right) \quad (16)$$

□

The Fokker-Planck equation of this SDE is

$$\frac{\partial \rho}{\partial t} = -\langle \boldsymbol{\eta}, \nabla_{\boldsymbol{\xi}} \rho \rangle + \nabla_{\boldsymbol{\eta}} \cdot ((\nabla_{\boldsymbol{\eta}} F(\boldsymbol{\eta}) + \nabla_{\boldsymbol{\xi}} U(\boldsymbol{\xi})) \rho) + \frac{1}{2} \Delta_{\boldsymbol{\eta}} \rho \quad (17)$$

Remark 1. Given a random vector X , who joint pdf is given by $f_X(x)$ and if $H : \mathbb{R}^n \mapsto \mathbb{R}^n$ is a $1-1$ differentiable function, then the random vector $Y = H(X)$ also has a density function given by

$$f_Y(y) = \frac{f_X(x)}{|\det \nabla H(x)|}, \quad x \in H^{-1}(y) \quad (18)$$