



Gajendra Purohit

Legend in CSIR-UGC NET & IIT-JAM

~ Unlock Code : GPSIR – PhD, CSIR NET (Maths) | Youtuber(800K+165K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

→ **TOP EDUCATOR ON UNACADEMY
FOR CSIR NET & IIT JAM**

YouTuber with 800K Subscribers

→ **AUTHOR OF BEST SELLER BOOK
FOR CSIR NET & IIT JAM**

**Get
10% Off**

Referral Code : GP SIR



CSIR

We heard you and upgraded the
Ask a Doubt feature to suit your needs!

Unacademy Ask a Doubt



The Ask a Doubt feature is now available on the Website and the Learner App . Getting your doubts clarified is easier and faster than before!

[ASK A DOUBT](#)

Detailed Course on Function of One and Several Variable IIT JAM' 23

 Batch starting from May 25, 2022



Scan QR code
to join class

Gajendra Purohit

[Enroll now](#)

Use code GPSIR
for 10% off





Detail Course on Numerical Analysis For CSIR NET 22

📅 May 10, 2022 | 4 PM

Gajendra Purohit



Enroll Now



Detail Course on Real Analysis For CSIR NET 22

📅 May 4, 2022 | 9 PM

Gajendra Purohit



Enroll Now

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months

SAVE 47%

₹908 /mo

Total ₹21,780

You get 6 months extra for free

Offer expires 15 Jun 2022

12 months

SAVE 54%

₹1,248 /mo

Total ₹14,974

You get 4 months extra for free

Offer expires 15 Jun 2022

9 months

SAVE 45%

₹1,497 /mo

Total ₹13,475

6 months

SAVE 25%

₹2,042 /mo

Total ₹12,252

3 months

SAVE 17%

₹2,269 /mo

Total ₹6,807

1 month

₹2,723 /mo

Total ₹2,723

To be paid as a one-time payment



Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months

SAVE 47%

₹817 /mo

₹21,780 ₹19,602

You get 6 months extra for free

Offer expires 15 Jun 2022

12 months

SAVE 54%

₹1,123 /mo

₹14,974 ₹13,477

You get 4 months extra for free

Offer expires 15 Jun 2022

9 months

SAVE 45%

₹1,348 /mo

₹13,475 ₹12,328

6 months

SAVE 25%

₹1,838 /mo

₹12,252 ₹11,027

3 months

SAVE 17%

₹2,042 /mo

₹6,807 ₹6,126

1 month

₹2,451 /mo




GPSIR

Awesome! You got 10% off!

Proceed to pay

After Using Referral Code





Gajendra Purohit

Legend in CSIR-UGC NET & IIT JAM

~ Unlock Code : GPSIR – PhD, CSIR NET (Maths) | Youtuber (1000k+230K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

77M Watch mins

4M Watch mins (last 30 days)

54K Followers

3K Dedications

USE
CODE
GPSIR

GET
10% OFF


**TOP EDUCATOR
ON UNACADEMY**

FOR CSIR-NET & IIT JAM

1M SUBSCRIBERS ON YOUTUBE

**AUTHOR OF BEST SELLER BOOK
OF CSIR-NET & IIT JAM**





Prices have dropped!

Get an Unacademy CSIR-UGC NET Subscription and save up to 21%*

Plus

Duration	Current Price	After 12% Price Slash	After 10% Unlock code	What you Save
24 months	₹ 25,410	₹ 22,361	₹ 20,125	₹ 5,285 (21%)
12 months	₹ 18,423	₹ 16,212	₹ 14,591	₹ 3,832 (21%)
6 months	₹ 14,738	₹ 12,969	₹ 11,672	₹ 3,066 (21%)

Iconic


Duration	Current Price	After 12% Price Slash	After 10% Unlock code	What you Save
24 months	₹ 58,273	₹ 51,280	₹ 46,152	₹ 12,121 (21%)
12 months	₹ 33,858	₹ 29,795	₹ 26,816	₹ 7,042 (21%)
6 months	₹ 23,694	₹ 20,851	₹ 18,766	₹ 4,928 (21%)



Subscribe Now

USE CODE
GPSIR





Save up to 20% on your IIT JAM Subscriptions!*

Now available at the lowest prices!*

Plus

Duration	Current Price	Price Drop	What you Pay	What you save
3 Months	₹ 6,807	₹ 1,361	₹ 5,446	20%

Hurry, offer valid till November 24

[Subscribe Now](#)

Use code
GPSIR

Call **8585858585**
for more details

*T&C apply, as available on the platform.





DETAILED COURSE ON **THEORY OF ESTIMATION** FOR CSIR-NET 2023

Gajendra Purohit

22 November

Enroll Now



DETAILED COURSE 2.0 ON **INTEGRAL CALCULUS** FOR IIT-JAM 2023

1 December | 9:00 AM

Gajendra Purohit

Enroll Now

Use code GPSIR for 10% off



*T&C apply, as available on the platform.



FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

Directional Derivative :

Let f be a function of n -variables defined on D and let $u = (u_1, u_2, \dots, u_n)$ be a unit vector then the directional derivative of f at $\alpha = (x_1, x_2, \dots, x_n)$ in the direction of u denoted by $D_u f(x_1, x_2, \dots, x_n)$ is defined by

$$D_u f(\alpha) = \lim_{h \rightarrow 0} \frac{f(\alpha + hu) - f(\alpha)}{h} \text{ if limit exist,}$$

Where $\alpha = (x_1, x_2, \dots, x_n)$, $u = (u_1, u_2, \dots, u_n)$

$$\alpha + hu = (x_1 + hu_1, x_2 + hu_2, \dots, x_n + hu_n)$$

Note :

- (1) If direction is not unit vector, then firstly we convert into unit vector.
- (2) If direction is not mention in question then we take arbitrary direction as

$$u = \cos\theta \hat{i} + \sin\theta \hat{j}$$

‘Some important result :

- (1) If $f(x, y)$ is differentiable at (a, b) then all directional derivative exist at (a, b) but converse need not be true.
- (2) If $f(x, y)$ is continuous at (a, b) then all directional derivative of f at (a, b) but converse need not be true.
- (3) If partial derivative exist and continuous at (a, b) then f is differentiable at (a, b)

Q1. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function. Then which of the following statements is/are TRUE? **IIT-JAM 2017**

- (a) If f is differentiable at $(0, 0)$, then all directional derivatives of f exists at $(0, 0)$
- (b) If all directional of f exists at $(0, 0)$, then f is differentiable at $(0, 0)$
- (c) If all directional derivative of f exists at $(0, 0)$, then f is continuous at $(0, 0)$
- (d) If the partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ exists and are continuous in a disc centred at $(0,0)$ then f is differentiable at $(0,0)$

Q2.

$$\text{Let } f(x, y) = \begin{cases} \frac{x^3 + y^3}{x^2 - y^2}, & x^2 - y^2 \neq 0 \\ 0, & x^2 - y^2 = 0 \end{cases}$$

$$\alpha + h\gamma = (0, 0 + h) \rightarrow (4, 7)$$

$$\alpha = (0, 0)$$

$$\gamma = \left(\frac{4}{5}, \frac{7}{5} \right)$$

Then the directional derivative of f at $(0, 0)$ in the

direction of $\frac{4}{5}\hat{i} + \frac{3}{5}\hat{j}$ is

IIT-JAM 2019

(a) 2.5 to 2.7

(b) 2.4 to 2.8

(c) 2.6 to 2.6

(d) 2.8 to 2.9

$$= \lim_{h \rightarrow 0} \frac{(4h\hat{i})^3 + (3h\hat{j})^3}{(4h\hat{i})^2 - (3h\hat{j})^2} h$$

$$= \frac{(4^3\hat{i}^3 + 3^3\hat{j}^3)}{(4^2\hat{i}^2 - 3^2\hat{j}^2)} = \frac{64 + 27}{5(16 - 9)} = \frac{91}{5 \times 7} = \frac{13}{5}$$

$$D_u(f) = \lim_{h \rightarrow 0} \frac{f(\alpha + h\gamma) - f(\alpha)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f((0, 0) + h(4, 7)) - f(0, 0)}{h}$$

Q3. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} \left(1 + \frac{x}{y}\right)^2, & y \neq 0 \\ 0, & y = 0 \end{cases} \text{ If the directional derivative } D_u(\mathbf{r}) = \lim_{h \rightarrow 0} \frac{f(r+hu) - f(r)}{h}$$

derivative of f at $(0, 0)$ exists along the direction

$\cos \alpha \hat{i} + \sin \alpha \hat{j}$, where $\sin \alpha \neq 0$, then the value of $\cot \alpha$ is

IIT-JAM 2015

- (a) 1
- (b) -1
- (c) 2
- (d) 0

$$\boxed{1 + (\cot \alpha)^2 = 0}$$

$(\cot \alpha)^2 = 1$

$$= \lim_{h \rightarrow 0} \left(1 + \frac{K(s)h}{K(h)^2} \right)$$

$$= \lim_{h \rightarrow 0} \left(1 + (92) \right) \quad \textcircled{C}$$

COMPLETE COURSE ON

MATHEMATICS

FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINEAR ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



**After Using
My Referral
Code**



GPSIR

Awesome! You get 10% off

Proceed to pay

Maxima and minima of two variables function

1. Maxima of function :

Let (a, b) be a point in the domain of $f(x, y)$, then $f(x, y)$ has a maxima at (a, b) iff $(a, b) \geq f(x, y)$; for all $(x, y) \in \text{domain}$

$$\begin{aligned}f(x,y) &= \sqrt{x^2+y^2} \\D &= \{(x,y) | x^2+y^2 \leq 1\} \\f'(0,0) &= \infty \\f(1,0) &= 1 \\f(-1,0) &= 1\end{aligned}$$

2.

Minima of function :

Let (a, b) be a point in the domain of $f(x, y)$, then $f(x, y)$ has a minima at (a, b) if $f(a, b) \leq f(x, y)$; for all $(x, y) \in$ domain

3.

Extremum point :

$f(x, y)$ is said to have an extremum at a point (a, b) if it has either a maximum and minimum at (a, b) .

4. Critical point :

A point (a, b) is said to be a critical point for the function $f(x, y)$ if $f_x(a, b) = 0$ & $f_y(a, b) = 0$

Note : A critical point is an extremum point of function but converse may not true.

$$\textcircled{0,0}$$

$$\textcircled{1,0}$$

$$\textcircled{0,1}$$


$$f = x^2 + y^2$$
$$\frac{\partial f}{\partial x} = 2x = 0$$
$$\frac{\partial f}{\partial y} = 2y = 0$$

$$n = \infty$$
$$T = \infty$$

Procedure of finding maxima & minima

Let $f(x, y)$ be a function of two variables and $r = f_{xx}(a, b)$,
 $s = f_{xy}(a, b)$, $t = f_{yy}(a, b)$, then

- (i) f has a minima at (a, b) if $rt - s^2 > 0$ & $r > 0$
- (ii) f has a maxima at (a, b) if $rt - s^2 > 0$ & $r < 0$
- (iii) f has neither maxima nor minima at (a, b) if $rt - s^2 < 0$
- (iv) f may or may not have an extrema if $rt - s^2 = 0$



$$\begin{aligned}r &= \frac{\partial^2 f}{\partial x^2} \\s &= \frac{\partial^2 f}{\partial x \partial y} \\t &= \frac{\partial^2 f}{\partial y^2}\end{aligned}$$

$$D = \frac{\partial^2 f}{\partial x^2} \cdot \frac{\partial^2 f}{\partial y^2} - \left(\frac{\partial^2 f}{\partial x \partial y}\right)^2$$

$$\begin{aligned}r &= \frac{\partial^2 f}{\partial x^2} \\s &= \frac{\partial^2 f}{\partial x \partial y} \\t &= \frac{\partial^2 f}{\partial y^2}\end{aligned}$$

$$D = \frac{\partial^2 f}{\partial x^2} \cdot \frac{\partial^2 f}{\partial y^2} - \left(\frac{\partial^2 f}{\partial x \partial y}\right)^2$$

$$f = 1 - \frac{y^2 - y^2}{y^2}$$

$$\frac{\partial f}{\partial y} = -\frac{2y}{y^2} \Rightarrow$$

$$\frac{\partial f}{\partial y} = -\frac{2y}{y^2} \Rightarrow$$

$$(0, 0)$$

$$\gamma = \frac{\partial^2 f}{\partial y^2} = -2$$

$$\delta = \frac{\partial^2 f}{\partial x \partial y} = 0$$

$$\kappa = \frac{\partial^2 f}{\partial x^2} = -1$$

$$\begin{cases} y=0 \\ y=\infty \end{cases}$$

$$\gamma L - \gamma U = 4 - 0 > 0$$

$$\gamma = -2 < 0$$

maximum

(0, 0) true IP

maximum

$$\max f = 1 - 0 - 0$$

$$= 1$$

$$z = \frac{x^3 - 2xy^2}{x^2}$$

$$\frac{\partial z}{\partial x} = 3x^2 - 2y^2 = 0 \Rightarrow$$

$$\frac{\partial z}{\partial y} = -4xy = 0$$

0:5

$$y = \frac{\partial z}{\partial x} = 3$$

$$P = \frac{\partial z}{\partial y} = -4x$$

$$t = \frac{\partial P}{\partial x} = -4$$

$$x = t = y$$

$$x = 0, y = 0$$

$$x = 0, y = 0$$

at (0,0)

$$y = 0$$

$$P = 0$$

$$t = 0$$

$$\begin{aligned} f(0,0) &= 0 \\ f'(1,0) &= 1 \\ f''(1) &= -2 \\ y + -j^2 &= 0 \end{aligned}$$

Saddle Point :

A point (a, b) is said to be saddle point if it is neither maxima nor minima.

Q.1. Consider the function $f(x, y) = 5 - 4 \sin x + y^2$ for $0 \leq x \leq 2\pi$ and $y \in \mathbb{R}$. The set of critical points of $f(x, y)$ consists of

IT JAM – 2017

- (a) a point of local maxima and a point of local minima
- (b) a point of local maxima and a saddle point
- (c) a point of local maxima, a point of local minima and a saddle point
- (d) a point of local minima and a saddle point.

at $(\pi, 0)$

$y=4, s=0, t=-1$

$y+s-t=8>0$
min.

at $(\pi, 0)$

$y=-1, s=0, t=1$

$y+s-t=-8<0$

$y+s-t=-8<0$

$$\frac{\partial f}{\partial x} = -4 \cos x = 0$$

$$\frac{\partial f}{\partial y} = 2y = 0$$

$$\frac{1}{4}, \frac{3\pi}{2}$$

$$(\pi_2, 0), (\pi_2, 0)$$

$$\gamma = \frac{\partial^2 f}{\partial x^2} = 4 \sin x$$

$$\delta = \frac{\partial^2 f}{\partial x \partial y} = 0$$

$$\tau = \frac{\partial^2 f}{\partial y^2} = 2$$

Q.2 $f(x, y) = x^3 - y^3 - 3x^2 + 3y^2 + 7, (x, y) \in \mathbb{R}^2$. Then the local minimum(m) and local Maximum (M) of f are given by IIT JAM – 2017

(a) $m = 3, M = 7$

(b) $m = 4, M = 11$

(c) $m = 7, M = 11$

(d) $m = 3, M = 11$

$$(0, 0)$$

$$\gamma = -1$$

$$\delta = 0$$

$$\lambda = 1$$

$$\gamma + \delta = -1 < 0$$

$$(0, 2)$$

$$\gamma = -6 < 0$$

$$\delta = 0$$

$$\lambda = -6$$

$$\gamma + \delta = -6 > 0$$

$$\max f = -8 + 12 + 7 \\ m = 11$$

$$(2, 0)$$

$$\gamma = 6$$

$$\delta = 0$$

$$\lambda = 6$$

$$\gamma + \delta = 6 > 0$$

$$\min f = 8 - 12 + 7$$

$$m = 3$$

$$\frac{\partial f}{\partial x} = 3x^2 - 6x = 0$$

$$\frac{\partial f}{\partial y} = -3y + 6 = 0$$

$$y = 2$$

$$\gamma = \frac{\partial y}{\partial x}, \quad \gamma_{yy} = (y-6)$$

$$\gamma_{yy} = 0$$

$$\lambda = \frac{\partial y}{\partial x} - \gamma + 1$$

Q3. Let f given by $f(x, y) = x^2 + xy + y^2 - x - 100$. Find the points of local maxima and local minima (if any) of f .

IIT JAM - 2011

(a) Only $\left(\frac{2}{3}, -\frac{1}{3}\right)$

(c) $\left(\frac{2}{3}, \frac{1}{3}\right)$

(x_3, y_3)

(b) Only $\left(-\frac{2}{3}, -\frac{1}{3}\right)$

(d) None of these

$$\frac{\partial f}{\partial x} = 2x + y - 1 = 0$$

$$\frac{\partial f}{\partial y} = x + 2y = 0$$

$$x = -2y$$

$$\gamma = \frac{\partial^2 f}{\partial x^2} =$$

$$\rho = \frac{\partial^2 f}{\partial x \partial y} = 1 \quad -4y + 1 = 1 \Rightarrow$$

$$\delta = \frac{\partial^2 f}{\partial y^2} =$$

$$-3 = 1 \quad \begin{cases} y = \frac{1}{3} \\ y = -\frac{1}{3} \end{cases}$$

$$x + y - 4 - 1 = 3 > 0$$

$$\delta = 2 > 0$$

Q4. Let $f(x, y) = x^3 + y^3 + 3xy$ is a function of two variables. Which of the following is true?

ITT - JAM 2012

(a) maxima at $(-1, -1)$

(c) maxima at $(0, 0)$

(d) saddle point at $(-1, -1)$

$\Delta < 0$

$r = \infty$

$P = 3$

$K = \infty$

$$\lambda - \mu^2 = -1 < 0$$

$\Delta < 0$

$$\mu = -\lambda$$

$$\beta = \lambda$$

$$k = -1$$

$$\lambda - \mu^2 = \lambda > 0$$

$$\gamma = \frac{\partial f}{\partial x} = 3x + 3y$$

$$\beta = \frac{\partial f}{\partial y} = 3y + 3x$$

$$k = -\frac{\partial f}{\partial y} = -3y$$

$$\frac{\partial f}{\partial x} = 3x^2 + 3y = 0$$

$$\frac{\partial f}{\partial y} = 3y^2 + 3x = 0$$

$$3y^2 = -3x$$

$$3y^2 = -3x$$

$$\lambda = \pm \gamma$$

Q5. Let $f(x, y) = x^4 - 2x^2y^2 + y^4 + x^2 - 6xy + 9y^2$

then which of the following is true

- (a) f has a minima (global) at (0,1)
- (b) f has a minima (global) at (0,0)
- (c) f has a minima (global) at (2,0)
- (d) None of these

Q6. The critical point of $f(x, y) = x^3 + y^2 - 12x - 6y + 40$
are **IIT JAM – 2006**

- (a) saddle point (b) maxima
- (c) minima (d) none of these

Q7. Let $f(x, y) = 4xy - x^3y - xy^3$ for $(x, y) \in \mathbb{R}^2$. The value of f at local minimum in the rectangular region

$$R = \left\{ (x, y) \in \mathbb{R}^2 \mid |x| < \frac{3}{2}, |y| < \frac{3}{2} \right\}$$
 is

- (a) -2
- (b) -3
- (c) -7/8
- (d) 0

hit

FOUNDATION COURSE OF

MATHEMATICS

FOR CSIR-NET



DETAILED COURSE ON **THEORY OF ESTIMATION** FOR CSIR-NET 2023

Gajendra Purohit

22 November

Enroll Now



DETAILED COURSE 2.0 ON **INTEGRAL CALCULUS** FOR IIT-JAM 2023

1 December | 9:00 AM

Gajendra Purohit

Enroll Now

Use code GPSIR for 10% off



*T&C apply, as available on the platform.



Educator Profile



Gajendra Purohit

#5 Educator in CSIR-UGC NET

[Follow](#)

Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



CSIR-UGC NET

[SEE ALL](#)

HINDI MATHEMATICAL SCIENCES

Course on Linear Algebra, Partial Diff. Equation & Calculus

Starts on Mar 1, 2021 • 24 lessons

Gajendra Purohit

HINDI MATHEMATICAL SCIENCES

Course on Complex Analysis & Integral Equation

Starts on Jan 14, 2021 • 16 lessons

Gajendra Purohit

HINDI MATHEMATICAL SCIENCES

Foundation Course on Mathematics for CSIR 2021

Starts on Dec 7, 2020 • 20 lessons

Gajendra Purohit

Educator highlights

- 📍 Works at Pacific Science College
- 📍 Studied at M.Sc., NET, PhD(Algebra), MBA(Finance), BEd
- 📍 PhD, NET | Plus Educator For CSIR NET | Youtuber (260K+Subs.) | Director Pacific Science College |
- 📍 Lives in Udaipur, Rajasthan, India
- 📍 Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR

Maxima and minima of two variables function

1. Maxima of function :

Let (a, b) be a point in the domain of $f(x, y)$, then $f(x, y)$ has a maxima at (a, b) iff $(a, b) \geq f(x, y)$; for all $(x, y) \in \text{domain}$

2.

Minima of function :

Let (a, b) be a point in the domain of $f(x, y)$, then $f(x, y)$ has a minima at (a, b) if $f(a, b) \leq f(x, y)$; for all $(x, y) \in$ domain

3.

Extremum point :

$f(x, y)$ is said to have an extremum at a point (a, b) if it has either a maximum and minimum at (a, b) .

4. Critical point :

A point (a, b) is said to be a critical point for the function $f(x, y)$ if $f_x(a, b) = 0$ & $f_y(a, b) = 0$

Note : A critical point is an extremum point of function but converse may not true.

Procedure of finding maxima & minima

Let $f(x, y)$ be a function of two variables and $r = f_{xx}(a, b)$,
 $s = f_{xy}(a, b)$, $t = f_{yy}(a, b)$, then

- (i) f has a minima at (a, b) if $rt - s^2 > 0$ & $r > 0$
- (ii) f has a maxima at (a, b) if $rt - s^2 > 0$ & $r < 0$
- (iii) f has neither maxima nor minima at (a, b) if $rt - s^2 < 0$
- (iv) f may or may not have an extrema if $rt - s^2 = 0$

Saddle Point :

A point (a, b) is said to be saddle point if it is neither maxima nor minima.

Q.1. Consider the function $f(x, y) = 5 - 4 \sin x + y^2$ for

$0 < x < 2\pi$ and $y \in \mathbb{R}$. The set of critical points of

$f(x, y)$ consists of

IT JAM – 2017

- (a) a point of local maxima and a point of local minima
- (b) a point of local maxima and a saddle point
- (c) a point of local maxima, a point of local minima and a saddle point
- (d) a point of local minima and a saddle point.

COMPLETE COURSE ON

MATHEMATICS

FOR IIT-JAM 2022

TARGETED AUDIENCE

- IIT-JAM
- M.Sc. Entrance Exam

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINEAR ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months

SAVE 47%

₹908 /mo

Total ₹21,780

You get 6 months extra for free

Offer expires 15 Jun 2022

12 months

SAVE 54%

₹1,248 /mo

Total ₹14,974

You get 4 months extra for free

Offer expires 15 Jun 2022

9 months

SAVE 45%

₹1,497 /mo

Total ₹13,475

6 months

SAVE 25%

₹2,042 /mo

Total ₹12,252

3 months

SAVE 17%

₹2,269 /mo

Total ₹6,807

1 month

₹2,723 /mo

Total ₹2,723

To be paid as a one-time payment



Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months

SAVE 47%

₹817 /mo

₹21,780 ₹19,602

You get 6 months extra for free

Offer expires 15 Jun 2022

12 months

SAVE 54%

₹1,123 /mo

₹14,974 ₹13,877

You get 4 months extra for free

Offer expires 15 Jun 2022

9 months

SAVE 45%

₹1,348 /mo

₹13,475 ₹12,328

6 months

SAVE 25%

₹1,838 /mo

₹12,252 ₹11,027

3 months

SAVE 17%

₹2,042 /mo

₹6,807 ₹6,126


1 month

₹2,451 /mo

Awesome! You got 10% off!

GPSIR

Proceed to pay



After Using Referral Code

Q.2 $f(x, y) = x^3 - y^3 - 3x^2 + 3y^2 + 7, (x, y) \in \mathbb{R}^2$. Then the local minimum(m) and local Maximum (M) of f are given by **IT JAM – 2017**

- (a) $m = 3, M = 7$
- (b) $m = 4, M = 11$
- (c) $m = 7, M = 11$
- (d) $m = 3, M = 11$

Q3. Let f given by $f(x, y) = x^2 + xy + y^2 - x - 100$. Find the points of local maxima and local minima (if any) of f .

IIT JAM – 2011

- (a) Only $\left(\frac{2}{3}, -\frac{1}{3}\right)$
- (b) Only $\left(-\frac{2}{3}, -\frac{1}{3}\right)$
- (c) $\left(\frac{2}{3}, \frac{1}{3}\right)$
- (d) None of these

Q4. Let $f(x, y) = x^3 + y^3 + 3xy$ is a function of two variables. Which of the following is true?

ITT – JAM 2012

- (a) maxima at $(-1, -1)$
- (b) minima at $(-1, -1)$
- (c) maxima at $(0, 0)$
- (d) saddle point at $(-1, -1)$

Q5. Let $f(x, y) = x^4 - 2x^2y^2 + y^4 + x^2 - 6xy + 9y^2$

then which of the following is true

- (a) f has a minima (global) at (0,1)
- (b) f has a minima (global) at (0,0)
- (c) f has a minima (global) at (2,0)
- (d) None of these

Q6. The critical point of $f(x, y) = x^3 + y^2 - 12x - 6y + 40$
are **IIT JAM – 2006**

- (a) saddle point (b) maxima
- (c) minima (d) none of these

Q7. Let $f(x, y) = 4xy - x^3y - xy^3$ for $(x, y) \in \mathbb{R}^2$. The value of f at local minimum in the rectangular region

$$R = \left\{ (x, y) \in \mathbb{R}^2 \mid |x| < \frac{3}{2}, |y| < \frac{3}{2} \right\}$$
 is

- (a) -2
- (b) -3
- (c) -7/8
- (d) 0

hit

CSIR

We heard you and upgraded the
Ask a Doubt feature to suit your needs!

Unacademy Ask a Doubt



The Ask a Doubt feature is now available on the Website and the Learner App . Getting your doubts clarified is easier and faster than before!

[ASK A DOUBT](#)

Detailed Course on Function of One and Several Variable IIT JAM' 23

 Batch starting from May 25, 2022



Scan QR code
to join class

Gajendra Purohit

[Enroll now](#)

Use code GPSIR
for 10% off





Detail Course on Numerical Analysis For CSIR NET 22

📅 May 10, 2022 | 4 PM

Gajendra Purohit



Enroll Now



Detail Course on Real Analysis For CSIR NET 22

📅 May 4, 2022 | 9 PM

Gajendra Purohit



Enroll Now

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months

SAVE 47%

₹908 /mo

Total ₹21,780

You get 6 months extra for free

Offer expires 15 Jun 2022

12 months

SAVE 54%

₹1,248 /mo

Total ₹14,974

You get 4 months extra for free

Offer expires 15 Jun 2022

9 months

SAVE 45%

₹1,497 /mo

Total ₹13,475

6 months

SAVE 25%

₹2,042 /mo

Total ₹12,252

3 months

SAVE 17%

₹2,269 /mo

Total ₹6,807

1 month

₹2,723 /mo

Total ₹2,723

To be paid as a one-time payment



Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months

SAVE 47%

₹817 /mo

₹21,780 ₹19,602

You get 6 months extra for free

Offer expires 15 Jun 2022

12 months

SAVE 54%

₹1,123 /mo

₹14,974 ₹13,877

You get 6 months extra for free

Offer expires 15 Jun 2022

9 months

SAVE 45%

₹1,348 /mo

₹13,475 ₹12,328

6 months

SAVE 25%

₹1,838 /mo

₹12,252 ₹11,027

3 months

SAVE 17%

₹2,042 /mo

₹6,807 ₹6,126


1 month

₹2,451 /mo

GPSIR

Proceed to pay

Awesome! You got 10% off!



After Using Referral Code

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR