

Detailed Course 2.0 on Function of One and Several Variable - IIT JAM, 23



Gajendra Purohit



Legend in CSIR-UGC NET & IIT-JAM

- Unlock Code: GPSIR - PhD, CSIR NET (Maths) | Youtuber(800K+165K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

TOP EDUCATOR ON UNACADEMY FOR CSIR NET & IIT JAM

YouTuber with 800K Subscribers

AUTHOR OF BEST SELLER BOOK FOR CSIR NET & IIT JAM

> Get 10% Off

Referral Code: GP SIR





Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

November 3

Enroll Now





Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26 9:00 AM

Gajendra Purohit

Enroll Now

Use code GPSIR for 10% off





FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months	₹ 908 / mo
Save 67%	Total ₹ 21,780
You get 6 months extra for free	Offer expires 15 Jun 2022

You g	et 6 months extra for free	Offer expires 15 Jun 2022
Sa	ve 54%	Total ₹ 14,974
Ø 12	months	₹ 1,248 / mo

9 months	₹ 1,497 / mo
Save 45%	Total ₹ 13,475

6 months	₹ 2,042 / mo
Save 25%	Total ₹ 12,252

₹ 2,269 / mo
Total ₹ 6,807

1 month	₹2,723 / mo
	Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay



After Using My Referral Code

No cost EMI available on 6 months & above subscription plans

You get 6 months extra for free	Offer expires 15 Jun 2022
Save 67%	₹ 21,700 ₹ 19,602
24 months	₹ 817 / mo

2 12 months	₹ 1,123 / mo
Save 54%	₹ 14,974 ₹ 13,477
You get 6 months extra for free	Offer expires 15 Jun 2022

9 months	₹ 1,348 / mo
Save 45%	₹ 13,475 ₹ 12,128

6 months	₹ 1,838 / mo
Save 25%	₹-12,252 ₹ 11,027


3 months	₹ 2,042 / mo
Save 17%	₹ -6,807 ₹ 6,126



Proceed to pay

Mean Value Theorem:

- (1) Rolle's Theorem: Let f be a function defined on [a, b] s.t.
 - (a) f is continuous on [a, b]
 - (b) f is differentiable on (a, b)
 - (c) f(a) = f(b) then $\exists c \in (a, b)$ s.t. f(c) = 0



Q.1. If
$$f(x) = \begin{cases} 1+x & \text{if } x < 0 \\ (1-x)(px+q) & \text{if } x \ge 0 \end{cases}$$
 satisfies the

assumption of Rolle's theorem in the interval [-1, 1] then the order pair (p, q) is IIT JAM 2017

- (a)(2,-1)
- (b) (-2, -1)
- (c)(-2,1)
- (d)(2,1)

Using Rolle's theorem, the equation $a_0x^n + a_1x^{n-1}$ Q.3. $+a_2x^{n-2}+....+a_n=0$ has at least one root between 0 and 1, If

(a)
$$\frac{a_0}{n} + \frac{a_1}{n-1} + \dots \cdot a_{n-1} = 0$$

(b)
$$\frac{a_0}{n-1} + \frac{a_1}{n-2} + \dots \cdot a_{n-2} = 0$$

(c)
$$\frac{a_0}{n+1} + \frac{a_1}{n} + \dots + a_n = 0$$

(d) $a_0 n + a_1 (n-1) + \dots + a_{n-1} = 0$

(d)
$$a_0 n + a_1 (n-1) + \dots a_{n-1} = 0$$


Lagrange's mean value theorem (LMVT):

Let f be a function defined on [a, b] s.t.

- (i) f is continuous on [a, b]
- (ii) f is differentiable on (a, b),

then $\exists c \in (a, b)$

s.t.
$$\frac{f(b)-f(a)}{b-a}=f`(c)$$



Q.3. Let $f : [a, b] \to R$ be a differentiable function. there exist point $c_1, c_2 \in (a, b)$ then which of the following is true

IIT JAM 2005

(a)
$$3f(c_1) f(c_1) = f'(c_2) [f(a) - f(b)]$$

(b)
$$4f(c_1) f(c_1) = f''(c_2) [f(a) + f(b)]$$

(c)
$$5f(c_1) f''(c_1) = f'(c_2) [f(a) + f(b)]$$

(d)
$$2f(c_1) f'(c_1) = f'(c_2) [f(a) + f(b)]$$

- Q.4. For a, b ∈ R with a < b, let f : [a, b] → R be continuous on [a, b] and twice differentiable on (a, b). Further, assume that the graph of f intersects the straight line segment joining the points (a, f(a)) and (b, f(b)) at point (c, f(c)) for a < c < b. Then which of the following is always true IIT JAM 2012</p>
 - (a) There exists a real number $\xi \in (a, b)$ such that $f'(\xi) = 0$
 - (b) For all real number $\xi \in (a, b)$ such that $f'(\xi) \neq 0$
 - (c) we can't say
 - (d) None of these



Cauchy's Mean Value Theorem:

Let f & g be two function defined on [a, b] s.t.

- (i) f and g are continuous in [a, b]
- (ii) f & g are differentiable in (a, b)
- (iii) $g'(x) \neq 0$ for each $x \in (a, b)$ and $g(a) \neq g(b)$.

Then \exists at least one point $c \in (a, b)$ s.t. $\frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)}$

Q.5. The value of ξ in the mean value theorem of $f(b) - f(a) = (b-a)f'(\xi)$ for $f(x) = Ax^2 + Bx + C$ in (a, b) is

$$(a) b + a$$

(b)
$$b - a$$

(c)
$$\frac{(b+a)}{2}$$

(d)
$$\frac{(b-a)}{2}$$



Q.6 A function $f(x) = 1 - x^2 + x^3$ is defined in the closed interval [-1, 1]. The value of x, in the open interval (-1, 1) for which the mean value theorem is satisfied is

(a) - 1/2

(b) -1/3

(c) 1/3

(d) 1/2

COMPLETE COURSE ON MATHEMATICS FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINAER ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months	₹ 908 / mo
Save 67%	Total ₹ 21,780
You get 6 months extra for free	Offer expires 15 Jun 2022

You g	et 6 months extra for free	Offer expires 15 Jun 2022
Sa	ve 54%	Total ₹ 14,974
Ø 12	months	₹ 1,248 / mo

9 months	₹ 1,497 / mo
Save 45%	Total ₹ 13,475

6 months	₹ 2,042 / mo
Save 25%	Total ₹ 12,252

₹ 2,269 / mo
Total ₹ 6,807

1 month	₹2,723 / mo
	Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay



After Using My Referral Code

No cost EMI available on 6 months & above subscription plans

You get 6 months extra for free	Offer expires 15 Jun 2022
Save 67%	₹ 21,700 ₹ 19,602
24 months	₹ 817 / mo

2 12 months	₹ 1,123 / mo
Save 54%	₹ 14,974 ₹ 13,477
You get 6 months extra for free	Offer expires 15 Jun 2022

9 months	₹ 1,348 / mo
Save 45%	₹ 13,475 ₹ 12,128

6 months	₹ 1,838 / mo
Save 25%	₹-12,252 ₹ 11,027

3 months	₹ 2,042 / mo
Save 17%	₹ -6,807 ₹ 6,126



Proceed to pay

FOUNDATION COURSE OF MATHEMATICS FOR CSIR-NET

Taylor's infinite series:

Let $n \in N$, I = [a,b] and $f : I \to R$ be a function f, f', f''....f(n) are continuous on I and that f(n+1) exist on (a, b). then

$$f(x) = f(a) + (x-a)f'(a) + \dots + \frac{(x-a)^{n-1}}{(n-1)!}f^{(n-1)}(a) + \frac{(x-a)^n}{(n)!}f^{(n)}(a) + \dots$$

This is called Taylor's infinite series about x = a.

Maclaurin's infinite series:

In Taylor's series put a = 0.

So,
$$f(x) = f(0) + xf'(0) + \dots + \frac{x^n}{n!} f^{(n)}(0)$$

which is called Maclaurin's infinite series

which is called Maclaurin's infinite series

Tricks: If f(x) is continuous function and it is vanishes at countably infinte numbers then it will

be identically zero

- Let S be the set of all continuous function $f: [-1,1] \rightarrow$ Q.7. R satisfying the following three conditions
 - f is infinitely differentiable on the open (i) interval (-1,1)
 - The Taylor's series (ii)

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots \text{ of } f \text{ at } 0$$

$$\text{converges to } f(x) \text{ for each } x \in (-1,1)$$

$$(iii) \ f\left(\frac{1}{n}\right) = 0 \text{ for all } n \in \mathbb{N}$$

(b)
$$f'\left(\frac{1}{2}\right) = 0$$
 for all $f \in S$

(iii)
$$f\left(\frac{1}{n}\right) = 0$$
 for all $n \in \mathbb{N}$

(c)
$$\exists f \in S$$
 such that $f'\left(\frac{1}{2}\right) \neq 0$

Which of the following is trueIIT JAM 2022

(a)
$$f(0) = 0$$
 for all $f \in S$

(d)
$$\exists f \in S$$
 such that $f(x) \neq 0$ for some $x \in [-1,1]$

2.8. Let α be the real number such that the coefficient of x^{125} in Maclaurin's series of $(x + \alpha^3)e^{x}$ is $\frac{28}{(124)!}$, then

α IIT JAM 2020

- (a) 15
- (c) 25

(b) 20

(d) 39

Q.9. Let $f(x) = \sqrt{x} + 6\alpha$, x > 0 and $g(x) = a_0 + a_1(x - 1) + a_2(x - 1)^2$ be the sum of the first three terms of the taylor series of f(x) around x = 1. If g(3) = 3, then α is? **IIT JAM 2019**

(a) I

(b) 1/2

(c) 1/4

(d) 3/4

Q.10. The coefficient of x^2 in the Maclaurin's series expansion of the function $f(x) = xe^x$.

(a) 0

(b) 1

(c)2

(d) 3



Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

November 3

Enroll Now





Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26 9:00 AM

Gajendra Purohit

Enroll Now

Use code GPSIR for 10% off





Educator Profile





Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



Follow

CSIR-UGC NET



HINDI MATHEMATICAL SCIENCES

Course on Linear Algebra, Partial Diff. Equation & Calculus

Starts on Mar 1, 2021 - 24 lessons

Gajendra Purohit



HINDI MATHEMATICAL SCIENCES

Course on Complex Analysis & Integral Equation

Starts on Jan 14, 2021 • 16 lessons

Gajendra Purohit



HINDI MATHEMATICAL SCIENCES

Foundation Course on Mathematics for CSIR 2021

Starts on Dec 7, 2020 • 20 lessons

Gajendra Purohit

Educator highlights

SEE ALL

Works at Pacific Science College

- Studied at M.Sc., NET,
 PhD(Algebra), MBA(Finance),
 BEd
- PhD, NET | Plus Educator For CSIR NET | Youtuber
 (260K+Subs.) | Director Pacific Science College |
- Lives in Udaipur, Rajasthan,
 India
- Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months	₹ 908 / mo
Save 67%	Total ₹ 21,780
You get 6 months extra for free	Offer expires 15 Jun 2022

You g	et 6 months extra for free	Offer expires 15 Jun 2022
Sa	ve 54%	Total ₹ 14,974
Ø 12	months	₹ 1,248 / mo

9 months	₹ 1,497 / mo
Save 45%	Total ₹ 13,475

6 months	₹ 2,042 / mo
Save 25%	Total ₹ 12,252

₹ 2,269 / mo
Total ₹ 6,807

1 month	₹2,723 / mo
	Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay



After Using My Referral Code

No cost EMI available on 6 months & above subscription plans

You get 6 months extra for free	Offer expires 15 Jun 2022
Save 67%	₹ 21,700 ₹ 19,602
24 months	₹ 817 / mo

2 12 months	₹ 1,123 / mo
Save 54%	₹ 14,974 ₹ 13,477
You get 6 months extra for free	Offer expires 15 Jun 2022

9 months	₹ 1,348 / mo
Save 45%	₹ 13,475 ₹ 12,128

6 months	₹ 1,838 / mo
Save 25%	₹-12,252 ₹ 11,027

3 months	₹ 2,042 / mo
Save 17%	₹ -6,807 ₹ 6,126



Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR