



Gajendra Purohit

Legend in CSIR-UGC NET & IIT-JAM

- Unlock Code : GPSIR ~ PhD, CSIR NET (Maths) | Youtuber(800K+165K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

→ **TOP EDUCATOR ON UNACADEMY
FOR CSIR NET & IIT JAM**

YouTuber with 800K Subscribers

→ **AUTHOR OF BEST SELLER BOOK
FOR CSIR NET & IIT JAM**

**Get
10% Off**

Referral Code : GP SIR





Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

📅 November 3

[Enroll Now](#)



Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26
9:00 AM

Gajendra Purohit

[Enroll Now](#)

Use code GPSIR for 10% off



FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



**After Using
My Referral
Code**



GPSIR

Awesome! You get 10% off

Proceed to pay

Differentiability

Let $f : [a, b] \rightarrow \mathbb{R}$ is a real valued function it is said to be a differentiable at $x = c$.

If $\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$ finitely exist.

Right Hand Derivative :

$$Rf'(c) = \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h}$$

Left Hand Derivative :

$$Lf'(c) = \lim_{h \rightarrow 0} \frac{f(c-h) - f(c)}{-h}$$

Note : If $f(x)$ is differentiable at $x = c$

Iff $Rf'(c) = Lf'(c)$

$$f(x) = \begin{cases} x & \text{for } x \neq 0 \\ 0 & \text{for } x = 0 \end{cases}$$

$$\lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h} = \lim_{h \rightarrow 0} \frac{f(c+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{c+h - 0}{h} = \lim_{h \rightarrow 0} 1 = 1$$

DNE

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{for } x \neq 0 \\ 0 & \text{for } x = 0 \end{cases}$$

cont.
 \Rightarrow

$$f(x) = |x| = \begin{cases} -x & x < 0 \\ x & x \geq 0 \end{cases}$$

$$f'(x) = \begin{cases} -1 & x < 0 \\ 1 & x \geq 0 \end{cases}$$

$$f(x) = |x|^2 = \begin{cases} -x^2 & x < 0 \\ x^2 & x \geq 0 \end{cases}$$

$$f'(x) = \begin{cases} -2x & x < 0 \\ 2x & x \geq 0 \end{cases}$$

$$f(x) = |x|^\alpha$$

$\alpha > 1$

$$f(x) = \begin{cases} x \\ 2x-1 \end{cases}$$

$$x \in \varphi$$

$$x \in \varphi^c$$

$$\gamma^4 = 4\gamma$$

$$\gamma^4 - 4\gamma = 0$$

$$\gamma^2(\gamma^2 - 4) = 0$$

$$\begin{array}{l} n=0, 0 \\ \gamma = 2 \end{array}$$

$$f(x) = \begin{cases} \gamma^4 & n \in \varnothing \\ 4\gamma^2 & n \in \varphi^c \end{cases}$$

$$\frac{\gamma^2 = 2n}{\gamma^2 = 2n - 2}$$

$$\frac{2n}{2n-2} \approx 1$$

$$\gamma^2 = 2n-1$$

$$\gamma^2 = 2n+1 = 0$$

$$(n-1) \approx 0$$

$$\begin{array}{l} n=0 \\ \gamma = 1, 1 \end{array}$$

Q.1. Let $f(x) = \begin{cases} \frac{\sin x}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$, then f is

$\alpha > 0$

- (a) Discontinuous
- (b) Continuous but not differentiable
- (c) Differentiable only once
- (d) Differentiable more than once.

$$x^2 \sin \frac{1}{x}$$


$\alpha > 1$

$$f'(x) = 2x \sin \frac{1}{x}$$



Necessary condition for differentiable :

If a function is differentiable at $x = c$, then it is continuous at $x = c$ but converse may not be true.



Conclusion :

$$f(x) = \begin{cases} x^\alpha \sin \frac{1}{x^\beta}; & x \neq 0 \\ 0; & x = 0 \end{cases}$$

- (i) $f(x)$ is continuous at $x = 0$ for $\alpha > 0$
- (ii) $f(x)$ is differentiable at $x = 0$ for $\alpha > 1$

Result :

(1) $f(x) = \begin{cases} x^\alpha \sin \frac{1}{x^\beta}; & x \neq 0 \\ 0; & x = 0 \end{cases}$

then $f(x)$ is differentiable $\left[\frac{p}{q} \right]$ times and

$$f^{\left[\frac{p}{q} \right]} = \begin{cases} \text{continuous if } p \text{ is odd} \\ \text{discontinuous if } p \text{ is even} \end{cases}$$

$$f'(x)$$

$$\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x} = \lim_{x \rightarrow 0} \frac{x^\alpha \sin \frac{1}{x^\beta}}{x} = \lim_{x \rightarrow 0} x^{\alpha-\beta} \sin \frac{1}{x^\beta}$$

$$f'(0) = \lim_{n \rightarrow \infty} \frac{1}{n} \sin \frac{1}{n^\beta} \neq 0$$

$$\left[\frac{\alpha}{\beta} \right]$$

$$\left[\frac{3}{2} \right] = 1$$

(2) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \begin{cases} \phi_1(x) & x \in Q \\ \phi_2(x) & x \in Q^c \end{cases}$

$f(x)$ is differentiable at the double root of $\phi_1(x) - \phi_2(x) = 0$

- (3) A function is not differentiable at that point at which graph of function is sharp edge.

Q.2. Consider the function $f(x) = |\cos x| + |\sin(2 - x)|$.

At which of the following points is f not differentiable?

- (a) $\left\{(2n+1)\frac{\pi}{2} : n \in \mathbb{Z}\right\}$ (b) $\{n\pi : n \in \mathbb{Z}\}$
- (c) $\{n\pi + 2 : n \in \mathbb{Z}\}$ (d) $\left\{\frac{n\pi}{2} : n \in \mathbb{Z}\right\}$

$$\delta_{2n}(\pi) = 0$$

$$C_{2n} = 0$$

$$C_{2n} = C_{2(n)} \cancel{\pi}$$

$$\cancel{\pi} = (2n+1)\pi$$

$$\pi - L = \pi \cancel{\pi}$$

$$\cancel{\pi} = n\pi + \cancel{\pi}$$

Q.3. The function $f(x) = a_0 + a_1|x| + a_2|x|^2 + a_3|x|^3$ is differentiable at $x = 0$

(a) for no values of a_0, a_1, a_2, a_3

(b) for any value of a_0, a_1, a_2, a_3

(c) only if $a_1 = 0$

(d) only if both $a_1 = 0$ and $a_3 = 0$

$|M|$

COMPLETE COURSE ON

MATHEMATICS

FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINEAR ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



**After Using
My Referral
Code**



GPSIR

Awesome! You get 10% off

Proceed to pay

**FOUNDATION COURSE OF
MATHEMATICS
FOR CSIR-NET**

Q.4. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function with $f(0) = 0$. If for all $x \in \mathbb{R}$, $1 < f'(x) < 2$, then which one of the following statements is true on $(0, \infty)$?

IIT JAM 2015

- (a) f is unbounded
- (b) f is increasing and bounded
- (c) f has at least one zero
- (d) f is periodic

$$f(x) = \frac{3}{2}x$$

$$f'(x) = \frac{3}{2}$$

Q.5. Let the function $f : \mathbb{R} \rightarrow \mathbb{R}$ as

$$f(x) = \begin{cases} x^2 & x \in (0,2) \cap \mathbb{Q}, \\ 2x-1 & x \in (0,2) \cap (\mathbb{R}/\mathbb{Q}) \end{cases}$$

Which one of the following is not true?

- (a) f is continuous at $x = 1$
- (b) f is differentiable at $x = 1$
- (c) f is not differentiable at $x = 1$
- (d) f is differentiable only at $x = 1$

$$\underline{\underline{y^2 = 2m}}$$

$$\underline{\underline{y^2 - 2m + 1 = 0}}$$

$$\underline{\underline{m=1}}$$

Q.6. Let S be the set of all function $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $|f(x) - f(y)|^2 \leq |x - y|^3$ for all x, y in \mathbb{R} . Then which of the following is /are true IIT JAM 2022

(a) Every function in S is differentiable

(b) \exists a function $f \in S$ s.t. f is twice differentiable but f is not thrice differentiable

(c) \exists a function $f \in S$ s.t. f is differentiable but f is not twice differentiable

(d) f is infinitely differentiable

$$f'(y) = 6$$

$$f(y) = C$$

$$\lim_{n \rightarrow y} \left| \frac{f(n) - f(y)}{n - y} \right| \stackrel{\sim}{\leq} \lim_{n \rightarrow y} (n - y)$$

$$|f'(y)| \stackrel{\sim}{\leq} 0$$

$$\left| \frac{f(x) - f(y)}{|x - y|^2} \right| \stackrel{\sim}{\leq} |x - y|$$

Q.7. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be such that f , f' and f'' are all continuous functions with $f > 0$, $f' \geq 0$ and $f''(0) < 0$, then

$$\lim_{x \rightarrow \infty} \frac{f(x) + f'(x)}{2}$$

IIT JAM 2020

- (a) 0
- (b) 1
- (c) 2
- (d) 3

$$\lim_{n \rightarrow \infty} \frac{e^n + e^{-n}}{2}$$

$$\lim_{n \rightarrow -\infty} \frac{e^{-n} + e^n}{2} = \frac{e^{\infty} - 0}{2} = \infty$$

$f = e^{2x}$

$f' = e^{2x}$

$f'' = e^{2x}$

Q.8. Let $f : [-1, 3] \rightarrow \mathbb{R}$ be a continuous function such that

f differentiable on $(-1, 3)$, $|f'(x)| \leq \frac{3}{2}$ for all $x \in (-1, 3)$,

$f(-1) = 1$ and $f(3) = 7$ then $f(1) = ?$ IIT JAM 2020

(a) 2

(b) 3

(c) 4

(d) 5

$$|f'(x)| \leq \frac{3}{2} \quad (x \in (-1, 3))$$

$$\begin{aligned} f(x) &= \frac{3}{2}x + \frac{5}{2} \\ f(1) &= \frac{3}{2} + \frac{5}{2} = \end{aligned}$$

$$\begin{aligned} |f'(x)| &\leq \frac{3}{2} \\ |f'(1)| &\leq \frac{3}{2} \\ |f'(1)| &= \frac{7-1}{3-1} (1+1) \\ |f'(1)| &= \frac{3}{2} (2) \\ |f'(1)| &= 3 \end{aligned}$$

Q.9. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function such that $f'(x) > f(x)$, for all $x \in \mathbb{R}$ and $f(0) = 1$, then $f(1)$ lies in the interval.

- (a) $(0, e^{-1})$
- (b) (\sqrt{e}, e)
- (c) (e^{-1}, \sqrt{e})
- (d) (e, ∞)

Q.10. Let $f : [0, 1] \rightarrow \mathbb{R}$ be a continuous function such that

$$f\left(\frac{1}{2}\right) = -\frac{1}{2} \text{ and } |f(x) - f(y) - (x - y)| \leq \sin(|x - y|^2), \forall$$

$x, y \in [0, 1]$. then $\int_0^1 f(x) dx$ IIT JAM 2020

- (a) $-1/2$
- (b) $-1/4$
- (c) $1/4$
- (d) $1/2$



Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

📅 November 3

[Enroll Now](#)



Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26
9:00 AM

Gajendra Purohit

[Enroll Now](#)

Use code GPSIR for 10% off



Educator Profile



Gajendra Purohit

#5 Educator in CSIR-UGC NET

[Follow](#)

Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



CSIR-UGC NET

[SEE ALL](#)

HINDI MATHEMATICAL SCIENCES
Course on Linear Algebra, Partial Diff. Equation & Calculus
Starts on Mar 1, 2021 • 24 lessons
Gajendra Purohit

HINDI MATHEMATICAL SCIENCES
Course on Complex Analysis & Integral Equation
Starts on Jan 14, 2021 • 16 lessons
Gajendra Purohit

HINDI MATHEMATICAL SCIENCES
Foundation Course on Mathematics for CSIR 2021
Starts on Dec 7, 2020 • 20 lessons
Gajendra Purohit

Educator highlights

- 📍 Works at Pacific Science College
- 📍 Studied at M.Sc., NET, PhD(Algebra), MBA(Finance), BEd
- 📍 PhD, NET | Plus Educator For CSIR NET | Youtuber (260K+Subs.) | Director Pacific Science College |
- 📍 Lives in Udaipur, Rajasthan, India
- 📍 Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



**After Using
My Referral
Code**



GPSIR

Awesome! You get 10% off

Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR