



Ace your preparation with Top Educators

Get **20% off*** on **IIT JAM** subscriptions


Limited period offer



Unlimited access to Live
and Recorded Classes



PYQs, Live Test
and Quizzes



In class doubt solving
by Top Educators



Structured Batches
& courses

Subscribe Now

Use code
GPSIR

For more details, contact: **8585858585**

*T&C apply, as available on the platforms.


20



Ace your preparation with Top Educators

Get **20% off*** on **CSIR-UGC NET** subscriptions

Limited period offer



Unlimited access to Live
and Recorded Classes



Learn from Top
Educators



Comprehensive
Notes and PDFs



Prepare with Unacademy
Lite Subscription

Subscribe Now

Use code

For more details, contact: **8585858585**

*T&C apply, as available on the platforms.

20



Gajendra Purohit

Legend in CSIR-UGC NET & IIT-JAM

- Unlock Code : GPSIR ~ PhD, CSIR NET (Maths) | Youtuber(800K+165K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

→ **TOP EDUCATOR ON UNACADEMY
FOR CSIR NET & IIT JAM**

YouTuber with 800K Subscribers

→ **AUTHOR OF BEST SELLER BOOK
FOR CSIR NET & IIT JAM**

**Get
10% Off**

Referral Code : GP SIR





RANK BOOSTER COURSE UNIT 2 CSIR NET 2022

23rd AUGUST

Gajendra Purohit

[Enroll Now](#)

USE CODE
GPSIR
FOR 10% OFF



DETAILED COURSE 2.0 LINEAR ALGEBRA FOR IIT JAM 2023

8th SEPTEMBER

Gajendra Purohit

[Enroll Now](#)

USE CODE
GPSIR
FOR 10% OFF



FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



**After Using
My Referral
Code**



GPSIR

Awesome! You get 10% off

Proceed to pay



~~Transpose of a matrix :~~

If $A = [a_{ij}]_{m \times n}$ then transpose of A is denoted by A^T is defined as

$$A^T = [b_{ij}]_{n \times m}.$$

Example : Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 1 \end{bmatrix}$ then $A^T = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 3 & 1 \end{bmatrix}.$

$$2A = \begin{bmatrix} 2a & 2b & 2c \\ 2d & 2e & 2f \end{bmatrix}_{2 \times 3}$$

$$2A^T = 2 \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}_{3 \times 2}$$

~~Properties of transpose of matrix :~~

If A & B are two matrices and k is any number then

(1) $(kA)^T = kA^T$

$$2A^T = 2A$$

$$(kA)^T = kA$$

(2) $(A + B)^T = A^T + B^T$

(3) $\underline{(AB)^T = B^T A^T}$

(4) If A is square matrix then $\underline{(A^n)^T = (A^T)^n}$.

$$(A^3)^T = \underline{(A^T)^3}$$

~~Conjugate of a matrix :~~

If $A = [a_{ij}]_{m \times n}$ then conjugate of A denoted by \bar{A} .

$$z = x + iy$$

$$\bar{z} = x - iy$$

Conjugate transpose of a matrix :

If $A = [a_{ij}]_{m \times n}$ then conjugate transpose of A denoted by A^θ as

$$A^\theta = (\bar{A})^T.$$

(2) $(A + B)^\theta = A^\theta + B^\theta$

(3) $(AB)^\theta = B^\theta A^\theta$

$$\underline{(\bar{A})^T = A^\theta}$$

Trace of a square matrix : Sum of all diagonal elements of a square matrix A is known as Trace of A.

$$A = \begin{bmatrix} 1 & 7 & 5 \\ 3 & 2 & 9 \\ 10 & 5 & 6 \end{bmatrix}$$

$$\text{Tr } A = 1 + 2 + 6 = 9$$

Q.1 Let $A = \begin{bmatrix} a & 0 & 1 \\ 0 & 1 & 2 \\ 1 & 2 & 3 \end{bmatrix}$ be a matrix of order 3 whose Trace is 5, then value of a is

- (a) 0
- (b) 1
- (c) 2
- (d) -1

$$\begin{aligned} \text{Tr } A &= 5 \\ a + 1 + 3 &= 5 \\ a + 4 &= 5 \\ a &= 5 - 4 \Rightarrow \end{aligned}$$

Properties of Trace of matrix (A) :

(1) If A be a square matrix of order n and k is any number
then $\text{Tr}(kA) = k\text{Tr}(A)$

(2) If A & B are square matrix of same order n, then
 $\text{Tr}(A + B) = \text{Tr}(A) + \text{Tr}(B)$

(3) $\text{Tr}(AB) = \text{Tr}(BA)$

(4) If $A = [a_{ij}]_{m \times n}$ is any matrix then $\text{Trace}(AA^T) = \text{Sum of squares of every element of } A$.

(5) $\text{Tr}(AA^T) = \text{sum of the squares of modulus of each element of } A$.

$$|1+1|^2$$

$$|1+1|$$

$$|2i| = 2$$

Q2. A real $n \times n$ matrix $A = [a_{ij}]$ is defined as

$$\begin{cases} a_{ij} = i & \forall i = j \\ 0 & \text{Otherwise} \end{cases}$$

then Trace(A) is

(a) $\frac{n(n+1)}{2}$

(b) $\frac{n(n-1)}{2}$

(c) $\frac{n(n+1)(2n+1)}{2}$

(d) n^2

$$ad_1 = 0$$

$$a_{12} = 2$$

$$a_{23} = 3$$

$$a_{11} = 1$$

$$a_{ij} = 1$$

$$1 + 2 + 3 + \dots + n$$

$$\frac{n(n+1)}{2}$$


A, D, C, B, E

~~TV (B < D E A)~~

~~TV (B < A D E)~~


~~TV (D < C E A B)~~

~~TV (D < B A C)~~



Q.3 If A, B & C are square matrix of same order then which of the following is/are equal to $\text{Tr}(ABC)$ is

- (a) $\text{Tr}(ACB)$
- (b) $\text{Tr}(BCA)$
- (c) $\text{Tr}(BAC)$
- (d) $\text{Tr}(CAB)$



a c
b d
c b
d g

Q.4. If $S = \left\{ A = [a_{ij}]_{3 \times 3} \mid \underline{AA^T} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \text{ & } a_{ij} \in R \right\}$. Then

is

- (a) Empty set
- (c) Countably infinite set

- (b) Singleton set
- (d) Uncountable set

$$\text{Tr}(AA^T) = 0$$

$$A = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}$$

$$\text{Tr}(AA^T) = a^2 + b^2 + c^2 + d^2 + e^2 + f^2 + g^2 + h^2 + i^2$$

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$AA^T = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

COMPLETE COURSE ON

MATHEMATICS

FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINEAR ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

**FOUNDATION COURSE OF
MATHEMATICS
FOR CSIR-NET**

Definition of Determinant : Let $A = [a_{ij}]_{n \times n}$ be a matrix, then $|A|$ is called a determinant of order n.

(1) Determinant of order 2 :

Let $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$

$|A| = a_{11} \cdot a_{22} - a_{21} \cdot a_{12}$

$$A = \begin{pmatrix} 5 & 7 \\ 2 & 3 \end{pmatrix}$$

$$|A| = 15 - 14 = 1$$


(2) ~~Determinant of order 3 :~~

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$

Then $|A| = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$


Singular matrix : If $|A| = 0$ then this matrix is called Singular matrix

Non – Singular matrix : If $|A| \neq 0$, then this matrix is called non – singular matrix



$$(6+0+4) - (0+6+3)$$

$$\overline{4-3=1}$$



$$(14 + 18 + 20) - (48 + 105 - 1)$$

$$24 - 48 - 105 + 1$$

$$\underline{-128}$$

Q .5. The number of distinct real values of x for which the

matrix $\begin{pmatrix} 0 & 1 & 1 \\ 1 & x & 1 \\ 1 & 1 & x \end{pmatrix}$ is singular is

- (a) 1
(b) 2
(c) 3
(d) infinite

$$(1 + \gamma^2 + 1) - (n + n + \gamma) = 0$$

$$\gamma^2 + 2 - 3\gamma = 0$$

$$\gamma^2 - 3\gamma + 2 = 0$$

$$1, 1, -1$$

100-2

Property of determinant :

(1) Determinant of diagonal matrix is product of all diagonal element


(2) Determinant of Identity matrix is always one


(3) Determinant of upper or lower triangular matrix is always product of diagonal elements

(4) The value of determinant does not change when rows and columns are interchanged

(5) If any two rows or columns are identically then value of determinant is zero

$$\overline{\det A} = \det A^T$$





(6) $|A^T| = |A|$

(7) $|A+B|$ and $|A| + |B|$ may not be equal

(8) Let $A = [a_{ij}]_{n \times n}$ & then $|kA| = k^n |A|$

(9) Let A & B are two matrix then $|AB| = |A| \cdot |B|$

$\det(A_{4 \times 4}) = 10$

$\det(2A) = \frac{4}{2 \times 10}$

$= 15 \times 10$

$= 150$

Q.6.

Let P be 4×4 matrix whose determinant is 10. The determinant of the matrix $-3P$ is

- (a) -810
- (c) 30

(b) -30

(d) 810

$$\begin{aligned} \text{det } P &= 10 \\ \text{det } (-3P) &= (-3)^4 \cdot 10 \\ &= \underline{\underline{810}} \end{aligned}$$

Q7.

Let $A = \begin{bmatrix} 2 & 3 & 1 \\ 5 & 1 & 2 \\ 1 & 0 & 0 \end{bmatrix}$ then

- (a) A is invertible
- (b) $|A|$ is odd
- (c) $|A|$ is divisible by 2
- (d) $|A|$ is prime number

$$\begin{aligned}(0+0+\cancel{1}) - (0+1+\cancel{0}) \\ \cancel{1}-1 = \cancel{5}\end{aligned}$$

Q.8. It is known that $X = X_0 \in M_2(\mathbb{Z})$ is a solution of $AX - XA = A$

for some $A \in \left\{ \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}, \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \right\}$. Which of the

following values are not possible for the determinant of X_0 ?

CSIR NET FEB 2022

- (a) $\det(X_0) = 0$
- (b) $\det(X_0) = 2$
- (c) $\det(X_0) = 6$
- (d) $\det(X_0) = 10$

Q.9. Let M & N be any two 4×4 matrices with integer entries.

Satisfying $MN = 2$. Then the maximum value of

$$\begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

$\det(M) + \det(N)$ is

- (a) 16
- (b) 17
- (c) 18
- (d) 19

$$\underline{\det(MN)} = 16$$

$$\underline{\det M} \underline{\det N} = \frac{1 \times 16}{2 \times 8}$$

$$\frac{4 \times 4}{4 \times 4}$$

17
10

8

$$A_{n \times n} = (a_{ij}), \quad n \geq 2$$

$$a_{ij} = (b_i^2 - b_j^2)$$

$\det(A)$



(a) $\pi(b_i - b_j)$

(b) $\pi(b_i + b_j)$

(c) $A = \begin{bmatrix} 0 & b_1^2 - b_2^2 & b_1^2 - b_3^2 \\ b_2^2 - b_1^2 & 0 & b_2^2 - b_3^2 \\ b_3^2 - b_1^2 & b_3^2 - b_2^2 & 0 \end{bmatrix}$

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

$|A| = 0$



RANK BOOSTER COURSE UNIT 2 CSIR NET 2022

23rd AUGUST

Gajendra Purohit

Enroll Now

USE CODE
GPSIR
FOR 10% OFF



DETAILED COURSE 2.0 LINEAR ALGEBRA FOR IIT JAM 2023

8th SEPTEMBER

Gajendra Purohit

Enroll Now

USE CODE
GPSIR
FOR 10% OFF



Educator Profile



Gajendra Purohit

#5 Educator in CSIR-UGC NET

[Follow](#)

Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



CSIR-UGC NET

[SEE ALL](#)

HINDI MATHEMATICAL SCIENCES
Course on Linear Algebra, Partial Diff. Equation & Calculus
Starts on Mar 1, 2021 • 24 lessons
Gajendra Purohit

HINDI MATHEMATICAL SCIENCES
Course on Complex Analysis & Integral Equation
Starts on Jan 14, 2021 • 16 lessons
Gajendra Purohit

HINDI MATHEMATICAL SCIENCES
Foundation Course on Mathematics for CSIR 2021
Starts on Dec 7, 2020 • 20 lessons
Gajendra Purohit

Educator highlights

- 📍 Works at Pacific Science College
- 📍 Studied at M.Sc., NET, PhD(Algebra), MBA(Finance), BEd
- 📍 PhD, NET | Plus Educator For CSIR NET | Youtuber (260K+Subs.) | Director Pacific Science College |
- 📍 Lives in Udaipur, Rajasthan, India
- 📍 Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR