

Gajendra Purohit



Unlock Code: GPSIR ~ PhD, CSIR NET (Maths) | Youtuber(800K+165K
 Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

TOP EDUCATOR ON UNACADEMY FOR CSIR NET & IIT JAM

YouTuber with 800K \$ubscribers

AUTHOR OF BEST SELLER BOOK FOR CSIR NET & IIT JAM

Get 10% Off

Referral Code: GP SIR





Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

November 3

Enroll Now





Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26 9:00 AM

Gajendra Purohit

Enroll Now

Use code GPSIR for 10% off







Save up to 20% on your IIT JAM Subscriptions!*

Now available at the lowest prices!*

Plus

Duration	Current Price	Price Drop	What you Pay	What you save
3 Months	₹ 6,807	₹ 1,361	₹ 5,446	20%



Subscribe Now

Use code GPSIR Call 8585858585 for more details

^{*}T&C apply, as available on the platform.

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months	₹ 908 / mo
Save 67%	Total ₹ 21,780
You get 6 months extra for free	Offer expires 15 Jun 2022

You g	et 6 months extra for free	Offer expires 15 Jun 2022
Sa	ve 54%	Total ₹ 14,974
Ø 12	months	₹ 1,248 / mo

9 months	₹ 1,497 / mo
Save 45%	Total ₹ 13,475

6 months	₹ 2,042 / mo
Save 25%	Total ₹ 12,252

₹ 2,269 / mo
Total ₹ 6,807

1 month	₹2,723 / mo
	Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay



After Using My Referral Code

No cost EMI available on 6 months & above subscription plans

You get 6 months extra for free	Offer expires 15 Jun 2022
Save 67%	₹ 21,700 ₹ 19,602
24 months	₹ 817 / mo

2 12 months	₹ 1,123 / mo
Save 54%	₹ 14,974 ₹ 13,477
You get 6 months extra for free	Offer expires 15 Jun 2022

9 months	₹ 1,348 / mo
Save 45%	₹ 13,475 ₹ 12,128

6 months	₹ 1,838 / mo
Save 25%	₹-12,252 ₹ 11,027

3 months	₹ 2,042 / mo
Save 17%	₹ -6,807 ₹ 6,126



Proceed to pay

Function of several variable:

Definition of n-tuples: The Euclidean n-space R^n is the set of all n-tuples $(x_1, x_2, ..., x_n)$ of real numbers on which the two operation of vector addition and scalar multiplication are defined as follow:

(i) Vector addition of two n-tuples:

Let $(x_1, x_2, ..., x_n)$ & $(y_1, y_2,, y_n)$ are two vectors then

$$(x_1, x_2, ..., x_n) + (y_1, y_2, ..., y_n) = (x_1 + y_1, x_2 + y_2, + x_n + y_n)$$

(ii) Scalar multiplication:

Let
$$(x_1, x_2,, x_n) \in R^n \& \alpha \in R$$
, then
$$\alpha(x_1, x_2,, x_n) = (\alpha x_1, \alpha x_2,, \alpha x_n)$$

Real valued function on n-variables:

Let $S \subseteq \mathbb{R}^n$, then a map $f: S \to \mathbb{R}$ is called real valued function on n variables.

Limit of function of two variable:

Let $f: \mathbb{R}^2 \to \mathbb{R}$ be a function & $(a, b) \in \mathbb{R}^2$, $l \in \mathbb{R}$, then we


usually denote limit by $\lim_{x \to \infty} f(x, y) = l$.


Important method:

Path method:


If f(x, y) gives same values 'l' along all paths $y = \phi(x) / \gamma = \sqrt{\gamma^2 + \gamma^2}$

Then
$$\lim_{(x,y)\to(a,b)} f(x,y) = l$$
.






Dr.Gajendra Purohit (PhD,NET) Referral Code [F] 5







Convert function from cartisian form to polar form:

Let f(x, y) is a function of two variable in cartisian form, then put $x = r \cos \theta \& y = r \sin \theta$

Then f(x, y) convert into $f(r, \theta)$ which is called in polar form.

Bounded function:

A function of two variables f(x, y) is said to be bounded iff $\exists M > 0$ s.t. $|f(x, y)| \le M$; for all (x, y)

Results: If a function f(x, y) is bounded then limit of this function need not be exist.

Results: If limit of function exist then function need not be bounded.

Function from Rn to Rm:

Let
$$f: \mathbb{R}^n \to \mathbb{R}^m$$

Let
$$f: \mathbb{R}^n \to \mathbb{R}^m$$

s.t. $f(x_1, x_2,, x_n) = (\phi_1, \phi_2,, \phi_m)$, where $\phi_i: \mathbb{R}^n \to \mathbb{R}$

Limit of a function from $\mathbb{R}^n \to \mathbb{R}^m$:

Let
$$f: \mathbb{R}^n \to \mathbb{R}^m$$

s.t.
$$f(x_1, ..., x_n) = (\phi_1, ..., \phi_m)$$

f has a limit at point (a, b) iff all ϕ_i have limit at (a, b).

Q.1. Let
$$L = \lim_{(x,y)\to(2,-2)} \frac{\sqrt{x-y}-2}{(x-y)-4}$$
, then L is

IIT JAM 2016

(a) 1/2

(b) 1/4

(c) 1/8

(d) 1

COMPLETE COURSE ON MATHEMATICS FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINAER ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months	₹ 908 / mo
Save 67%	Total ₹ 21,780
You get 6 months extra for free	Offer expires 15 Jun 2022

You g	et 6 months extra for free	Offer expires 15 Jun 2022
Sa	ve 54%	Total ₹ 14,974
Ø 12	months	₹ 1,248 / mo

9 months	₹ 1,497 / mo
Save 45%	Total ₹ 13,475

6 months	₹ 2,042 / mo
Save 25%	Total ₹ 12,252

₹ 2,269 / mo
Total ₹ 6,807

1 month	₹2,723 / mo
	Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay



After Using My Referral Code

No cost EMI available on 6 months & above subscription plans

You get 6 months extra for free	Offer expires 15 Jun 2022
Save 67%	₹ 21,700 ₹ 19,602
24 months	₹ 817 / mo

2 12 months	₹ 1,123 / mo
Save 54%	₹ 14,974 ₹ 13,477
You get 6 months extra for free	Offer expires 15 Jun 2022

9 months	₹ 1,348 / mo
Save 45%	₹ 13,475 ₹ 12,128

6 months	₹ 1,838 / mo
Save 25%	₹-12,252 ₹ 11,027

3 months	₹ 2,042 / mo
Save 17%	₹ -6,807 ₹ 6,126



Proceed to pay

FOUNDATION COURSE OF MATHEMATICS FOR CSIR-NET

Q.3. Let
$$I = \lim_{(x,y)\to(0,0)} \frac{x^2y^2}{x^2+y^2}$$
 and Consider the set $A = \{x \in R \mid e^x = l\}$. Then

 $(a) A = \phi$

- (b) A is singleton
- (c) A is countably infinite
- (d) A is uncountable

Q.3. Let
$$L = \lim_{(x,y)\to(0,0)} \frac{x^2y}{x^4 + y^2}$$
, then L is

(a) 0

(b) 1

(c) 1/2

(d) does not exist

For $t \in \mathbb{R}$, let [t] denote the greatest integer less than or equal to t. Define function

$$h \ : \ R^2 \ \to \ R \ and \ g \ : \ R \ \to \ R \ by$$

$$h(x,y) = \begin{cases} \frac{-1}{x^2 - y} & \text{if } x^2 \neq y \\ 0 & \text{if } (x^2 = y) \end{cases}$$

and
$$g(x, y) = \begin{cases} \frac{\sin x}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$
 then which of the

(a)
$$\lim_{(x,y)\to(\sqrt{2},\pi)} \cos\left(\frac{x^2y}{x^2+1}\right) = \frac{-1}{2}$$

(b)
$$\lim_{(x,y)\to(\sqrt{2},2)} e^{h(x,y)} = 0$$

(c)
$$\lim_{(x,y)\to(e,e)} \log(x^{y-[y]}) \neq e-2$$

(d)
$$\lim_{(x,y)\to(0,0)} e^{2y} g(x) = 1$$

Dr.Gajendra Purohit (PhD,NET) Referral Code [FP5]

Q.5. Let S be the set of $(\alpha, \beta) \in \mathbb{R}^2$ s.t. $\frac{x^{\alpha}y^{\beta}}{\sqrt{2}}$


$$(x, y) \rightarrow (0, 0)$$
, then S is


(a)
$$\{(\alpha, \beta); \alpha > 0, \beta > 0\}$$

(b)
$$\{(\alpha, \beta) ; \alpha > 2, \beta > 2\}$$

(c)
$$\{(\alpha, \beta); \alpha + \beta > 1\}$$

(d)
$$\{(\alpha, \beta); \alpha + 4\beta > 1\}$$






Q.6. Statement – 1: $\lim_{(x,y)\to(0,0)} \frac{y^2y^2}{(x,y)\to(0,0)} = \exp(-\frac{y^2y^2}{2} + \frac{y^2}{2})^2$ exists.

Statement – 2:
$$\lim_{(y,y)\to(0,0)} \frac{x^2 y^2}{(x^2+v^2)^{3/2}}$$
 exists.

Then

- (a) Statement 1 is true but statement 2 is not
- (b) Statement 2 is true but statement 1 is not
- (c) Both statements are true
- (d) Both statements are false



- 0). Where $f(x, y) = \frac{x^3}{100}$

(b) for $\alpha = 2$, $\beta = 2$

(d) None of these

73+ m3y (+> 2my 4(n+ my3) 2+m342-+3mx (41-m43)

Dr.Gajendra Purohit (PhD,NET) Referral Code [FPS]



Q.8. Let
$$f(x,y) = \frac{x \cdot y^2}{x^2 + y^4}$$
. Then $\lim_{(x,y)\to(0,0)} f(x,y)$.

- (a) is equal to 1/2 (b) is equal to 2/5
- (d) none of the above

Q.9. Let $R^2 \rightarrow R$ be defined by

$$f(x,y) = \begin{cases} \frac{x^2 - v^2}{x^2 + y^2}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$$
. Then is equal to

(a) 1/3

(b) 2/3

(c) 4/3

(d) None of the above



Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

November 3

Enroll Now





Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26 9:00 AM

Gajendra Purohit

Enroll Now

Use code GPSIR for 10% off





Educator Profile





Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



Follow

CSIR-UGC NET



HINDI MATHEMATICAL SCIENCES

Course on Linear Algebra, Partial Diff. Equation & Calculus

Starts on Mar 1, 2021 • 24 lessons

Gajendra Purohit



HINDI MATHEMATICAL SCIENCES

Course on Complex Analysis & Integral Equation

Starts on Jan 14, 2021 • 16 lessons

Gajendra Purohit



HINDI MATHEMATICAL SCIENCES

Foundation Course on Mathematics for CSIR 2021

Starts on Dec 7, 2020 • 20 lessons

Gajendra Purohit

Educator highlights

SEE ALL

Works at Pacific Science College

- Studied at M.Sc., NET,
 PhD(Algebra), MBA(Finance),
 BEd
- PhD, NET | Plus Educator For CSIR NET | Youtuber
 (260K+Subs.) | Director Pacific Science College |
- Lives in Udaipur, Rajasthan,
 India
- Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

You get 6 months extra for free	Offer expires 15 Jun 2022
Save 67%	Total ₹ 21,780
24 months	₹ 908 / mo

12 months	₹ 1,248 / mo
Save 54%	Total ₹ 14,974
You get 6 months extra for free	Offer expires 15 Jun 2022

9 months	₹ 1,497 / mo
Save 45%	Total ₹ 13,475

6 months	₹ 2,042 / mo
Save 25%	Total ₹ 12,252

3 months	₹ 2,269 / mo
Save 17%	Total ₹ 6,807

1 month	₹ 2,723 / mo
	Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay



After Using My Referral Code

No cost EMI available on 6 months & above subscription plans

You get 6 months extra for free	Offer expires 15 Jun 2022
Save 67%	₹ 21,700 ₹ 19,602
24 months	₹ 817 / mo



9 months	₹ 1,348 / mo
Save 45%	₹ 13,475 ₹ 12,128

6 months	₹ 1,838 / mo
Save 25%	₹-12,252 ₹ 11,027

3 months	₹ 2,042 / mo
Save 17%	₹ -6,807 ₹ 6,126



Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR