



Gajendra Purohit

Legend in CSIR-UGC NET & IIT-JAM

- Unlock Code : GPSIR ~ PhD, CSIR NET (Maths) | Youtuber(800K+165K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

→ **TOP EDUCATOR ON UNACADEMY
FOR CSIR NET & IIT JAM**

YouTuber with 800K Subscribers

→ **AUTHOR OF BEST SELLER BOOK
FOR CSIR NET & IIT JAM**

**Get
10% Off**

Referral Code : GP SIR





Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26
9:00 AM

Gajendra Purohit

[Enroll Now](#)

Use code GPSIR for 10% off



DETAILED COURSE 2.0 GROUP THEORY FOR IIT JAM 2023

6th OCTOBER

Gajendra Purohit

[Enroll Now](#)

USE CODE
GPSIR
FOR 10% OFF



FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

Subsequence : Let $\langle a_n \rangle$ be a given sequence. If $\langle n_k \rangle$ is strictly

increasing sequence of natural numbers, then $\langle a_{n_k} \rangle$ is called a

subsequence of $\langle a_n \rangle$.

2 1 5 3 4

$$a_n = \langle n \rangle = \langle 1, 2, 3, 4, \dots \rangle$$

$$a_{2n} = \langle 2n \rangle = \langle 2, 4, 6, \dots \rangle$$

$$a_{2n+1} = \langle 2n+1 \rangle = \langle 1, 3, 5, \dots \rangle$$

$$a_{n^2} = \langle 1, 4, 9, 16, \dots \rangle$$

$$a_{n^2+1} = \langle 2, 5, 10, 17, \dots \rangle$$

$$a_{2^n} = \langle 2, 4, 8, 16, \dots \rangle$$

$$\langle a_n \rangle = \langle 1^n \rangle = \underbrace{\langle 1, 1, 1, \dots \rangle}_{\text{if } n \in \mathbb{N}}$$

$$\cancel{\langle a_{2n} \rangle} = \cancel{\langle 1^{2^n} \rangle} = \langle 1, 1, 1, \dots \rangle$$


$$\cancel{\langle a_{2n+1} \rangle} = \cancel{\langle 1^{2^n+1} \rangle} = \langle 1, 1, 1, \dots \rangle$$

$$\langle a_n \rangle = \begin{cases} 2 & \text{if } n \in \mathbb{N} \text{ form} \\ n & \text{if } n \notin \mathbb{N} \text{ - } \end{cases}$$

$$= \langle 1, 2, \cancel{2}, 4, 2, 6, \cancel{2}, 8, 9, 10, 2, \dots \rangle$$

$$\cancel{a_n} = \langle 2, 2, 2, \dots \rangle$$

$$a_{n \neq p} = \langle 1, 4, 6, 8, 10, \dots \rangle$$





Complementary Subsequence : If all the terms of two subsequences are distinct then these subsequences are called complementary subsequences.

$$\begin{aligned}\langle a_m \rangle &= \langle 1^n \rangle \\ \cancel{\langle a_n \rangle} &= \langle 1, 1, 1, \dots \rangle \\ \cancel{\langle a_{2n+1} \rangle} &= \langle 1, 1, 1, \dots \rangle\end{aligned}$$


Important result :

- (1) If a sequence $\langle a_n \rangle$ converges to l , then every subsequence of $\langle a_n \rangle$ converges to l . Converse may not be true
- (2) If two complementary subsequences of a sequence $\langle a_n \rangle$ are converges to same limit l , then $\langle a_n \rangle$ converge to l . This is best process for show that sequence is not converge.

Important result :

- (1) If a sequence $\langle a_n \rangle$ converges to l , then every subsequence of $\langle a_n \rangle$ converges to l .

i.e. all subsequences of a convergent sequence converge to same limit but converse need not be true.




(2) If two complementary subsequences of a sequence $\langle a_n \rangle$ are converges to same limit l , then $\langle a_n \rangle$ converges to l .

(3) Every subsequence of divergent sequence is divergent but converse need not true.

(4) Every subsequence of a monotonic sequence are monotonic.

(5) A non-monotonic sequence can have monotonic subsequence.

(6) Every subsequence of bounded sequence is bounded but converse need not be true.



(7) Every sequence have atleast one monotonic subsequence.

(8) A bounded sequence have atleast one convergent subsequence.

(9) If two subsequence are convergent then sequence need not be convergent.

Cauchy Sequence : A sequence is called a cauchy sequence iff it is convergent sequence.



$1 + (-1)^n$

$\langle a_0, a_1, \dots \rangle$

$\langle 3, 2, -1, \dots \rangle$

Q1. Let $\langle a_n \rangle$, $\langle b_n \rangle$ and $\langle c_n \rangle$ be sequences of real numbers such that
 $b_n = a_{2n}$ and $c_n = a_{2n+1}$. Then $\langle a_n \rangle$ is convergent.

- (a) Implies $\langle b_n \rangle$ is convergent but $\langle c_n \rangle$ need not be convergent.
- (b) Implies $\langle c_n \rangle$ is convergent but $\langle b_n \rangle$ need not be convergent.
- (c) Implies both $\langle b_n \rangle$ and $\langle c_n \rangle$ are convergent.
- (d) If both $\langle b_n \rangle$ and $\langle c_n \rangle$ are convergent.


▲ 1 • Asked by Megha

Dekhiye sir



Q2. If $\langle x_n \rangle$ is a convergent sequence in \mathbb{R} and $\langle y_n \rangle$ is a bounded sequence in \mathbb{R} , then we can conclude that

- (a) $\langle x_n + y_n \rangle$ is convergent
- (b) $\langle x_n + y_n \rangle$ is bounded
- (c) $\langle x_n + y_n \rangle$ has no convergent subsequence
- (d) $\langle x_n + y_n \rangle$ has no bounded subsequence.



COMPLETE COURSE ON

MATHEMATICS

FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINEAR ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

FOUNDATION COURSE OF

MATHEMATICS

FOR CSIR-NET


~~Q3.~~ Which of the following is true?

(a) Every sequence that has convergent subsequence is a Cauchy sequence.

(b) Every sequence that has convergent subsequence is bounded sequence

(c) The sequence $\langle \sin n \rangle$ has convergent subsequence.

(d) The sequence $\left\langle n \cos \frac{1}{n} \right\rangle$ has a convergent subsequence.



$a_{n\pi}$

Q.4. Which of the following is false?

- (a) Every bounded sequence has a convergent subsequence.
- (b) Every sequence has a monotonic sub-sequence.
- (c) Every sequence has a limit point.
- (d) Limit points of a sequence is always limit of sequence.



Q.5. Which of the following statements about a sequence of real numbers are true

- (a) Every bounded sequence has a convergent subsequence
- (b) Every sequence has monotonic subsequence
- (c) Every sequence has a limit point
- (d) Every sequence has countable number of terms



Q.6. Let $\langle a_n \rangle$ be a bounded sequence of real numbers. then

- (a) There is a subsequence of $\langle a_n \rangle$ which is convergent
- (b) Every subsequence of $\langle a_n \rangle$ is convergent
- (c) There is exactly one subsequence of $\langle a_n \rangle$ which is convergent
- (d) None of these

$$a_{2n}$$

$$a_{3n+1}$$

$$a_n = \underbrace{\langle 1 + (-1)^n \rangle}_{\text{converges to } 1}$$
$$a_{3n} = \underbrace{\langle 1 + (-1)^{3n} \rangle}_{\text{converges to } 1}$$
$$a_{2n} = \underbrace{\langle 1 + (-1)^{2n} \rangle}_{\text{converges to } 2}$$
$$a_{3n+1} = \underbrace{\langle 1 + (-1)^{3n+1} \rangle}_{\text{converges to } 0}$$

Q.7. If a sequence $\langle a_n \rangle$ is bounded, then the sequence

(E11)

- ~~(a) has no convergent subsequence~~
- ~~(b) has a convergent subsequence~~
- ~~(c) must be converge~~
- ~~(d) is monotonic~~

Q.8. Consider the statement :

S_1 : Every Cauchy sequence of R is convergent.

S_2 : Every convergent sequence is a Cauchy sequence.

S_3 : Every Cauchy sequence is bounded.

Which of the following is true?

- (a) Only S_1
- (b) Only S_1 and S_3
- (c) Only S_2 and S_3
- (d) All the statements



Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26
9:00 AM

Gajendra Purohit

[Enroll Now](#)

Use code GPSIR for 10% off



DETAILED COURSE 2.0 **GROUP THEORY FOR IIT JAM 2023**

6th OCTOBER

Gajendra Purohit

[Enroll Now](#)

**USE CODE
GPSIR
FOR 10% OFF**



Educator Profile



Gajendra Purohit

#5 Educator in CSIR-UGC NET

[Follow](#)

Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



CSIR-UGC NET

[SEE ALL](#)

HINDI MATHEMATICAL SCIENCES

Course on Linear Algebra, Partial Diff. Equation & Calculus

Starts on Mar 1, 2021 • 24 lessons

Gajendra Purohit

HINDI MATHEMATICAL SCIENCES

Course on Complex Analysis & Integral Equation

Starts on Jan 14, 2021 • 16 lessons

Gajendra Purohit

HINDI MATHEMATICAL SCIENCES

Foundation Course on Mathematics for CSIR 2021

Starts on Dec 7, 2020 • 20 lessons

Gajendra Purohit

Educator highlights

- 📍 Works at Pacific Science College
- 📍 Studied at M.Sc., NET, PhD(Algebra), MBA(Finance), BEd
- 📍 PhD, NET | Plus Educator For CSIR NET | Youtuber (260K+Subs.) | Director Pacific Science College |
- 📍 Lives in Udaipur, Rajasthan, India
- 📍 Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR