

Riemann Integration

Detailed Course 2.0 on Function of One and Several Variable - IIT JAM, 23



Gajendra Purohit

Legend in CSIR-UGC NET & IIT-JAM

- Unlock Code : GPSIR ~ PhD, CSIR NET (Maths) | Youtuber(800K+165K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

→ **TOP EDUCATOR ON UNACADEMY
FOR CSIR NET & IIT JAM**

YouTuber with 800K Subscribers

→ **AUTHOR OF BEST SELLER BOOK
FOR CSIR NET & IIT JAM**

**Get
10% Off**

Referral Code : GP SIR





Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

📅 November 3

[Enroll Now](#)



Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26
9:00 AM

Gajendra Purohit

[Enroll Now](#)

Use code GPSIR for 10% off



FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off


Proceed to pay

Differentiability

Darboux Theorem :

If a function is differentiable in a closed interval $[a, b]$ and $f'(a), f'(b)$ are of opposite sign, then there exist at least one point c in the open interval (a, b) s.t. $f(c) = 0$.

$$f(x_1) = \underline{v^2} \quad E_{11}$$



$$f'(c) = 2c = 0$$

$c = 0$

$$f'(x_1) = 2$$

~~$f'(-1) = -2$~~

~~$f'(1) = 2$~~

$$f_{xx} = \delta_{nn}$$

$$n \in [0, \pi]$$

$$f'(n) = \cos n$$

$$f'(0) = 1$$

$$f'(\pi) = -1$$


$$f'(\zeta) = \cos \zeta = 0$$

$$\zeta = \pi/2$$

~~Local maxima and local minima :~~

Let $f : I \rightarrow \mathbb{R}$ and c be an interior point of the interval I , then

- (i) $f(c)$ is said to be a local maximum value of the function f , if there exist some neighbourhood $(c - \delta, c + \delta)$ of c , such that $f(c) > f(x)$, for all $(c - \delta, c + \delta)$ then c is point of local maxima.



$f_{max}, \forall x \in (c - \delta, c + \delta)$

$f_{max} > f(c)$

- (2) $f(c)$ is said to be local minimum value of the function f , if there exist some neighbourhood of $(c - \delta, c + \delta)$ of c s.t. $f(c) < f(x)$, for all $x \in (c - \delta, c + \delta)$, and c is called point of local minima.

(3)

Extreme Value :

$f(c)$ is said to be extreme value of f , if it is either a maximum or minimum value.

i.e. a point $x = c$ is said to be extreme point if $f'(c) = 0$




$$y = \sin x$$
$$f'(x) = \cos x = 0$$

Interior extremum theorem :

Let c be the interior point of the interval I at which $f : I \rightarrow \mathbb{R}$

has an extremum value. If $f(c) = 0$.



Note :

- (1) If $f(x)$ is continuous on $[a, b]$ and $f'(x) > 0$ in (a, b) ,
then $f(x)$ is increasing in $[a, b]$.

$$f(x_1) = n^- \quad [0, 4]$$

- (2) If $f(x)$ is continuous in $[a, b]$ and $f'(x) < 0$ in (a, b)
then $f(x)$ is decreasing in $[a, b]$.

$$f(x_1) = n^- \quad E^{410}$$


$$f'(x_1) = 2n \quad (-f_1)$$

$$f'(x_1) = -2x < 0 \quad \underline{(-f_1)}$$

$$f(x_1) = \text{min}$$

$$f'(x_1) = 0$$


$$f''(x_1) < 0$$



$$x \in (\pi_2, 3\pi_2)$$

$$x \in (\pi_1, \pi)$$

$$(\pi_1, m)$$




First Derivative test for extreme values :


Let a function f be differentiable in a neighbourhood of c ,

where f has an extreme value at c , then $f(c)$ is a maximum


value if the sign of f' change from positive to negative.




$$f(x_1) = f_{\text{min}}$$



$$f(x_1) = f_{\text{min}} \quad [0 \ \theta]$$



And $f(c)$ is a minimum value of the sign of f changes from negative to positive.



Second test for extreme value :

(a) If $f'(c) = 0$ and $f''(c) < 0$ then f has a maximum value at $x = c$.


(b) If $f(c) = 0$ and $f''(c) > 0$ then f has a minimum value at $x = c$.

$$f(n) = -2(2-n)^2$$

$$f'(n) = 4(2-n) = 0$$

$$n=2$$

$$f''(n) = -4 < 0$$



$$y = f(x)$$

$$\frac{dy}{dx} = f'(x) \Rightarrow$$

$$f''(x) < 0$$


$$\underline{\text{min}}$$

$$f(x) = (x - 2)^2$$

$$f'(x) = 2(x - 2) = 0$$

$$f''(x) = 2 > 0$$


$$\text{at } x=2$$



Q.1 Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function such that ϕ' is strictly increasing with $\phi'(1) = 0$. Let α and β denote the minimum and maximum values of $\phi(x)$ on the interval $[2, 3]$, respectively. Then which one of the following is TRUE?

IIT JAM 2017

- (a) $\beta = \phi(3)$
- (b) $\alpha = \phi(2.5)$
- (c) $\beta = \phi(2.5)$
- (d) $\alpha = \phi(3)$



COMPLETE COURSE ON

MATHEMATICS

FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINEAR ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



**After Using
My Referral
Code**



GPSIR

Awesome! You get 10% off

Proceed to pay

FOUNDATION COURSE OF

MATHEMATICS

FOR CSIR-NET

Q.2. Let $f : \left(0, \frac{\pi}{2}\right)$ be given by $f(x) = (\sin x)^\pi - \pi \sin x + \pi$



π. Then which of the following statements is/are

TRUE? IIT JAM 2019

- (a) f is an increasing function
- (b) f is a decreasing function
- (c) $f(x) > 0$ for all $x \in \left(0, \frac{\pi}{2}\right)$
- (d) $f(x) < 0$ for some $x \in \left(0, \frac{\pi}{2}\right)$

$$f'(x) = \pi(\sin x)^{\pi-1} - \pi \cos x$$

$$= \pi \sin x \cdot [(\sin x)^{\pi-1} - 1]$$

$$f'\left(\frac{\pi}{4}\right) = \pi \frac{1}{\sqrt{2}} \left[\left(\frac{1}{\sqrt{2}}\right)^{\pi-1} - 1\right] < 0$$

$$\therefore (\sin x)^\pi < 1$$

Q.3. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function with $f(0) = 0$. If for all $x \in \mathbb{R}$, $1 < f'(x) < 2$, then which one of the following statements is true on $(0, \infty)$ IIT

JAM 2015

~~(a) f is unbounded~~

~~(b) f is increasing and bounded~~

~~(c) f has at least one zero~~

~~(d) f is periodic~~

$$f(x) = \frac{3}{2}x$$

Q.4. For $x \in \mathbb{R}$, let $f(x) = \begin{cases} x^3 \sin\left(\frac{1}{x}\right) & x \neq 0 \\ 0 & x = 0 \end{cases}$. Then which

one of the following is FALSE?

IIT JAM 2017

(a) $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 0$

(b) $\lim_{x \rightarrow 0} \frac{f(x)}{x^2} = 0$

(c) $\frac{f(x)}{x^2}$ has infinitely many maxima and minima on the interval $(0, 1)$

(d) $\frac{f(x)}{x^4}$ is continuous at $x = 0$ but not differentiable at $x = 0$.





Q.5. Let $f(x) = \begin{cases} x + x^2 \cos\left(\frac{\pi}{x}\right) & x \neq 0 \\ 0 & x = 0 \end{cases}$ & consider the

following statements :

- (i) $f'(0)$ exists & is equal to 1
- (ii) f is not increasing in any neighbourhood of 0
- (iii) $f'(0)$ does not exist
- (iv) f is increasing on \mathbb{R}

How many of the above statement is/are true?

- (a) 0
- (b) 1
- (c) 2
- (d) 3

$$\begin{aligned}
 f'(0) &= \lim_{n \rightarrow 0} \frac{f(n) - f(0)}{n - 0} \\
 &= \lim_{n \rightarrow 0} \frac{n + n^2 \cos\left(\frac{\pi}{n}\right) - 0}{n} \\
 &= \lim_{n \rightarrow 0} (1 + n \cos\left(\frac{\pi}{n}\right)) \\
 &= 1
 \end{aligned}$$


Q.6. Let $f(x) = \begin{cases} x+1 & x < 0 \\ (x-1)^2 & x \geq 0 \end{cases}$

$$y = (x-1)^2$$

Which one of the following is TRUE?

- (a) f is differentiable on \mathbb{R}
- (b) f has neither a local maximum nor a local minimum in \mathbb{R}
- (c) f is bounded on \mathbb{R}
- (d) f is not differentiable at $x = 0$ and $f(x)$ has local maximum at $x = 0$

$$y = (x-1)^2$$



Q.7. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function such that

~~$f(x+y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$, then~~

(a) f is increasing if $f(1) \geq 0$ and decreasing if $f(1) \leq 0$

$f'(x) = k$

(b) f is increasing if $f(1) \leq 0$ and decreasing if $f(1) \geq 0$

$f(x) = kx$

(c) f is not an increasing function

$f(1) = k \geq 0$

(d) f is neither an increasing nor a decreasing function



Q.8. Let f be a twice differentiable function on \mathbb{R} . Given that $f''(x) > 0$ for all $x \in \mathbb{R}$, then

- (a) $f(x) = 0$ has exactly two solution on \mathbb{R} .
- (b) $f(x) = 0$ has a positive solution if $f(0) = 0$ & $f'(0) = 0$
- (c) $f(x) = 0$ has no positive solution if $f(0) = 0$ and $f'(0) > 0$
- (d) $f(x) = 0$ has no positive solution if $f(0) = 0$ and $f'(0) < 0$



Detailed Course on Group Theory For CSIR NET 2023

Gajendra Purohit

📅 November 3

[Enroll Now](#)



Detailed Course 2.0 on Sequence and Series For IIT JAM' 23

October 26
9:00 AM

Gajendra Purohit

[Enroll Now](#)

Use code GPSIR for 10% off



Educator Profile



Gajendra Purohit

#5 Educator in CSIR-UGC NET

[Follow](#)

Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



CSIR-UGC NET

[SEE ALL](#)

HINDI MATHEMATICAL SCIENCES
Course on Linear Algebra, Partial Diff. Equation & Calculus
Starts on Mar 1, 2021 • 24 lessons
Gajendra Purohit

HINDI MATHEMATICAL SCIENCES
Course on Complex Analysis & Integral Equation
Starts on Jan 14, 2021 • 16 lessons
Gajendra Purohit

HINDI MATHEMATICAL SCIENCES
Foundation Course on Mathematics for CSIR 2021
Starts on Dec 7, 2020 • 20 lessons
Gajendra Purohit

Educator highlights

- 📍 Works at Pacific Science College
- 📍 Studied at M.Sc., NET, PhD(Algebra), MBA(Finance), BEd
- 📍 PhD, NET | Plus Educator For CSIR NET | Youtuber (260K+Subs.) | Director Pacific Science College |
- 📍 Lives in Udaipur, Rajasthan, India
- 📍 Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR