

Doubt Clearing Session

Detailed Course 2.0 on Linear Algebra For IIT JAM' 23


Gajendra Purohit • Lesson 6 • Sept 16, 2022



Ace your preparation with Top Educators

Get **20% off*** on **IIT JAM** subscriptions


Limited period offer



Unlimited access to Live
and Recorded Classes



PYQs, Live Test
and Quizzes



In class doubt solving
by Top Educators



Structured Batches
& courses

Subscribe Now

Use code
GPSIR

For more details, contact: **8585858585**

*T&C apply, as available on the platform.

20




Ace your preparation with Top Educators

Get **20% off*** on **CSIR-UGC NET** subscriptions

Limited period offer



Unlimited access to Live
and Recorded Classes



Learn from Top
Educators



Comprehensive
Notes and PDFs



Prepare with Unacademy
Lite Subscription

Subscribe Now

Use code

For more details, contact: **8585858585**

*T&C apply, as available on the platform.

20



Gajendra Purohit

Legend in CSIR-UGC NET & IIT-JAM

- Unlock Code : GPSIR ~ PhD, CSIR NET (Maths) | Youtuber(800K+165K Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author

50M Watch mins

3M Watch mins (last 30 days)

44K Followers

2K Dedications

→ **TOP EDUCATOR ON UNACADEMY
FOR CSIR NET & IIT JAM**

YouTuber with 800K Subscribers

→ **AUTHOR OF BEST SELLER BOOK
FOR CSIR NET & IIT JAM**

**Get
10% Off**

Referral Code : GP SIR





RANK BOOSTER COURSE UNIT 2 CSIR NET 2022

23rd AUGUST

Gajendra Purohit

[Enroll Now](#)

USE CODE
GPSIR
FOR 10% OFF



DETAILED COURSE 2.0 LINEAR ALGEBRA FOR IIT JAM 2023

8th SEPTEMBER

Gajendra Purohit

[Enroll Now](#)

USE CODE
GPSIR
FOR 10% OFF



FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay


$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 1 & 4 & 3 & 5 \end{pmatrix}$$

$$= \frac{(12)(34)}{2}$$

$\text{Tr}(A) = 1$

$$|A| = (-1)^{\frac{n(n-1)}{2}} \text{Transposition}$$

$$= (-1)^{\frac{2}{2}} = 1$$



$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 2 & 3 \\ 5 & 4 & 3 \end{pmatrix}}_{\sim} \begin{pmatrix} 4 & 5 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 15 & 24 \\ 15 & 14 \end{pmatrix}$$

$\text{Tr}(A) = 1$

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix} = ((2)(4)^3, (4)(5))$$

$$= \underline{(12453)}$$

$$\underline{(13)} \underline{(15)} \underline{(14)} \underline{(12)}$$

$|A| = (-1)^4 = 1$

~~A~~

$\text{Tr}(A) = 0$

$$A = \begin{bmatrix} & \\ & \end{bmatrix}$$

.

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} = (132)$$
$$= \underline{(12)(13)}$$

$$\|A\| = \sqrt{1^2} = 1$$

$$\text{Tr}(A) = 0$$

Matrix and their properties

Block Matrix : Let $A = [a_{ij}]_{m \times n}$, $B = [b_{ij}]_{m \times p}$, $C = [c_{ij}]_{x \times n}$

& $D = [d_{ij}]_{x \times p}$ are matrix then a matrix

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}_{(m+x) \times (n+p)}$$

is called block matrix.

$$M =$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$



Block Diagonal Matrix : Let A & B are two matrix then a

matrix of type $M = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$ is called block diagonal matrix.

$$\det M = \det(A) \det(B)$$

Block Upper Triangular matrix : If all blocks are above the diagonal then this matrix is called block upper triangular matrix.

$$\begin{bmatrix} A & B \\ 0 & C \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 2 & 5 & 5 \\ 6 & -1 & 2 & 0 \\ 0 & 5 & 7 & 8 \end{bmatrix}$$

$$\det M = (4)(-6)(2)(-3)$$

$$= -24 - 28$$

$$= -52$$

$$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$$

$$= \underline{(12)} \underline{(3)}$$

$$= \underline{(x^2-1)(x-1)}$$

$$\begin{bmatrix} 0-\lambda & 1 & 0 \\ 1 & -\lambda & 0 \\ 0 & 0 & 1-\lambda \end{bmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$$

$$= \underline{(1)} \underline{(2)} \underline{(3)}$$

$$(x-1) \ (m) \ (m)$$

$$(m)^2$$

$$\begin{bmatrix}
 0 & 0 & 0 & 1 & 0 \\
 1 & 0 & 0 & 0 & 0 \\
 0 & 1 & 0 & 0 & 0 \\
 0 & 0 & 1 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0
 \end{bmatrix} = \underbrace{\left(\begin{smallmatrix} 1 & 2 & 3 \\ 4 & 1 & 3 \end{smallmatrix} \right) \left(\begin{smallmatrix} 4 & 5 \\ 5 & 2 \end{smallmatrix} \right)}_{= (1452)(3)} = \frac{(1452)(3)}{(14)(m-1)} = 0$$

$$(q^2-1)(q^2+1)(x_1)=0$$

~~$n \geq k+1$~~ , $n = t\lambda^6, 1$

$$\text{oval containing } \langle 1, -1, 1, -1 \rangle$$

$$\left(\begin{array}{cccccc} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{array} \right)$$

which of following is

correct

(a) t^1, t^1, t^1

$$\text{der}(A) = \frac{t^1 t^3}{t^1}$$

(c) $\text{der}(A) = 1$

$$(1 \ 2 \ 3 \ 4 \ 5 \ 6) \\ (4 \ 5 \ 6 \ 1 \ 2)$$

~~$(14)(25)(36)$~~

t^1, t^1, t^1

$$(n^2 - 1) (n^2 - 1) (n^2 - 1) = 0$$

$$n = t^1 \\ n = t^1 \\ n = t^1$$

- (a) a, c
- (b) b, c
- (c) b, d
- (d) a, d

$$\begin{bmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \end{bmatrix} \quad 4 \times 4$$

$$\gamma^4 - \gamma^3 - (-3)\gamma^2 - 2\gamma - (-1)$$

(a)

$$\gamma^4 + 3\gamma^2 + 2\gamma + 1$$

(b)

$$\gamma^4 - 3\gamma^2 - 2\gamma - 1$$

(c)

$$\gamma^4 - 3\gamma^2 + 2\gamma + 1$$

(d)

$$\gamma^4 + 3\gamma^2 - 2\gamma + 1$$

$$\gamma^4 + 3\gamma^2 - 2\gamma + 1$$

$$\begin{bmatrix} 0 & 0 & 0 & 3 \\ 1 & 0 & 0 & 7 \\ 0 & 1 & 0 & -5 \\ 0 & 0 & 1 & -2 \end{bmatrix}$$

$$y^4 + 2y^3 + 5y^2 - 7y - 3 = 0$$

~~Property :~~

- (1) Trace of block matrix is sum of trace of all diagonal block of matrix.
- (2) Determinant of block diagonal matrix is product of determinant of all diagonal block of matrix.

Companion matrix :

Let $P(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1} + a_nx^n$ be a polynomial then

a matrix A corresponding to $P(x)$ is called companion matrix if

$$A = \begin{bmatrix} 0 & 0 & \cdots & -a_0 \\ 1 & 0 & \cdots & -a_1 \\ 0 & 1 & \cdots & -a_2 \\ \vdots & & & \\ 0 & 0 & \cdots & 1-a_{n-1} \end{bmatrix}.$$

$$\det(A) = (-1)^n a_0$$

Property :

- (1) Characteristic polynomial and minimal polynomial are same which equal to given polynomial.
- (2) Suppose $p(x) = \underline{a_0 + a_1x + \dots + a_{n-1}x^{n-1}} + \underline{\cancel{a_n}x^n}$, then
 $\text{Trace}(A) = -(a_{n-1})$ and $|A| = (-1)^m \underline{a_0}$.

Similar Matrix : A & B are said to be similar matrix if \exists a non-singular matrix P s.t. $A = P^{-1}BP$

Congruent Matrix : A & B are said to be congruent matrix if \exists a matrix O s.t. $A = P^TBP$

Adjoint of matrix : Adjoint of a matrix A is the transpose of the cofactor matrix of A denoted as $\text{adj}(A)$.

Property :

$$(1) \quad A(\text{adj } A) = |A| I_n$$

$$(2) \quad |A(\text{adj } A)| = | |A| I_n | = |A|n$$

$$\Rightarrow |A| |\text{adj } A| = |A|n$$

$$\Rightarrow |\text{adj } A| = |A|^{n-1}$$

In general $|\text{adj } \text{adj} \dots \text{adj } A| = |A|^{(n-1)^k}$.
(k-times)

$$(3) \quad (\text{adj } A)^T = \text{adj } (A^T)$$

$$(4) \quad \text{adj}(AB) = \text{adj}(B) \cdot \text{adj}(A)$$

$$(5) \quad |\text{adj } (kA)| = |kA|^{n-1} = k^{n(n-1)} |A|^{n-1}$$

- Q.2.** Suppose A be a 3×3 matrix of determinant 6, then
determinant of $(\text{adj } A)$
- (a) 36
 - (b) 9
 - (c) 25
 - (d) None of these

Q.3. Let A be a 5×5 skew-symmetric matrix with entries in R and B be the 5×5 symmetric matrix whose $(i, j)^{\text{th}}$ entry is

the binomial coefficient $\binom{i}{j}$ for $1 \leq i \leq j \leq 5$. Consider

the 10×10 matrix, given in block form by

$$C = \begin{pmatrix} A & A+B \\ 0 & B \end{pmatrix}. \text{ Then}$$

- (a) $\det C = 1 \text{ or } -1$
(c) trace of C is 0

- (b) $\det C = 0$
(d) trace of C is 5

$\det(C) = \frac{\det(A)}{\det(B)}$
 $= 0 \times \det B = 0$

~~$\begin{pmatrix} 0 & 0 & 0 & \dots & 0 \\ 0 & 2 & 0 & \dots & 0 \\ 0 & 0 & 2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 2 \end{pmatrix}$~~

$$A = \begin{bmatrix} 0 & 3 & 7 \\ 2 & 1 & 3 \\ 0 & 2 & 4 \end{bmatrix}$$

$$\det(A) = 108$$

$$\text{Tr}(A) = 14$$

$$|a-b|$$

(@) 2

(c) 4 (d) 5

$$\det(M) = (a)(2b) = 108$$

$$10 \times 5 = 50$$

$$50 = 10$$

$$a + 5 + 2 + b = 14$$

$$\cancel{a+b=7}$$

$$\begin{cases} a=5 \\ b=2 \end{cases}$$

COMPLETE COURSE ON

MATHEMATICS

FOR IIT-JAM 2022

TOPICS TO BE COVERED

- REAL ANALYSIS
- FUNCTION OF ONE & TWO VARIABLE
- LINEAR ALGEBRA
- MODERN ALGEBRA

TOPICS TO BE COVERED

- SEQUENCE & SERIES
- INTEGRAL CALCULUS
- VECTOR CALCULUS
- DIFFERENTIAL EQUATION

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



**After Using
My Referral
Code**



GPSIR

Awesome! You get 10% off

Proceed to pay

FOUNDATION COURSE OF
MATHEMATICS
FOR CSIR-NET

Q.4. Let $A = \begin{bmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$, then which of the following is true?

- (a) $|A| = 0$
(b) $|A| = 1$
(c) $|A| = 2$
(d) $|A| = -1$

$\left(\text{HIT.}\right)$

$\begin{bmatrix} 0 & 0 & 0 & 0 & -5 \\ 1 & 0 & 0 & 0 & 7 \\ 0 & 1 & 0 & 0 & 3 \\ 0 & 0 & 1 & 0 & 5 \\ 0 & 0 & 0 & 1 & 5 \end{bmatrix}$

$\xrightarrow{\left(\begin{array}{cc} 1 & 5 \\ -1 & 5 \end{array} \right)}$

Q5. Let $A = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$ then characteristic polynomial.

(a) $x^3 - 2x^2 + 1$
~~(c) $x^3 - 2x^2 - 1$~~

(b) $x^2 - 2x + 1$
~~(d) None of these~~

$\begin{array}{r} x^3 \\ -11 \quad -1 \\ -1 \quad \rightarrow = 1 \end{array}$

$$\lambda^3 + 2\lambda^2 - 5\lambda - 1$$

$$\lambda^3 + 2\lambda^2 - 1 = 0$$

Q6. Let $\alpha, \beta, \gamma, \delta$ be the eigenvalues of the matrix

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix}. \text{ Then } \alpha^2 + \beta^2 + \gamma^2 + \delta^2 = \underline{\hspace{2cm}}.$$

(a) 4

(b) 9

(c) 1 + 1 + 1 + 4

$$\lambda^4 - 2\lambda^3 - \lambda^2 + 2\lambda = 0$$

$$(\lambda - 1)^3(\lambda + 2)$$

$$(\lambda^2 - 1)(\lambda + 2)$$

$$\lambda(\lambda^2 - 1)(\lambda + 2)$$

$$\lambda(\lambda - 1)(\lambda + 1)(\lambda + 2)$$

$$\lambda = -2, 1, -1, 2$$

Rank of matrix : If A is matrix then a non-negative integer is said to be rank of A. if \exists a non-singular submatrix of order r of A and all matrix of order greater than r are singular.

Note :

- (1) A matrix $A_{n \times n}$ is a non-singular iff $\text{rank}(A) = n$
- (2) Rank of matrix is denoted by $\rho(A)$
- (3) If $|A| \neq 0$ then $\rho(A) = n$ where n is order of A

Elementary Transformation : Those transformation that does not effect on rank of matrix are called elementary transformation.

Elementary matrix : A matrix obtained by a single elementary operation over identity matrix is known as elementary matrix.

Q.1. Find elementary matrix

(a) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$

(d) $\begin{bmatrix} 3 & 2 & 0 \\ 1 & 2 & 0 \\ 1 & 2 & 2 \end{bmatrix}$

Note :

- (1) Every elementary matrix is non-singular.
- (2) Every permutation matrix is elementary matrix.

Q.2. Let $A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 151 & 262 & 373 & 484 \end{bmatrix}$.

Then $\text{Rank}(A)$ is

- (a) 1
- (b) 2
- (c) 3
- (d) 4



RANK BOOSTER COURSE UNIT 2 CSIR NET 2022

23rd AUGUST

Gajendra Purohit

Enroll Now

USE CODE
GPSIR
FOR 10% OFF



DETAILED COURSE 2.0 LINEAR ALGEBRA FOR IIT JAM 2023

8th SEPTEMBER

Gajendra Purohit

Enroll Now

USE CODE
GPSIR
FOR 10% OFF



Educator Profile



Gajendra Purohit

#5 Educator in CSIR-UGC NET

[Follow](#)

Dr.Gajendra Purohit PhD, CSIR NET (Maths) | Youtuber(330K+30k Sub.)/Dr.Gajendra Purohit (Maths), 17+ Yr. Experience, Author of Bestseller

11M Watch mins

1M Watch mins (last 30 days)

22k Followers

1k Dedications



CSIR-UGC NET

[SEE ALL](#)

HINDI MATHEMATICAL SCIENCES

Course on Linear Algebra, Partial Diff. Equation & Calculus

Starts on Mar 1, 2021 • 24 lessons

Gajendra Purohit

HINDI MATHEMATICAL SCIENCES

Course on Complex Analysis & Integral Equation

Starts on Jan 14, 2021 • 16 lessons

Gajendra Purohit

HINDI MATHEMATICAL SCIENCES

Foundation Course on Mathematics for CSIR 2021

Starts on Dec 7, 2020 • 20 lessons

Gajendra Purohit

Educator highlights

- 📍 Works at Pacific Science College
- 📍 Studied at M.Sc., NET, PhD(Algebra), MBA(Finance), BEd
- 📍 PhD, NET | Plus Educator For CSIR NET | Youtuber (260K+Subs.) | Director Pacific Science College |
- 📍 Lives in Udaipur, Rajasthan, India
- 📍 Unacademy Educator since

FEE DETAILS FOR IIT JAM SUBSCRIPTION

No cost EMI available on 6 months & above subscription plans

24 months ₹ 908 / mo
Save 67%
Total ₹ 21,780

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,248 / mo
Save 54%
Total ₹ 14,974

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,497 / mo
Save 45%
Total ₹ 13,475

6 months ₹ 2,042 / mo
Save 25%
Total ₹ 12,252

3 months ₹ 2,269 / mo
Save 17%
Total ₹ 6,807

1 month ₹ 2,723 / mo
Save 0%
Total ₹ 2,723

To be paid as a one-time payment

Have a referral code?

Proceed to pay

No cost EMI available on 6 months & above subscription plans

24 months ₹ 817 / mo
Save 67%
Total ₹ 21,700 ₹ 19,602

You get 6 months extra for free Offer expires 15 Jun 2022

✓ 12 months ₹ 1,123 / mo
Save 54%
Total ₹ 13,477

You get 6 months extra for free Offer expires 15 Jun 2022

9 months ₹ 1,348 / mo
Save 45%
Total ₹ 12,128

6 months ₹ 1,838 / mo
Save 25%
Total ₹ 11,027

3 months ₹ 2,042 / mo
Save 17%
Total ₹ 6,126



After Using
My Referral
Code



GPSIR

Awesome! You get 10% off

Proceed to pay

THANK YOU VERY MUCH EVERYONE

GET THE UNACADEMY PLUS SUBSCRIPTION SOON.

TO GET 10% DISCOUNT IN TOTAL SUBSCRIPTION AMOUNT

USE REFERRAL CODE: GPSIR