

# Urban Indian Environment in the Context of a Pandemic

Abhijith. S.\*, Akshara S. N.\* and P. P. Nikhil Raj\*\*†

\*Department of Computer Science and Engineering, Amrita School of Computing, Coimbatore, Amrita Vishwa Vidyapeetham, India

\*\*Department of Chemical Engineering and Materials Science, Amrita School of Engineering, Coimbatore, Amrita Vishwa Vidyapeetham, India

ORCID: 0000-0002-5104-4607

†Corresponding author: P. P. Nikhil Raj; cb.en.u4cse19102@cb.students.amrita.edu; pp\_nikhilraj@cb.amrita.edu

**Nat. Env. & Poll. Tech.**  
 Website: [www.neptjournal.com](http://www.neptjournal.com)

Received: 09-06-2023

Revised: 19-07-2023

Accepted: 27-07-2023

## Key Words:

COVID-19, Air Quality Index, Population, Deaths

## ABSTRACT

The spread of the Coronavirus disease 2019 (COVID-19) has impacted human life severely since November 2019. The urban centers in the world, especially, were highly affected by the diseases. Several socioeconomic and environmental factors probably enhanced the spread of the pandemic and consequent mortality. Many studies examining environmental factors, such as air quality, in urban centers indicate the roles of those factors in the spread of diseases and consequent mortality. However, other socioeconomic factors that directly or indirectly elevate the mass death of people are seldom studied. The present study explores the socioeconomic factors and air quality influencing COVID-19 deaths in urban India. We randomly selected 19 Indian cities and collected each city's socioeconomic and air quality data from reliable and open sources. The data were analyzed using multivariate data analysis techniques using R statistics. The results showed significant positive relationships, population, and total area of the urban centers.

## INTRODUCTION

The world has witnessed the intensity of the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-COV-2), also known as COVID-19, since November 2019. So far, the pandemic (declared by the World Health Organization on March 11, 2020) has led to the deaths of more than six million people around the globe (covid19.who.int). The virus is highly versatile and mutating (Balachandran et al. 2022). Its known transmission modes include contact, droplet, airborne, fomite, fecal-oral, bloodborne, mother-to-child, and animal-to-human transmission (www.who.int). The environmental setup is also a crucial factor in determining the spread. Since the Coronavirus attacks the respiratory tract, people exposed to highly polluted air and suffering from respiratory ailments (Subbarao et al. 2020) and people with other comorbidities, heart diseases, lung cancer, etc., are highly vulnerable to COVID-19 infection. People living in urban centers with considerably low air quality are also of prime concern (Conticini et al. 2020, Cao et al. 2020). For example, the highest cases of COVID-19 deaths reported in urban areas in China, South Korea, Iran, and northern Italy were exacerbated by poor air quality (Jiang

et al. 2020, Gupta et al. 2021). The spread of the disease in a region is also linked to its geographical location, economy and public awareness. It is also associated with the region's health systems' capacity, readiness and ability to curtail the spread (Kejela 2020). At an individual level, community consciousness, age, and social (in fact, physical distancing) distancing are other significant factors influencing the spread of the disease (Raj & Azeez 2020, Lakshmi & Suresh 2020, Priya et al. 2021).

Regarding the Air Quality Index in 2020, the air quality in world cities, in general, is almost five times above the WHO exposure recommendations (WHO 2021). Six major pollutants, Carbon Monoxide, Lead, Nitrogen Oxide, Ozone, Particulate matter (PM), and Sulphur dioxide, are highly likely to affect a population's short and long-term health. Of the PMs (PM<sub>10</sub> and PM<sub>2.5</sub>), PM<sub>2.5</sub> is considered the most lethal; chronic exposure to PM<sub>2.5</sub> increases the risk of acute respiratory diseases among the older population (Benmarhnia 2020) since that can penetrate the bronchi deeper (www.who.int). As per a recent report by the European Environment Agency, around 0.30 million premature deaths in urban areas of 27 European countries were due to high PM<sub>2.5</sub>. Vehicular

emission (25%) followed by agriculture (22%), domestic fuel burning (20%), natural dust and salt (18%), and industrial activities (15%) significantly contribute to PMs (PM<sub>10</sub>, PM<sub>2.5</sub>, etc.) in cities (Karagulian et al. 2015).

Due to the poor air quality, approximately 1.24 million people lose their lives in India annually (Balakrishnan et al. 2019). It is reported that high pollution levels in the ambient air contribute to respiratory dysfunctions among urban dwellers (Balakrishnan et al. 2014, Venkataraman et al. 2018, Khilnani & Tiwari 2018). In 2019, air pollution accounted for 17.8 % of total mortality in India, of which 5% of deaths were attributed to PM pollution. However, the deaths due to household air pollution decreased from 74.2 million in 1990 to 0.61 million in 2019, around 64.2% of the total death cases (Pandey et al. 2021). In India, it is reported that the primary sector contributing significantly to ambient air quality is the transport sector, which substantially contributes (7-43%) to ambient air PM (Gurjar et al. 2016). Khilnani & Tiwari (2018) reported that the ambient air pollution in the major city centers of the country is showing an alarmingly rising trend.

From different studies carried out by the Central Pollution Control Board (CPCB), it has been found that 46 Indian cities are severely polluted (Nasir et al. 2016). It is widely known that most Indian metro cities are highly polluted, with 14 out of 20 Indian cities examined being shortlisted as among the world's most air-polluted cities (Sikarwar & Rani 2020). As a fast-developing country (fifth-largest economy in terms of nominal GDP), India has an active industrial sector and fossil fuel-based power plants. From 2010 to 2015, it was observed that the emission of CO<sub>2</sub> gas significantly increased in most parts of the country (Girdhar et al. 2021).

Air pollution is one of the significant outcomes of the current economic development model (Aditya et al. 2022). While the developed nations were historically massive emitters, the rapidly growing economies are increasingly contributing to air pollution since high emission level of air-polluting gasses like CO<sub>2</sub> is usual evils in the current economic development model. WHO and OECD reported that Europe's economic cost of premature death and disability from air pollution in 2015 was nearly USD 1.6 trillion (Manjula 2018). The adverse effects of air pollution are diverse, including social and psychological (Chen & Jin 2019, Heissel et al. 2022). Studies have also shown that a 10% rise in PM<sub>2.5</sub> caused a 2.4% fall in the price of houses in China during 2005-2013 (Sivarethinamohan et al. 2021). Present-day economic developmental models are committed to the financial growth calculated in terms of the Gross Domestic Productivity of a region and that, most times, fail to ensure the socioeconomic welfare of the people, except perhaps by the trickle-down effect. The major

cities in developing countries like India are deprived of scientific urban planning and crucial amenities required for overall welfare, face urban chaos, and are highly vulnerable to natural and human-made disasters. Inadequacy of clean drinking water, clean air, green space, hygienic living conditions, hospitals, and other life-sustaining institutions in urban centers increases the vulnerability.

There is an inverse relationship between the quality of air and the COVID-19 mortality rate (Naqvi et al. 2021). For instance, it has been proven that PM can act as a carrier for SARS-COV-2 (Setti et al. 2020, Borisova & Komisarenko 2021, Pivato et al. 2021). The finer ones (PM<sub>2.5</sub>) have a higher potential to spread the virus (Nor et al. 2021). However, such studies have not considered other possible socioeconomic factors influencing the COVID-19 death toll in urban centers. In that context, the present study examines the relationships between various socioeconomic factors and air quality that influenced the COVID-19 death in Indian cities.

## MATERIALS AND METHODS

For the present study, we have selected 19 Indian cities based on the data available. These cities were Hyderabad, Visakhapatnam, Surat, Bangalore, Kochi, Bhopal, Indore, Jabalpur, Mumbai, Nagpur, Nashik, Pune, Jaipur, Chennai, Coimbatore, Kolkata, Delhi, Patna and Meerut (Fig. 1). These city centers in the corresponding districts are ascribed to be highly populous areas surrounded by less populated rural or sub-urban settlements. These city centers act as the centers of each district's economic, social, and administrative processes. In this study, the district-level COVID-19 death rate reported till 17 March 2022 was collected from the National Center for Disease Control under the Ministry of Health and Family Welfare, Government of India (<https://www.ncdc.gov.in/Mortality/Home.html>). We used the district-level data to represent the COVID-19 death cases of the cities since there were no reliable data available exclusively on the city-level COVID-19 death cases. Further, we assumed that the district-level data primarily represent the death cases reported from the urban centers; it is known that in India, rural COVID-19 death cases were not reported during the first wave and seldom reflected during the second wave (Jha et al. 2022).

The air quality data relating to SO<sub>2</sub>, NO<sub>2</sub>, and PM<sub>10</sub> for ten years (2008-2019) corresponding to each of those cities was collected from the Indian Institute of Tropical Meteorology (IITM) (<https://cpcb.nic.in/annual-report.php>). Using the average values of SO<sub>2</sub>, NO<sub>2</sub>, and PM<sub>10</sub>, the AQI for each city was calculated using the AQI matrix provided by the Central Pollution Control Board at [https://app.cpcbcr.com/ccr\\_docs/AQI%20-Calculator.xls](https://app.cpcbcr.com/ccr_docs/AQI%20-Calculator.xls). The city's

socioeconomic data, area, population, population density, per capita income, and the number of hospitals and hospital beds were collected from <https://www.indiastat.com/>, the annual death rates from <https://csorgi.gov.in/annual-report.html>, and the GDP of the cities from <https://ca.finance.yahoo.com/photos/the-top-15-indian-cities-by-gdp-1348807591-slideshow.html>.

Since there are no routine and exhaustive censuses conducted by the Census of India (Government of India) for 2021, we used the population data for the earlier 3 census periods, 1991, 2001, and 2011, and estimated the current population, making use of the growth trend and the well-known equation is given below.

$$PN = P_0 \left(1 + \frac{r}{100}\right)^t$$

Where  $PN$  is the projected population for the year  $t$ ,

$P_0$  is the initial population

$r$  is the population growth rate, and  $t$  is the year.

The growth rate was calculated using the equation

$$r = \{(P_n - P_0)/P_0\}100\}/N$$

Where  $N$  is the projected year.

Since there are no updated data on the number of hospitals and the total number of hospital beds available,

we used the census data available for the years 2001 and 2011 procured from the office of the registrar general and census commissioner, India <https://censusindia.gov.in/census.website/data/census-tables>. While comparing the data between 2001 and 2011, we found that for some cities, there was a decrease in the number of hospitals and bed availability; hence, we used the latest census data (i.e. 2011) for the analysis.

Multiple correlation analysis was done to reveal the relationships among the variables. Multiple linear regression analysis was carried out to check the affinity of each variable to the COVID death rate since multiple regression analysis is a widely used simple model to predict an outcome variable ( $Y$ ) from a group of predictor variables ( $X_n$ ). In general, multiple regressions can be represented by the following equation:

$$\{Y = B_0 + B_1X_1 + B_2X_2 + \dots B_nX_n + E\}$$

Where,

$Y$  is the dependent predicted variable.

$B_0$  is the Y-intercept or value of  $Y$  when all  $X$  values are 0.

$B_1, B_2$ , and  $B_n$  are the regression coefficients for  $X_1, X_2$ , and  $X_n$ , respectively.

$E$  is the residual standard deviation.

We used the R statistical package (Rstudio 2022.07.2) for all the analyses.




Fig. 1: Map of India showing the location of the 19 selected cities for the study.

## RESULTS AND DISCUSSIONS

The per capita income was highest in Surat, followed by Delhi and Mumbai, with the lowest per capita income in Jabalpur. The number of hospitals was highest in Mumbai, followed by Jabalpur, and the lowest number of hospitals in Jaipur. Mumbai has the highest number of hospital beds available, followed by Jabalpur and Kochi, and the lowest number was in Meerut. Among the 19 cities, the projected population is highest in Bangalore, followed by Mumbai, Delhi, Surat, Hyderabad, Pune, Jaipur, Chennai, Kolkata, Nagpur, Indore, Nashik, Patna, Visakhapatnam, Bhopal, Meerut, Jabalpur, Coimbatore, and Kochi. According to the population projection, Bangalore surpasses all other cities, including Mumbai and Delhi, due to Bangalore's consistently higher population growth rate, escalated possibly by immigration. The decadal population growth between 1991 and 2011 in Bangalore is more than five million. The immigration of highly skilled workers due to employment opportunities in the ICT sector, in the city, growing at an incredible pace (Narayana 2011), the livability of the city, mainly infrastructure amenities, climate, etc., could be the other factors for the population growth in Bangalore. It is important to note that, as suggested by the projection, the tier-2 cities in India, like Surat, Pune, and Jaipur, surpass the tier-1 cities, such as Chennai and Kolkata, in population. Sikarwar (2020) studied the population trend in Kolkata city and found that after 1971, the city's population showed a trend that, after 2001, slanted toward negative growth.

Moreover, cities like Bangalore (Ramachandra 2018) and Hyderabad (Das 2015) showed significant spatial growth during the last decade after the census in 2011. Among the cities, the GDP of Mumbai is the highest, followed by Delhi, Kolkata, and Bangalore. Jabalpur has the lowest GDP level among the cities.

The correlation analysis shows that all variables except the number of hospital beds positively correlate to the COVID-19 death rate in cities. GDP, population, the average death rate of the cities, and the concentration of NO<sub>2</sub> levels in the air are significantly correlated with the death rates. Although the number of hospital beds negatively correlated with the death rate, it is insignificant; very rational in that higher bed availability, a capital stock (Ravaghi et al 2020), and reflected better access to inpatient health care. Similarly, the SO<sub>2</sub>, PM<sub>10</sub>, Air Quality Index, per capita income, number of hospitals, urban area, and population density are positively correlated with the COVID-19 death rate of the cities but do not show any significant trend (Table 1).

The multiple regression analysis results between the COVID-19 death rate in the city with the other variables,

NO<sub>2</sub>, SO<sub>2</sub>, PM<sub>10</sub>, GDP, population, and normal death, of the cities indicated a positive impact. In contrast, AQI, income per capita, number of hospitals, number of hospital beds, area of the city, and the population density of the city show a negative relationship with the death rate. Of this, notably, only the population and area of the city have a significant level of predictability on the COVID-19 death in the city. From the model, COVID-19 deaths in the city can be estimated by the formula:

COVID death =  $5.8 \times 10^3 + 0.0016 \times \text{Population} - 21.9 \times \text{Area}$ , and the model is significant (Adjusted R-squared: 0.7126; p-value of F-statistic: 0.03; Table 2).

There is a significant positive correlation between the COVID-19 death rate and the population ( $r=0.68$ ;  $p=0.0564$ ). Among the selected cities, COVID-19 death was reported highest in Delhi, followed by Pune, Mumbai, and Bangalore, all densely populated megacities in India. The lowest number of personal losses due to COVID-19 was reported in Jabalpur. According to some studies, COVID-19 death cases were higher in cities with denser populations in Algeria (Kadi & Khelfaoui 2020). It has to be noted that in highly populated cities such as Kolkata, Chennai, Surat, and Hyderabad, the reported cases are fewer, which might be due to imprecise statistics or deliberate under-reporting by the authorities to reduce the instances (Jha et al. 2022, Anand et al. 2021, Lancet 2020). It is also reported that there is a lack of reporting the COVID-19 death cases in the older age group in India (Lewnard et al. 2022).

While checking for the significant correlation between the population and other variables, it is found that only two variables that relate to the city's economic status, the GDP and per capita income, are significantly and positively correlated with the population. In India, it is documented that the rural-to-urban mass migration for employment in the non-agriculture sector has increased many folds during the post-globalization of the Indian market economy (Siggel 2010). The city's urban centers offer many opportunities for people from the rural agricultural sector. Due to many factors, including higher input costs, insufficient returns, and an inadequate labor force, the rural agriculture sector has become unprofitable. The immigration of the poor rural labor eventually increased city centers' GDP and per capita income. Among the selected cities, the country's commercial capital, Mumbai, scores the highest GDP, followed by Delhi and Kolkata, and the GDP of Jabalpur is the lowest.

Similarly, the per capita income of Jabalpur is the lowest among the studied cities; Surat has the highest per capita income among these cities. It is reported that there is a direct connection between the availability of transportation networks with COVID-19 spread (Zhu & Guo 2021). The air



Table 1: Correlation matrix of the variables (Pearson correlation coefficients - r).

	CD	NO	SO	PM	AQI	IC	GDP	NH	NB	P	ND	AR	DEN
CD	1.00												
NO	0.53*	1.00											
SO	0.26	0.22	1.00										
PM	0.26	0.57*	-0.20	1.00									
AQI	0.27	0.59*	-0.17	1.00	1.00								
IC	0.30	0.16	0.30	0.06	0.08	1.00							
GDP	0.78*	0.43	-0.05	0.35	0.35	0.35	1.00						
NH	0.28	0.01	-0.11	-0.01	0.02	0.14	0.60	1.00					
NB	-0.02	-0.25	-0.25	-0.23	-0.21	0.05	0.25	0.77*	1.00				
P	0.68*	0.30	0.02	0.34	0.36	0.63*	0.75*	0.46	0.19	1.00			
ND	0.83*	0.45	-0.01	0.37	0.36	0.39	0.88*	0.31	0.04	0.79*	1.00		
AR	0.45	0.20	-0.03	0.13	0.16	0.34	0.51	0.35	0.17	0.77*	0.59*	1.00	
DEN	0.33	0.25	0.17	0.38	0.39	0.62*	0.43	0.15	-0.04	0.57*	0.47	0.04	1.00

CD: COVID deaths, NO = NO<sub>2</sub>, SO = SO<sub>2</sub>, PM = PM<sub>10</sub>, AQI = Air Quality Index, IC = Per capita income, GDP = GDP of the city, NH = Number of Hospitals, NB = Number of Hospital Beds, P = total Population, ND = Annual death in number, AR = Urban area, and DEN = Population density.

transportation networks significantly increased epidemics' spread in general (Colizza et al. 2006). In the case of COVID, due to global and local air travel, cities with higher GDP have a chance of higher spread since it is there that global investment and consequent travelers would be high, as reflected by the initial spread in cities like Mumbai and Delhi.

The highly populated cities in India are the country's economic hubs and industrial centers. Emissions, vehicular, and power generation are also higher in megacities due to higher per capita vehicle availability and higher energy consumption by diverse sectors. Further, household carbon emission is also a crucial issue in India (Beig et al. 2021). All these activities are closely related to air quality, and exposure to highly polluted air is one of the primary reasons of chronic respiratory illness of people, which is a potential factor significantly hiking the chances of death in COVID-affected individuals (Schraufnagel et al. 2019).

Especially since people suffering from other respiratory diseases exposed to air pollution are considered highly vulnerable to COVID-19 death due to comorbidities (Naqvi et al. 2021). Metro cities such as Delhi, Kolkata, and Mumbai are considered the most polluted areas in the world regarding air pollutants (Guttikunda & Nishadh 2022, Guttikunda et al. 2014). As per the recent report by IQAir (<https://www.iqair.com/>), based on the average annual PM<sub>10</sub> concentration, Delhi stands fourth most highly polluted city in the world, followed by Meerut, Patna, Kolkata, Jaipur, Mumbai, Indore, Visakhapatnam, Bhopal, Jabalpur, Pune, Nashik, Hyderabad, Bengaluru, Chennai, and Kochi. However, IQAir does not document the air quality for Surat, Coimbatore, and Nagpur.

A significant positive relationship between AQI and COVID-19 death has been reported by various researchers globally (Wu et al. 2020, Zhu et al. 2020, Conticini et al. 2020). A significant correlation is seen between industrial pollutants like PM<sub>2.5</sub> and COVID-19 deaths in metropolitan and industrial cities in India, as Travaglio et al. (2020) observed. While examining the disability-adjusted life years (DALYs) of the people living in Delhi and Mumbai in 2015, Maji et al. (2017b) found that for the low AQI, the mortality rate increased more than double the mortality observed during 1995 for both the cities. Kolluru et al. (2022) examined the significant relationship between air quality and the COVID spread during the second wave in Delhi. The present study also demonstrates a positive relationship between COVID-19 death and the AQI of cities, but it is not statistically significant. However, there is a significant correlation between the concentration of NO<sub>2</sub> in the urban air and COVID-19 death. The insignificance of AQI's relation with death might be due to the insignificance of the other variables, SO<sub>2</sub> and PM<sub>10</sub>, in determining the AQI of the city, possibly for the inaccuracy in the measurement of the parameters available in these official data sets.

Studies have already proved that prolonged exposure to high concentrations of NO<sub>2</sub> in the atmosphere can lead to other ailments (Ogen 2020), such as hypertension, diabetes (Shin et al. 2020), and other ischemic heart diseases (Qiu et al. 2013). A significant positive correlation between the average annual death rate and the COVID-19 death rate was observed in the present study. It is reported that 12.5% of India's average normal death rate is due to diseases




Fig. 2. Added variable plot for potential drivers for COVID death in urban India. AQI, per capita income, number of hospitals, number of hospital beds, the total area of the city, and population density showcase a negative relationship. Population and average annual death show a positive relationship, whereas per capita income, GDP, and the number of hospitals are almost neutral in association with COVID-19 deaths.

associated with high exposure to poor air quality (Guttikunda & Goel 2013). Further, Maji et al. (2017a) concluded that poor air quality directly influences the annual mortality rate in Delhi alone. As mentioned earlier, it is reported that people exposed to poor AQI are highly vulnerable to mortality due to COVID-19 (Naqvi et al. 2021). Further, the present results indicate that the country still lacks monitoring infrastructure and high-resolution pollution concentration maps relating to urban India's real-time air quality assessment (Agrawal et al. 2021).

It must be noted that all megacities are at the top of the list while checking the number of hospitals and hospital beds in the cities. However, when it comes to the per capita availability of hospital facilities in cities, including

megacities, is very low. It has been found that the average death rate before the COVID-19 scenario shows no significant correlation between the per capita availability of hospital facilities and the annual death rate among the cities.

This again shows that the number of hospital facilities needs to be improved for the population. The number of hospital beds per the census data includes private and public hospitals; in India, it is clear that most of the population accessing private facilities is still a financial concern (Duggal 2007). Recent studies on the Indian healthcare system in the wake of the COVID-19 pandemic also reveal the fact that the privatization of medical facilities has become a financial, geographical, administrative, and logistic burden for the middle and low-income groups, the majority of the

Table 2: Multiple regression analysis results.

Parameter	Estimate	Std.Error	t value	Pr(> t )
(Intercept)	5.89E+03	1.04E+04	0.569	0.5902
NO <sub>2</sub>	1.07E+02	1.00E+02	1.068	0.3264
SO <sub>2</sub>	3.69E+02	2.13E+02	1.733	0.1338
PM <sub>10</sub>	3.80E+01	2.99E+02	0.127	0.9028
AQI	-8.65E+01	4.09E+02	-0.211	0.8395
Income per capita	-1.24E-02	1.10E-02	-1.122	0.3049
GDP	6.18E+00	3.91E+01	0.158	0.8795
Hospital	-1.31E+00	5.32E+00	-0.247	0.8134
Hospital beds	-1.22E-02	2.22E-01	-0.055	0.9579
Population	1.59E-03	6.73E-04	2.359	*0.0564
Normal death	1.23E-01	1.06E-01	1.155	0.292
Area	-2.19E+01	9.85E+00	-2.222	*0.068
Density	-4.17E-01	2.20E-01	-1.899	0.1063

Significance codes: '\*\*' 0.1

Multiple R-squared: 0.9042, Adjusted R-squared: 0.7126

F-statistic: 4.719 on 12 and 6 DF, p-value: 0.03406

population in the country (Bhaduri 2020). The government's role has been highly restricted to financial aid for private parties (Chandwani 2021), while the government promotes the privatization of even such essential facilities.

It is worth noting that the present study does not testify to the inverse relationship between AQI and COVID-19 death, as reported by many others mentioned in the text. We presume this may be due to a shortage of data with adequate precision and the sampling points not being statistically representative. The Added variable plot (Fig. 2) between drivers and COVID-19 deaths in urban India also indicates the absence of precise data. For example, as per Fig. 2, COVID-19 death is negatively influenced by the AQI of the respective cities, which is unreasonable according to the general understanding (Naqvi et al. 2021). Similarly, the negative relationship between the city's population density and COVID-19 deaths, as shown in the plot, is illogical since the disease spread is likely to be enhanced by physical closeness among the people (Kadi & Khelfaoui 2020). As mentioned earlier, data reliability is one of the major lacunae in countries like India, which is one of the major limitations of this type of study. It may be noted that the Death and Birth Registration Authority accepts the inadequacy of reliable data on mortality in India (Basu & Adair 2021).

## CONCLUSION

Several studies from various parts of the globe documented the regional AQI in urban city centers and its influence

in spreading the COVID-19 epidemic. However, we hypothesized that several more underlying socioeconomic factors influence the spread of COVID-19. The present study evaluates COVID-19 in urban India by examining the socioeconomic factors and the AQI of 19 randomly selected urban centers. Our results showcased that the COVID death positively correlates with AQI, population, population density, the average annual death rate in the city, the total area of the urban centers, income per capita, and GDP; and the correlation is significant for GDP, population and average annual death rate of the city. It is also found that among the factors determining the AQI, the concentration of NO<sub>x</sub> is significantly correlated with COVID-19 death. COVID-19 death in cities is negatively correlated with the number of hospital beds in the city. The multiple regression analysis among the selected variables also showed significant positive relationships between COVID-19 death and the population and the total urban area in the city. The study also concluded that the lack of reliable data is a major lacuna while examining the crucial drivers of COVID-19 death in urban India.

## REFERENCES

- Aditya N.S., Nair, A.Y. and Veni, S. 2022. Determining the Effect of Correlation between Asthma/Gross Domestic Product and Air Pollution. In 2022 International Conference on Wireless Communications Signal Processing and Networking (WiSPNET) (pp. 44-48). IEEE
- Agrawal, G., Mohan, D. and Rahman, H. 2021. Ambient air pollution in selected small cities in India: Observed trends and future challenges. *IATSS Res.*, 45(1): 19-30.
- Anand, A., Sandefur, J. and Subramanian, A. 2021. Three new estimates of India's all-cause excess mortality during the COVID-19 pandemic.
- Balachandran, S., Moni, M., Sathyapalan, D.T., Varghese, P., Jose, M.P., Murugan, M.R., Rajan, C., Saboo, D., Nair, S.S., Varkey, R.A. and Balachandran, P. 2022. A comparison of clinical outcomes between vaccinated and vaccine-naïve patients of COVID-19 in four tertiary care hospitals of Kerala, South India. *Clinical Epidemiology and Global Health*, 13: 100971.
- Balakrishnan, K., Cohen, A. and Smith, K.R. 2014. Addressing the burden of disease attributable to air pollution in India: the need to integrate across household and ambient air pollution exposures. *Environ. Health Persp.*, 122(1): A6-A7.
- Balakrishnan, K., Dey, S., Gupta, T., Dhaliwal, R.S., Brauer, M., Cohen, A.J., Stanaway, J.D., Beig, G., Joshi, T.K., Aggarwal, A.N. and Sabde, Y. 2019. The impact of air pollution on deaths, disease burden, and life expectancy across the states of India: The global burden of disease study 2017. *Lancet Planet. Health*, 3(1): e26-e39.
- Basu, J.K. and Adair, T. 2021. Have inequalities in the completeness of death registration between states in India narrowed during two decades of civil registration system strengthening? *Int. J. Equity Health*, 20(1): 1-9.
- Beig, G., Korhale, N., Rathod, A., Maji, S., Sahu, S.K., Dole, S., Latha, R. and Murthy, B.S. 2021. On modeling the growing menace of household emissions under COVID-19 in Indian metros. *Environ. Pollut.*, 272: 115993.
- Benmarhnia, T. 2020. Linkages between air pollution and the health burden from COVID-19: methodological challenges and opportunities. *Am. J. Epidemiol.*, 189(11): 1238-1243.

- Bhaduri, S.D. 2020. Post-COVID healthcare reform in India: What to expect? *J. Family Med. Primary Care*, 9(11): 5427.
- Borisova, T. and Komisarenko, S. 2021. Air pollution particulate matter as a potential carrier of SARS-CoV-2 to the nervous system and/or neurological symptom enhancer: Arguments in favor. *Environ. Sci. Pollut. Res.*, 28(30): 40371-40377.
- Cao, Y., Chen, M., Dong, D., Xie, S. and Liu, M. 2020. Environmental pollutants damage airway epithelial cell cilia: Implications for the prevention of obstructive lung diseases. *Thoracic Cancer*, 11(3): 505-510.
- Chandwani, R. 2021. Stakeholders in the Indian Healthcare Sector. *Vikalpa*, 46(2): 65-70.
- Chen, S. and Jin, H. 2019. Pricing for the clean air: Evidence from the Chinese housing market. *J. Clean. Prod.*, 206: 297-306.
- Colizza, V., Barrat, A., Barthélemy, M. and Vespignani, A. 2006. The role of the airline transportation network in the prediction and predictability of global epidemics. *Proceed. National Acad. Sci.*, 103(7): 2015-2020.
- Conticini, E., Frediani, B. and Caro, D. 2020. Can atmospheric pollution be considered a co-factor in extremely high levels of SARS-CoV-2 lethality in Northern Italy? *Environ. Pollut.*, 261: 114465.
- Das, D. 2015. Hyderabad: Visioning, restructuring and making of a high-tech city. *Cities*, 43: 48-58.
- Duggal, R. 2007. Healthcare in India: changing the financing strategy. *Social Pol. Admin.*, 41(4): 386-394.
- Girdhar, A., Kapur, H., Kumar, V., Kaur, M., Singh, D. and Damasevicius, R. 2021. Effect of COVID-19 outbreak on urban health and environment. *Air Qual. Atmos. Health*, 14(3): 389-397.
- Gupta, A., Bherwani, H., Gautam, S., Anjum, S., Musugu, K., Kumar, N., Anshul, A. and Kumar, R. 2021. Is air pollution aggravating COVID-19 lethality? Exploration in Asian cities using statistical models. *Environ. Dev. Sustain.*, 23(4): 6408-6417.
- Gurjar, B.R., Ravindra, K. and Nagpure, A.S. 2016. Air pollution trends over Indian megacities and their local-to-global implications. *Atmos. Environ.*, 142: 475-495.
- Guttikunda, S. and Nishadh, K.A. 2022. Evolution of India's PM2.5 pollution between 1998 and 2020 using global reanalysis fields coupled with satellite observations and fuel consumption patterns. *Environ. Sci. Atmos.*, 15: 541.
- Guttikunda, S.K. and Goel, R. 2013. Health impacts of particulate pollution in a megacity—Delhi, India. *Environ. Dev.*, 6: 8-20.
- Guttikunda, S.K., Goel, R. and Pant, P. 2014. Nature of air pollution, emission sources, and management in the Indian cities. *Atmos. Environ.*, 95: 501-510.
- Heissel, J.A., Persico, C. and Simon, D. 2022. Does pollution drive achievement? The effect of traffic pollution on academic performance. *J. Hum. Resour.*, 57(3): 747-776.
- Jha, P., Deshmukh, Y., Tumbe, C., Suraweera, W., Bhowmick, A., Sharma, S., Novosad, P., Fu, S.H., Newcombe, L., Gelband, H. and Brown, P. 2022. COVID mortality in India: National survey data and health facility deaths. *Science*, 375(6581): 667-671.
- Jiang, Y., Wu, X.J. and Guan, Y.J. 2020. Effect of ambient air pollutants and meteorological variables on COVID-19 incidence. *Infect. Contr. Hosp. Epidemiol.*, 41(9): 1011-1015.
- Kadi, N. and Khelfaoui, M. 2020. Population density, a factor in the spread of COVID-19 in Algeria: statistic study. *Bull. National Res. Centre*, 44(1): 1-7.
- Karagulian, F., Belis, C.A., Dora, C.F.C., Prüss-Ustün, A.M., Bonjour, S., Adair-Rohani, H. and Amann, M. 2015. Contributions to cities' ambient particulate matter (PM): A systematic review of local source contributions at global level. *Atmos. Environ.*, 120: 475-483.
- Kejela, T. 2020. Probable factors contributing to the fast spread of the novel Coronavirus (COVID-19) in Ethiopia. *J. Infect. Dis. Epidemiol.*, 6: 169.
- Khilnani, G.C. and Tiwari, P. 2018. Air pollution in India and related adverse respiratory health effects: past, present, and future directions. *Curr. Opinion Pulm. Med.*, 24(2): 108-116.
- Kolluru, S.S.R., Nagendra, S.M., Patra, A.K., Gautam, S., Alshetty, V.D. and Kumar, P. 2022. Did unprecedented air pollution levels cause a spike in Delhi's COVID cases during the second wave? *Stoch. Environ. Res. Risk Assess.*, 15: 1-16.
- Lakshmi, P.S. and Suresh, M. 2020. Factors influencing the epidemiological characteristics of pandemic COVID-19: A TISM approach. *Int. J. Healthcare Manag.*, 13(2): 89-98.
- Lancet, T. 2020. COVID-19 in India: the dangers of false optimism. *Lancet (London, England)*, 396(10255): 867.
- Lewnard, J.A., Mahmud, A., Narayan, T., Wahl, B., Selvinayagam, T.S. and Laxminarayan, R. 2022. All-cause mortality during the COVID-19 pandemic in Chennai, India: an observational study. *Lancet Infect. Dis.*, 22(4): 463-472.
- Maji, K.J., Dikshit, A.K. and Deshpande, A. 2017b. Disability-adjusted life years and economic cost assessment of the health effects related to PM2.5 and PM10 pollution in Mumbai and Delhi, India, from 1991 to 2015. *Environmental Science and Pollution Research*, 24(5): 4709-4730.
- Maji, S., Ahmed, S., Siddiqui, W.A. and Ghosh, S. 2017a. Short-term effects of criteria air pollutants on daily mortality in Delhi, India. *Atmos. Environ.*, 150: 210-219.
- Manjula, P.M. 2018. Economics and air pollution analysis of Chennai city. *Int. Rev. Bus. Econ.*, 1(3): 17.
- Naqvi, H.R., Mutreja, G., Shakeel, A. and Siddiqui, M.A. 2021. Spatio-temporal analysis of air quality and its relationship with major COVID-19 hotspot places in India. *Remote Sens. Appl. Soc. Environ.*, 22: 100473.
- Nasir, H., Goyal, K. and Prabhakar, D. 2016. Review of air quality monitoring: A case study of India. *Indian J. Sci. Technol.*, 9(44): 1-7.
- Nor, N.S.M., Yip, C.W., Ibrahim, N., Jaafar, M.H., Rashid, Z.Z., Mustafa, N., Hamid, H.H.A., Chandru, K., Latif, M.T., Saw, P.E. and Lin, C.Y. 2021. Particulate matter (PM2.5) as a potential SARS-CoV-2 carrier. *Scientific Reports*, 11(1): 1-6.
- Ogen, Y. 2020. Assessing nitrogen dioxide (NO2) levels as a contributing factor to Coronavirus (COVID-19) fatality. *Science of the Total Environment*, 726: 138605.
- Pandey, A., Brauer, M., Cropper, M.L., Balakrishnan, K., Mathur, P., Dey, S., Turkoglu, B., Kumar, G.A., Khare, M., Beig, G. and Gupta, T. 2021. Health and economic impact of air pollution in the states of India: The global burden of disease study 2019. *Lancet Planet. Health*, 5(1): e25-e38.
- Pivato, A., Amoroso, I., Formenton, G., Di Maria, F., Bonato, T., Vanin, S., Marion, A. and Baldovin, T. 2021. Evaluating the presence of SARS-CoV-2 RNA in the particulate matter during the peak of COVID-19 in Padua, northern Italy. *Sci. Total Environ.*, 784: 147129.
- Priya, V., Samyuktha, S., Fernandez, S., Biju, A., Anand, G.S., Venniyoor, V., Mathew, K.A. and Marwaha, V. 2021. Knowledge, attitude, and practice towards COVID-19 among social media users in an engineering college in South India. *J. Commun. Dis.*, 3(2): 35-42.
- Qiu, H., Yu, I.T.S., Wang, X., Tian, L., Tse, L.A. and Wong, T.W. 2013. Season and humidity dependence of the effects of air pollution on COPD hospitalizations in Hong Kong. *Atmos. Environ.*, 76: 74-80.
- Raj, P.P.N. and Azeez, P.A. 2020. Economic Growth, the Pandemic, and the Convalescing Nature. Retrieved from <https://businessseconomics.in/economic-growth-pandemic-and-convalescing-nature>.
- Ravaghi, H., Alidoost, S., Mannion, R. and Bélorgeot, V.D. 2020. Models and methods for determining the optimal number of beds in hospitals and regions: A systematic scoping review. *BMC Health Services Res.*, 20(1): 1-13.



- Schraufnagel, D.E., Balmes, J.R., Cowl, C.T., De Matteis, S., Jung, S.H., Mortimer, K., Perez-Padilla, R., Rice, M.B., Riojas-Rodriguez, H., Sood, A. and Thurston, G.D. 2019. Air pollution and noncommunicable diseases: A review by the Forum of International Respiratory Societies' Environmental Committee, Part 2: Air pollution and organ systems. *Chest*, 155(2): 417-426.
- Setti, L., Passarini, F., De Gennaro, G., Barbieri, P., Lichen, S., Perrone, M.G., Piazzalunga, A., Borelli, M., Palmisani, J., Di Gilio, A. and Rizzo, E. 2020. Potential role of particulate matter in the spreading of COVID-19 in Northern Italy: first observational study based on initial epidemic diffusion. *BMJ Open*, 10(9): e039338.
- Shin, S., Bai, L., Oiamo, T.H., Burnett, R.T., Weichenthal, S., Jerrett, M., Kwong, J.C., Goldberg, M.S., Copes, R., Kopp, A. and Chen, H. 2020. Association between road traffic noise and incidence of diabetes mellitus and hypertension in Toronto, Canada: a population-based cohort study. *J. Am. Heart Association*, 9(6): e013021.
- Siggel, E. 2010. The Indian informal sector: the impact of globalization and reform. *Int. Lab. Rev.*, 149(1): 93-105.
- Sikarwar, A. and Rani, R. 2020. Assessing the immediate effect of COVID-19 lockdown on air quality: A case study of Delhi, India. *Journal of Environmental Geography*, 13(3-4): 27-33.
- Sikarwar, A. 2020. Falling population growth and subsequent land use pattern: A study of Kolkata. In Chattopadhyay, A. and Ghosh, S. (eds), *Population Dynamics in Eastern India and Bangladesh*, Springer, Singapore, pp. 383-392.
- Sivarethinamohan, R., Sujatha, S., Priya, S., Gafoor, A. and Rahman, Z. 2021. Impact of air pollution in health and socioeconomic aspects: review on future approach. *Mater. Today Proceed.*, 37: 2725-2729.
- Subbarao, K. and Mahanty, S. 2020. Respiratory virus infections: understanding COVID-19. *Immunity*, 52(6): 905-909.
- Travaglio, M., Yu, Y., Popovic, R., Selley, L., Leal, N.S. and Martins, L.M. 2021. Links between air pollution and COVID-19 in England. *Environ. Pollut.*, 268: 115859.
- Venkataraman, C., Brauer, M., Tibrewal, K., Sadavarte, P., Ma, Q., Cohen, A., Chaliyakunnel, S., Frostad, J., Klimont, Z., Martin, R.V. and Millet, D.B. 2018. Source influence on emission pathways and ambient PM 2.5 pollution over India (2015-2050). *Atmos. Chem. Phys.*, 18(11): 8017-8039.
- World Health Organization (WHO). 2021. New WHO Global Air Quality Guidelines Aim to Save Millions of Lives from Air Pollution. Retrieved from <https://www.who.int/news/item/22-09-2021-new-who-global-air-quality-guidelines-aim-to-save-millions-of-lives-from-air-pollution> (accessed October 1, 2021)
- Wu, X., Nethery, R.C., Sabath, M.B., Braun, D. and Dominici, F. 2020. Exposure to air pollution and COVID-19 mortality in the United States: A nationwide cross-sectional study. *MedRxiv*, 11: 225.
- Zhu, P. and Guo, Y. 2021. The role of high-speed rail and air travel in the spread of COVID-19 in China. *Travel Med. Infect. Dis.*, 42: 102097.
- Zhu, Y., Xie, J., Huang, F. and Cao, L. 2020. Association between short-term exposure to air pollution and COVID-19 infection: Evidence from China. *Sci. Total Environ.*, 727: 138704.