# Cardiovascular Disease Prevention Practices by U.S. Physicians for Patients with Diabetes

James B. Meigs, MD, MPH, Randall S. Stafford, MD, PhD

OBJECTIVE: Cardiovascular diseases account for the majority of morbidity and mortality in patients with type 2 diabetes mellitus. We describe patterns of cardiovascular disease primary prevention practices used for patients with diabetes by U.S. office-based physicians.

MEASUREMENTS AND MAIN RESULTS: We analyzed a representative sample of 14,038 visits from the 1995 and 1996 National Ambulatory Medical Care Surveys (NAMCS), including 1,489 visits by patients with diabetes. Physicians completed visit forms describing diagnoses, demographics, services provided, and current medications. Diabetes was defined by diagnostic codes; patients with ischemic heart disease or younger than 30 years were excluded. We estimated national visit volumes by extrapolation using NAMCS sampling weights. Independent determinants of prevention practices were evaluated using multiple logistic regression. Actual visits sampled translated into an estimated 407 million office visits in 1995 and 1996, of which 44.8 million (11%) were by patients with diabetes. Overall, patients with diabetes received more cardiovascular disease prevention services than patients without diabetes, including cholesterol reduction (8% vs 5%, P < .001) and exercise counseling (22% vs 13%, P < .001), blood pressure measurement (82% vs 72%, P < .001), and aspirin prescription (5% vs 2%, P < .001). Patients with diabetes and hyperlipidemia were more likely to receive lipid-lowering medications than patients without these diagnoses (67% vs 51%, P = .007), but those who had diabetes and hypertension or who smoked were no more likely than those without to receive antihypertensive medications or smoking cessation counseling, respectively. These effects persisted in multiple logistic regression analyses controlling for potential confounders.

CONCLUSIONS: Patients with diabetes visiting U.S. physicians in 1995 and 1996 received somewhat more cardiovascular disease prevention services than patients without diabetes. Absolute rates of services, however, remained lower than desired based on national recommendations. Current evidence suggests that wider implementation of these recommendations can be expected to reduce the burden of cardiovascular disease in patients with diabetes.

KEY WORDS: cardiovascular disease; type 2 diabetes; prevention.

J GEN INTERN MED 2000;15:220-228.

Received from the General Medicine Division, Massachusetts General Hospital, Medical Services, and Department of Medicine, Harvard Medical School, Boston, Mass (JBM, RSS).

Address correspondence and reprint requests to Dr. Meigs: General Medicine Division, Massachusetts General Hospital, 50 Staniford St., 9th Floor, Boston MA 02114 (e-mail: jmeigs@partners.org).

Diabetes mellitus is a common chronic condition affecting almost 8% of the U.S. population between the ages of 20 and 74 years. Care for diabetes is the fourth most common reason for ambulatory physician visits, the majority of which are to office-based primary care physicians. Well over 90% of people with diabetes have type 2 diabetes mellitus. Atherosclerotic cardiovascular disease (CVD) accounts for the majority of morbidity and mortality. In people with type 2 diabetes. Diabetes and its CVD complications are costly in economic terms as well as in human suffering. Estimated health care expenditures for diagnosed diabetes total nearly \$100 billion. At least 26% of costs attributable to diabetes in one managed care organization were due to CVD complications. 10

Clinical management of diabetes has traditionally focused on glycemic control. However, observational associations between blood glucose levels and CVD complications have been inconsistent, 11,12 and randomized, controlled trials of intensive glycemic control in patients with type 2 diabetes have not shown significant reductions in CVD event rates. 13-15 In contrast, clinical trials subgroup analyses of patients with type 2 diabetes have consistently demonstrated reductions in CVD event rates with intensive management of hyperlipidemia<sup>16-20</sup> and hypertension,<sup>21-26</sup> and with use of aspirin prophylaxis. 24,27,28 Cigarette smoking also has a pernicious effect among those with diabetes.<sup>29</sup> While interventions to improve these CVD risk factors have been widely recommended, 30-35 little is known about physician practices related to CVD risk factor modification among people with diabetes. In this report, we analyzed data from the 1992, 1995, and 1996 National Ambulatory Medical Care Surveys (NAMCS) to assess the prevalence, determinants, and trends over time in primary CVD prevention practices provided to patients with diabetes in the United States.

#### **METHODS**

We analyzed data from the NAMCS for 1992, 1995, and 1996.<sup>36-38</sup> Conducted by the National Center for Health Statistics (NCHS), these surveys assess the practices of U.S. office-based physicians. The American Medical Association and American Osteopathic Association master lists of all U.S patient care physicians were used to select a random physician sample stratified by geographic area and specialty. Annual participation rates among eligible physicians averaged 72%. Patient visits during a randomly selected week were sampled for each participating physician in each year. For the years we analyzed, the survey included 34,606 (1992), 36,875 (1995),

and 29,805 (1996) outpatient visits to 1,558 (1992), 1,883 (1995), and 1,500 (1996) physicians.

Participating physicians completed NAMCS encounter forms detailing the specific clinical services provided, new and ongoing medications, and patient demographics and diagnoses. For each recorded visit the NCHS provided a visit weight calculated from the physician and visitsampling rates, adjusted for nonresponse. Statistical aggregation using these visit weights allows extrapolation so that practice patterns and the volume of patient visits can be estimated at a national level. We modified these weights according to Potthoff et al.39 To derive effective sample sizes for use in statistical testing, we used NCHS relative standard errors to estimate the 95% confidence intervals for these national estimates.<sup>36-38</sup> Validation studies performed in earlier years confirm the accuracy of the information obtained.40 Missing data were limited to approximately 5% of most data fields. The NAMCS is a cross-sectional study and does not allow specific patients to be followed longitudinally. The unit of analysis is the patient visit, rather than the patient. The data do not include outpatient care provided by hospital outpatient departments, some health centers, emergency departments, or by nonphysician providers.

Our primary analytic goal was to compare rates of CVD prevention services provided to patients with diabetes and to those without diabetes. We identified visits by patients with diabetes mellitus by an International Classification of Diseases, Math Revisions, Clinical Modification (ICD-9-CM) diagnostic code of 250, by a NAMCS-specific "reason for visit code" for diabetes, by a check-off indicator for diabetes (in 1995 and 1996), or by the report of oral hypoglycemic or insulin therapy. We excluded visits by patients with reported coronary heart disease and visits by patients under the age of 30 years in order to assess practices related to primary CVD prevention for those most likely to have type 2 diabetes or type 1 diabetes of long duration. Patients with diabetes were classified into three groups according to whether they were treated with insulin, with an oral hypoglycemic agent alone, or with neither of these medications; the latter we defined as diet-treated diabetes. We further limited our analysis to visits made to primary care physicians, endocrinologists, and cardiologists, the specialties most likely to be expected to provide diabetes prevention services. Using these criteria, our analysis for 1995 and 1996 included 14,038 visits: 1,489 by patients with diabetes and 12,549 by patients without diabetes.

Our principal outcome measures were the provision of several specific ambulatory services related to CVD prevention. We assessed whether a variety of counseling and diagnostic testing services were performed, and whether patients were reported to be taking specific classes of medications. We analyzed physician practices as they applied to all patient visits in aggregate and to clinical subsets of patients for whom specific services would be most applicable. For all patients we examined exercise counsel-

ing, cholesterol reduction counseling, weight reduction counseling, cholesterol testing, blood pressure measurement, and the report of aspirin prescription. We also examined the following subsets: patients with hypertension (blood pressure measurement, angiotensin-converting enzyme [ACE] inhibitor use,  $\beta$ -blocker use, and thiazide diuretic use, and use of any blood pressure medication), patients with hyperlipidemia (cholesterol counseling and lipid-lowering medication use), cigarette smokers (smoking cessation counseling), and patients with obesity (weight reduction advice, exercise counseling, and obesity medications). For each service we used the  $\chi^2$  test to evaluate whether the three groups of patients with diabetes differed statistically from patients without diabetes.<sup>41</sup>

We further defined clinical, demographic, and physician characteristics hypothesized to influence the provision of CVD prevention services, including the presence of hypertension, smoking, obesity, and hyperlipidemia; patient age, race, and gender; geographic region of the country (four regions), whether the patient resided in a metropolitan statistical area (yes or no), and physician specialty. We used these characteristics both to describe patients with diabetes and to account for their potentially confounding effects in comparisons between patients with and patients without diabetes. We evaluated independent predictors of CVD services with a series of multiple logistic regression models assessing the effect of diabetes versus no diabetes on the likelihood of services being provided at a specific patient visit, controlling for effects of potentially confounding variables. 42 We ran logistic regression models using generalized estimating equations; this method accounts for potential correlation between observations (patient visits) recorded by the same physician.<sup>43</sup> Effective sample size weights were also included in regression models to further account for the complex NAMCS sampling design. We tested for statistical interactions between major demographic factors and provision of prevention services by introducing first-order interaction terms (e.g., race × diabetes) into regression models.

We also used data from 1992 to gauge whether changes have occurred since then in CVD prevention practices for patients with or patients without diabetes. We calculated the difference in the percentage of visits in which specific services were provided between 1992 and 1995–1996, assessing statistical differences with the  $\chi^2$  test for trend. We used SAS for all analyses,<sup>44</sup> and defined statistical significance as a two-tailed P < .05.

## **RESULTS**

In 1995 and 1996 actual visits translated into an estimated 407 million (95% confidence interval [CI], 371 to 443 million) visits to office-based primary care physicians, endocrinologists, and cardiologists in the United States by patients over 30 years of age not known to have coronary heart disease. Of these, 44.8 million (11%) were

by patients with diabetes mellitus. Patient characteristics are displayed in Table 1. Among patients with diabetes, 16% were treated with insulin, 35% with oral hypoglycemic medications alone, and 49% with no specific hypoglycemic medication. Patients with diabetes were more likely to be seen by endocrinologists and less likely to be seen by family and general practitioners compared with patients without diabetes. Patients with diabetes were older, and a greater proportion were not white, compared with those without diabetes. Hyperlipidemia (9%), hypertension (49%), and obesity (21%) were more common among patients with diabetes than among those without diabetes, but similar proportions of patients were cigarette smokers (11%).

The presence of diabetes made it moderately more likely for patients to receive some CVD prevention services during an office visit (Table 2). Diabetic patients were about twice as likely to receive counseling about exercise (21.7% of visits by diabetes patients vs 12.8% of visits by patients without diabetes, P=.001) and weight reduction (16.6% vs 7.1%, P=.001). Physicians were more likely to measure blood pressure for patients with diabetes compared with those without diabetes (81.9% vs 72.3%, P=.001). Among patients with diabetes, those treated with oral hypoglycemic medications were more likely to receive these services than were those treated with insulin or diet only. Although cholesterol advice (8.1 vs 4.7%, P=.001) and aspirin use (4.8% vs 2.3%, P=.001) were more likely to occur among patients with diabetes, these differences were limited to medication-treated patients

with diabetes. The likelihood of cholesterol testing among patients with diabetes (8.8% vs 7.4%, P = .001) was not significantly different from that among patients without diabetes.

Among patients with hypertension, blood pressure measurement was no more likely to occur among all patients with diabetes than among those without diabetes (86.4% vs 83.7%, P = .1), although blood pressure was more likely to be measured among the subgroup treated with oral hypoglycemic medications (Table 2). Overall, prescription of medication for hypertension was equally likely among patients with and patients without diabetes (52.7% vs 51.6%, P = .7), but use of antihypertensive medications was more common among medication-treated diabetic patients and less likely among patients treated with diet only. The classes of medications used to treat hypertension differed between patients with and patients without diabetes: hypertensive patients with diabetes were more likely to be taking ACE inhibitors (22.8% vs 18.2%, P = .02), but less likely to be taking  $\beta$ -blockers (9.3% vs 12.6%, P = .04) and thiazide diuretics (10.6% vs 14.3%, P = .03).

For patients with hyperlipidemia, there was no difference in the provision of cholesterol counseling between those with and those without diabetes (33.7% vs 36.5%, P=.6; Table 2). The use of lipid-lowering medications was more likely among patients with diabetes (66.9% vs 51.3%, P=.007), but was most likely among insulintreated diabetes patients. For patients smoking cigarettes, there were no differences between those with and those

Table 1. Characteristics of Office Visits by Patients With and Patients Without Diabetes, National Ambulatory Medical Care Surveys (NAMCS) 1995–1996

		Patients with Diabetes		
Characteristic	Patients Without Diabetes	Without Reported Hypoglycemic Medications	Treated with Oral Hypoglycemic Medications	Treated with Insulin
NAMCS observations	12,549	750	499	240
Total U.S. visits (in millions)	362.1	21.9	15.7	7.2
Percentage of all visits	89.0	5.4	3.9	1.8
Physician specialty, %				
Endocrinology	0.8	3.4*	4.2*	10.6*
Cardiology	3.3	4.0	2.7	4.9
Family practice/general practice	57.5	$51.2^{\dagger}$	$50.4^{\dagger}$	41.2*
General internal medicine	38.4	41.4	42.6	43.3
Female, %	60.5	60.6	55.1 <sup>‡</sup>	60.3
Age, %				
30-49 y	44.2	20.8*	21.9*	22.4*
50-64 y	24.3	$29.0^{\ddagger}$	32.9*	31.9*
65–74 y	17.3	31.5*	$23.8^{\dagger}$	$25.0^{\ddagger}$
75+ y	14.4	$18.8^{\dagger}$	21.4*	$20.6^{\ddagger}$
Nonwhite, %	15.0	21.9*	$20.2^{\dagger}$	26.3*
Smoker, %	12.7	12.2	$9.0^{\ddagger}$	10.1
Obesity, %	10.6	18.2*	24.9*	19.2*
Hyperlipidemia, %	5.7	7.3	10.2*	9.1
Hypertension, %	27.5	47.8*	48.3*	51.5*

 $<sup>^{*}\</sup>mathrm{P} < .001$ , compared with patients without diabetes.

 $<sup>^{\</sup>dagger}$ .001 < P < .01.

<sup>‡.01 &</sup>lt; P < .05.

Table 2. Cardiovascular Disease Prevention Services Among Patients With and Without Diabetes Mellitus, National Ambulatory Medical Care Surveys (NAMCS) 1995–1996

		Patients with Diabetes		
Cardiovascular Disease Prevention Service	Patients Without Diabetes	Without Reported Hypoglycemic Medications	Treated with Oral Hypoglycemic Medications	Treated with Insulin
Of all visits, percentage receiving				
Exercise counseling	12.8	16.2*	$30.2^{\dagger}$	$19.9^{\ddagger}$
Cholesterol advice	4.7	5.8	$12.8^{\dagger}$	4.8
Cholesterol test	7.4	8.6	9.6	7.4
Weight reduction advice	7.1	$12.5^{\dagger}$	$24.0^{\dagger}$	$13.0^{\ddagger}$
Blood pressure measurement	72.3	77.4*	86.8 <sup>†</sup>	85.1 <sup>†</sup>
Aspirin	2.3	2.7	$6.6^{\dagger}$	$7.2^{\dagger}$
Of visits by patients with hypertension, percentage receiving				
Blood pressure measurement	83.7	83.8	90.9*	84.4
Blood pressure medications	51.6	36.5 <sup>†</sup>	69.6 <sup>†</sup>	64.1*
Angiotensin-converting enzyme				
inhibitors	18.2	13.1	$34.4^{\dagger}$	23.1
β-Blockers	12.6	9.0	8.8	11.1
Thiazide diuretics	14.3	8.6*	14.3	8.8
Of visits by patients with hyperlipidemia, percentage receiving				
Cholesterol advice	36.5	28.2	45.9	17.1
Lipid-lowering medications	51.3	63.8	63.6	82.5*
Of visits by patients who are smokers, percentage receiving smoking				
counseling	26.9	25.8	21.6	19.8
Of visits by patients with obesity, percentage receiving				
Weight reduction counseling	38.3	43.2	$57.0^{\dagger}$	41.2
Exercise counseling	33.4	34.5	$49.8^{\ddagger}$	39.0
Obesity medications	5.9	0.3*	0.5*	0.8

<sup>\*.01 &</sup>lt; P < .05, compared with patients without diabetes.

without diabetes (23.6% vs 26.9%, P = .5) in the likelihood of receiving smoking cessation counseling.

Among patients reported to be obese, weight reduction advice (48.7% vs 38.3%, P=.006) and exercise counseling (41.6% vs 33.4%, P=.03) were more likely for patients with diabetes (Table 2). These differences were greatest for diabetic patients treated with oral agents. Pharmacologic treatment of obesity was rare, but was more than 10 times less common in patients with diabetes than in those without diabetes (0.5% vs 5.9%, P=.001).

After adjustment for potential confounders, patients with diabetes remained generally more likely to receive services compared with patients without diabetes (Table 3). The magnitude of these differences, however, was often modest. For all patients, those with diabetes were 12% to 98% more likely to receive exercise counseling, weight reduction advice, blood pressure measurement, cholesterol counseling, and aspirin therapy. The presence of diabetes did not affect the likelihood of cholesterol testing. For patients with hypertension, there was no independent effect of diabetes on blood pressure measurement or the overall

use of blood pressure medications. Significant differences in the selection of medications persisted, with diabetic patients more likely to receive ACE inhibitors and less likely to receive thiazide diuretics. Patients with diabetes and hyperlipidemia were twice as likely to be taking lipid-lowering medications as nondiabetic patients, but were no more likely to be counseled about cholesterol reduction. Smokers with diabetes were equally likely to receive smoking cessation advice as those without diabetes. Obese patients were just over 50% more likely to receive exercise counseling and weight reduction counseling if they had diabetes. Obese patients with diabetes, however, remained much less likely to be taking medications specifically intended to treat obesity.

Several other factors also influenced the provision of CVD prevention services independent of diabetes and other characteristics. Younger patients and male patients were generally more likely to receive prevention services. Patients residing in metropolitan areas tended to be less likely to receive services. Patients with hypertension, hyperlipidemia, or obesity were in general more likely to receive most services. Race, region of the country, and physician

<sup>†</sup>P < .001.

<sup>‡.001 &</sup>lt; P < .01.

Table 3. Independent Impact of Diabetes on Cardiovascular Disease Prevention Practices, National Ambulatory Medical Care Surveys (NAMCS) 1995–1996

	Adjusted Odds Ratio (95% CI)
Dependent Variable	Diabetes Versus No Diabetes*
Total population	
Exercise counseling	1.61 (1.37 to 1.90)
Weight reduction advice	1.98 (1.59 to 2.45)
Blood pressure measurement	1.12 (1.04 to 1.28)
Aspirin	1.46 (1.03 to 2.07)
Cholesterol advice	1.28 (0.99 to 1.66)
Cholesterol test	0.90 (0.71 to 1.15)
Patients with hypertension	
Blood pressure measurement	1.07 (0.84 to 1.37)
Blood pressure medications	1.04 (0.85 to 1.27)
Angiotensin-converting	
enzyme inhibitors	1.36 (1.07 to 1.74)
β-Blockers	0.71 (0.51 to 0.99)
Thiazide diuretics	0.70 (0.51 to 0.96)
Patients with hyperlipidemia	
Cholesterol advice	0.96 (0.59 to 1.56)
Lipid-lowering medications	1.87 (1.12 to 3.11)
Patients who are smokers	
Smoking counseling	0.81 (0.50 to 1.29)
Patients with obesity	
Weight reduction counseling	1.53 (1.12 to 2.11)
Exercise counseling	1.52 (1.09 to 2.10)
Obesity medications	0.11 (0.02 to 0.68)

<sup>\*</sup>Model included the additional independent variables of age group, physician speciality, gender, race, region, metropolitan statistical area, and clinical characteristics including hypertension, obesity, smoking status, and hyperlipidemia. CI indicates confidence interval.

specialty did not have significant impact on provision of prevention services. We found no consistent statistical interaction between diabetes and patient characteristics on provision of most prevention services.

Cardiovascular disease prevention practices in 1995 and 1996 were similar to those in 1992 (Table 4). Both for patients with and for patients without diabetes, changes over time were small in magnitude and did not necessarily suggest improvements. For diabetic patients provision of cholesterol advice, weight reduction advice, and antihypertensive medication prescription (among those with hypertension) actually became less common. For patients without diabetes, there were some improvements (exercise counseling and blood pressure measurement in all patients, and exercise counseling and weight reduction advice in obese patients), but also simultaneous declines in several other services including provision of cholesterol advice or antihypertensive medication prescription.

#### **DISCUSSION**

Despite their substantial risk of developing and dying from coronary heart disease or stroke, we have demon-

Table 4. Temporal Differences in Cardiovascular
Prevention Services between 1992 and 1995–1996 National
Ambulatory Medical Care Surveys (NAMCS)

Cardiovascular Disease Prevention ServicePatients Without DiabetesPatient With With DiabetesOf all visits, percentage difference Exercise counseling Cholesterol advice $2.8^*$ $-1.4$ $-1.4$ Cholesterol test Weight reduction advice Blood pressure measurement $-0.1$ $-1.1$
Exercise counseling $2.8^*$ $-1.4$ Cholesterol advice $-1.3^{\dagger}$ $-4.4^{\ddagger}$ Cholesterol test $1.3^{\ddagger}$ $-1.1$ Weight reduction advice $-0.1$ $-4.9^{\ddagger}$ Blood pressure measurement $2.3^{\ddagger}$ $3.3$
Exercise counseling $2.8^*$ $-1.4$ Cholesterol advice $-1.3^{\dagger}$ $-4.4^{\ddagger}$ Cholesterol test $1.3^{\ddagger}$ $-1.1$ Weight reduction advice $-0.1$ $-4.9^{\ddagger}$ Blood pressure measurement $2.3^{\ddagger}$ $3.3$
Cholesterol test $1.3^{\ddagger}$ $-1.1$ Weight reduction advice $-0.1$ $-4.9^{\ddagger}$ Blood pressure measurement $2.3^{\ddagger}$ $3.3$
Weight reduction advice $-0.1$ $-4.9^{\ddagger}$ Blood pressure measurement $2.3^{\ddagger}$ $3.3$
Blood pressure measurement $2.3^{\ddagger}$ $3.3$
Blood pressure measurement $2.3^{\ddagger}$ $3.3$
Aspirin 0.6 0.5
Of visits by patients with
hypertension, percentage difference
Blood pressure measurement 1.0 3.5
Blood pressure medications $-6.7^*$ $-19.0^*$
Angiotensin-converting enzyme
inhibitors $-4.1^{\dagger}$ $-4.2$
β-Blockers 0.2 1.5
Thiazide diuretics $-1.1$ $-12.9$ *
Of visits by patients with hyperlipidemia, percentage difference
Cholesterol advice $-13.0^{\dagger}$ $-2.7$
Lipid-lowering medications $-10.1^{\ddagger}$ 0.1
Of visits by patients who are smokers, percentage difference
Smoking counseling $-1.9$ 1.6
Of visits by patients with obesity, percentage difference
Weight reduction counseling $8.3^{\dagger}$ $2.0$
Exercise counseling 9.4* 7.2
Obesity medications 4.7* 0.4

 $<sup>^{*}\</sup>mathrm{P}<.001$ , testing the hypothesis that there is no difference between 1995–1996 and 1992 rates.

strated that patients with diabetes mellitus visiting U.S. physicians inconsistently and only modestly received more outpatient CVD prevention services than patients without diabetes. We have specifically focused on primary prevention, or services provided to patients without diagnosed CVD. Although patients with diabetes were more likely to receive services related to hypertension, hyperlipidemia, physical activity, and obesity, this was not consistent across different diabetes treatment groups. Smokers with diabetes were no more likely to receive counseling for smoking cessation than smokers without diabetes.

We defined diabetes using several coding options available on the NAMCS data record and excluded younger patients to minimize inclusion of those with type 1 diabetes of shorter duration, in whom metabolic or microvascular complications are the principal causes of morbidity and mortality. We included patients likely to have type 2 diabetes, treated in a variety of ways, including insulin, oral hypoglycemic medication, and those not receiving di-

<sup>†.001 &</sup>lt; P < .01.

<sup>‡.01 &</sup>lt; P < .05.

abetes-specific medications. We stratified our analyses by treatment group as a proxy for severity, hypothesizing that greater severity of diabetes would lead to more aggressive CVD prevention activities. Contrary to our expectations, CVD risk factor interventions did not necessarily increase in intensity for patients whose diabetes was treated more intensively. In general, diabetic patients treated with oral hypoglycemic medications tended to receive more services than patients receiving diet therapy only or insulin therapy. It may be that among patients on insulin therapy the focus on glycemic control superceded attention to reducing other risk factors.11 As patients with diabetes not receiving diabetes-specific therapy are also at substantially elevated risk of CVD,46 rates of CVD prevention practices equivalent to those provided to patients treated with diabetes medication were probably appropriate.

Although CVD prevention activities occurred more often among patients with diabetes, absolute rates of provision of many prevention services were still lower than expected based on national treatment recommendations. The American Diabetes Association, the Joint Committee on Detection, Evaluation, and Treatment of High Blood Pressure, and the National Cholesterol Education Program specifically identify diabetes as a high CVD-risk state; consensus statements or clinical practice recommendations encouraging aggressive treatment of hypertension, hyperlipidemia, and cigarette smoking were all published and available to clinicians during or before our sampling dates 1992-1996.30,34,47-49 Clinical trials results strongly supporting these recommendations were published subsequently. 16-20,22-26 While diabetic patients with hypertension had blood pressure measurements at the majority of office visits, at only half of these visits did patients receive an antihypertensive prescription, and, remarkably, rates of antihypertensive prescription fell by nearly 20% between 1992 and 1995. Rates of treatment for hyperlipidemia among diabetic patients were relatively high, with patients receiving lipid-lowering prescriptions at two thirds of visits. Cigarette smoking has especially adverse effects among patients with diabetes, yet cigarette smokers with diabetes received smoking cessation counseling at only a quarter of visits. Rates of prophylactic aspirin prescription were very low. While aspirin therapy guidelines for patients with diabetes were published only recently,33 clinical trials results demonstrating efficacy of prophylactic aspirin therapy in diabetes subgroups were available before or during our sampling period.<sup>27,28</sup> We found no significant increases, and in many cases significant decreases, in most CVD prevention practices between 1992 and 1995-1996.

The "right" rate for provision of CVD prevention services is not known. The unit of analysis in the NAMCS data is an individual patient visit; patients with more visits (e.g., diabetic patients) might be expected not to have all services provided at every visit, and therefore appear to have lower rates of some services provided. New and ongoing medication prescriptions are recorded at every

NAMCS visit; thus, rates of medication prescription in the NAMCS data are likely to reflect physician prescription practices for individual patients over time. The low rate of aspirin prescription, for example, is probably close to an accurate measure of limited use of this important prevention intervention. Low rates of patient screening and counseling services do probably reflect that these services were not provided at all of the relatively frequent visits by diabetic patients (estimated at about 5 ambulatory visits per year<sup>45</sup>). Also, some counseling and screening interventions may have been provided by nonphysicians (e.g., nurses or diabetes educators); provision of these services was not captured in the NAMCS data, and their rates of provision may be underestimated. Conversely, counseling and screening services may be overreported by physicians in the setting of the NAMCS study, where the Hawthorne effect causes physicians to improve practice behaviors while under observation. Nonetheless, it can be argued that certain counseling or screening services should be provided at most visits. For smoking cessation counseling, in particular, we think that every visit should provide the opportunity to encourage patients to overcome this deadly adverse health behavior. Another limitation of our NAMCS sample is our focus on visits to primary care physicians, endocrinologists, and cardiologists. Patterns of care provided to patients of these physicians may differ from patterns in the general population of patients with and patients without diabetes.

These NAMCS data add to accumulating evidence from local community surveys or from chart review studies that evidence-based guidelines are inconsistently applied to the care of patients with diabetes. It is notable that no published studies of diabetes care prior to this report have included any comprehensive, detailed information on CVD prevention practices; prior research in this area has focused almost exclusively on care related to glycemic control and screening for microvascular complications. The following few studies provided the only data we could find directly related to CVD prevention in diabetes. In the San Antonio Heart Study, fewer than 10% of participants with diabetes and hyperlipidemia surveyed in 1979 to 1988 reported hyperlipidemia treatment.50 In a 1991 community survey in Michigan, over 60% of diabetes patients reported having hypertension, and of these, about 80% reported receiving antihypertensive therapy. 51 Among nearly 400 diabetes patients cared for in two different large HMOs, chart review revealed that 86% to 90% of patients had at least 1 or 2 blood pressure checks during a 1- or 2-year sampling window during 1990 to 1994, but only 32% to 60% had at least 1 cholesterol determination. 52,53 Chart review data from 790 diabetes patients cared for at 42 different institutions between 1989 and 1993 revealed that 62% to 90% had a cardiovascular examination (presumably including blood pressure assessment) and 36% to 64% had an annual lipid profile. Among a national cohort of patients surviving acute myocardial infarction in 1994 or 1995, only 47% of 11,716 patients

with diabetes were discharged on  $\beta$ -blocker therapy; these patients were less likely (relative risk, 0.93; 95% CI, 0.91 to 0.95) than patients without diabetes to have received  $\beta$ -blockers. <sup>54</sup> All these investigators concluded that there is ample room for improvement in diabetes care overall. Our study provides specific data highlighting the need for improved provision of potentially life-saving CVD-specific prevention practices.

All major clinical guidelines focused on CVD prevention were available before or during our study period, and since 1995 a large number of clinical trials have solidified the evidence base for CVD risk factor management in diabetes. It is quite clear that aggressive management of hyperlipidemia almost certainly prolongs the lives of patients with diabetes and clinically evident coronary heart disease.  $^{16,18-20}$  Patients with diabetes but without evidence of coronary heart disease may also benefit from cholesterol lowering.  $^{17}$  Because these patients are also at very high risk of CVD events,  $^{55-57}$  the American Diabetes Association recently recommended a low-density lipoprotein cholesterol goal of  $<100~\rm mg/dL$  for all diabetic patients.  $^{32}$ 

Recent data from the U.K. Prospective Diabetes Study clearly demonstrate that aggressive blood pressure control lowers mortality rates in diabetic patients with hypertension.23 However, the best specific drug therapy for treatment of hypertension in diabetes remains a particularly confusing issue. Of the three classes of antihypertensive therapy we considered, ACE inhibitors were prescribed more commonly than thiazide diuretics or  $\beta$ -blockers, despite scant experimental evidence prior to 1998 that hypertension therapy with ACE inhibitors reduces cardiovascular or all-cause morbidity or mortality in either diabetic or nondiabetic patients. Such evidence did exist during our study period for low-dose thiazide diuretic and cardioselective  $\beta$ -blocker-based treatment regimens used as a primary prevention strategy in diabetes, 21,22 and for β-blocker therapy among diabetic patients with diagnosed CVD.<sup>58-62</sup> Both the U.K. Prospective Diabetes Study and the Captopril Prevention Project trial have now demonstrated that captopril and  $\beta$ -blocker-based treatment programs are equally effective for reducing complications and death related to diabetes. 23,25 The ACE inhibitors may have a special role in treatment for type 2 diabetes patients with microalbuminuria and either hypertension or normal blood pressure,  $^{63,64}$  but  $\beta$ -blocker therapy may have a similar beneficial effect.<sup>65</sup> We did not have information on microalbuminuria or diabetic nephropathy and could not stratify the analysis by this important complication. However, about one quarter of people with older-onset diabetes have microalbuminuria, with higher rates observed among insulintreated compared with non-insulin-treated subjects. 66 Paradoxically, we observed lower rates of ACE inhibitor use among hypertensive diabetic patients treated with insulin. However, ACE inhibitor therapy is not without risk,67 and while there has been concern about adverse metabolic or mortality effects of thiazide diuretic or β-blocker use in diabetes, 68-70 subsequent work suggests that these concerns are probably overstated. $^{22,71,72}$  The role of calcium channel blockers for treatment of hypertension in diabetes remains uncertain. Some studies suggest benefit with their use, $^{24,26}$  but others found worrisome adverse outcomes. $^{73,74}$  Current recommendations and expert opinion continue to recommend β-blocker or thiazide-based programs for treatment of hypertension in diabetes. $^{35,75}$  Our analysis demonstrates low overall rates of antihypertensive therapy among hypertensive diabetic patients, and further raises concern about especially low rates of use of β-blockers and thiazide diuretics.

Despite some limitations imposed by the NAMCS design, we have documented that patients with diabetes cared for by U.S. office-based physicians do not appear to have received appropriately aggressive management of CVD risk factors. The burden of morbidity, mortality, and resource use associated with the CVD complications of diabetes make better management of CVD risk factors a critical issue for quality improvement efforts in diabetes care. Among potential strategies, the traditional emphasis on glycemic control must be broadened to include a stronger evidence-based emphasis on CVD prevention. Wider physician education and enhanced information management tools also may help to expand use of interventions to lower blood pressure and lipoprotein cholesterol levels and increase smoking cessation counseling and prophylactic aspirin use-all areas where strong clinical trials results and observational evidence suggest significant reduction in morbidity and mortality for patients with diabetes.

This study was supported by an American Diabetes Association Clinical Research Grant (JBM) and a Mentored Clinical Scientist Development Award from the National Heart, Lung and Blood Institute (K08-HL03548, RSS). Assistance with statistical analyses was provided by Bismruta Misra, MPH.

### **REFERENCES**

- Harris MI, Flegal KM, Cowie CC, et al. Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults. The Third National Health and Nutrition Examination Survey, 1988–1994. Diabetes Care. 1998;21(4):518–24.
- Harris MI. Ambulatory medical care for diabetes. In: National Diabetes Data Group, ed. Diabetes in America: Diabetes Data Compiled 1984. Bethesda, Md: National Institutes of Health; 1985. Publication 85-1468.
- Harris MI. Testing for blood glucose by office-based physicians in the U.S. Diabetes Care. 1990;13:419–26.
- American Diabetes Association. Diabetes 1993 Vital Statistics. Alexandria, Va; American Diabetes Association; 1993.
- 5. UK Prospective Diabetes Study Group. UK Prospective Diabetes Study (UKPDS) 17: a 9-year update of a randomized, controlled trial on the effect of improved metabolic control on complications in non-insulin-dependent diabetes mellitus. Ann Intern Med. 1996;124(1 pt 2):136–45.
- Morrish NJ, Stevens LK, Head J, Fuller JH, Jarrett RJ, Keen H. A
  prospective study of mortality among middle-aged diabetic patients (the London cohort of the WHO Multinational Study of Vascular Disease in Diabetics), II: associated risk factors. Diabetologia. 1990;33:542–8.

- Moss SE, Klein R, Klein BEK. Cause-specific mortality in a population-based study of diabetes. Am J Public Health. 1991; 81:1158–62.
- Rubin RJ, Altman WM, Mendelson DM. Health care expenditures for people with diabetes mellitus, 1992. J Clin Endocrinol Metab. 1994:78:809A-F.
- American Diabetes Association. Economic consequences of diabetes mellitus in the U.S. in 1997. Diabetes Care. 1998;21:296–309.
- Selby JV, Ray GT, Zhang D, Colby CJ. Excess costs of medical care for patients with diabetes in a managed care population. Diabetes Care. 1997;20:1396–401.
- Barrett-Connor E. Does hyperglycemia really cause coronary heart disease? Diabetes Care. 1997;20:1620–3.
- Nathan DM, Meigs JB, Singer DE. The epidemiology of cardiovascular disease in type 2 diabetes mellitus: how sweet it is. or is it? Lancet. 1997;350 (suppl 1):4–9.
- Knatterud GL, Klimt CR, Levin ME, Jacobsen ME, Goldner MG. Effects of hypoglycemic agents on vascular complications in patients with adult-onset diabetes. VII: mortality and selected nonfatal events with insulin treatment. JAMA. 1978:240:37–42.
- 14. Abraira C, Colwell J, Nutall F, et al. Cardiovascular events and correlates in the Veterans Affairs diabetes feasibility trial: VA Cooperative Study on Glycemic Control and Complications in Type II Diabetes. Arch Intern Med. 1997;157:181–8.
- 15. UK Prospective Diabetes Study Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). Lancet. 1998;352:837-53.
- 16. Pyorala K, Pedersen TR, Kjekshus J, Faergeman O, Olsson AG, Thorgeirsson G. Cholesterol lowering with simvastatin improves prognosis of diabetic patients with coronary artery disease: a subgroup analysis of the Scandinavian Simvastatin Survival Study (4S). Diabetes Care. 1997;20:614–20.
- Downs JR, Clearfield M, Weis S, et al. Primary prevention of acute coronary events in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. JAMA. 1998;279:1615–22.
- 18. The Long-term Intervention with Pravastatin in Ischemic Disease (LIPID) Study Group. Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. N Engl J Med. 1998;339:1349–57.
- Goldberg RB, Mellies MJ, Sacks FM, et al. Cardiovascular events and their reduction with pravastatin in diabetic and glucose-intolerant myocardial infarction survivors with average cholesterol levels: subgroup analyses in the Cholesterol and Recurrent Events (CARE) trial. Circulation. 1998;98:2513–9.
- Rubins HB, Robins SJ, Collins D, et al. Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. N Engl J Med. 1999;341:410–8.
- 21. The Hypertension Detection and Follow-up Program Cooperative Research Group. Mortality findings for stepped-care and referredcare participants in the Hypertension Detection and Follow-up Program, stratified by other risk factors. Prev Med. 1985;14:312–35.
- 22. Curb JD, Pressel SL, Cutler JA, et al., for the Systolic Hypertension in the Elderly Program Cooperative Research Group. Effect of diuretic-based antihypertensive treatment on cardiovascular disease risk in older diabetic patients with isolated systolic hypertension. JAMA. 1996;276:1886–92.
- 23. UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. BMJ. 1998;317;703–13.
- 24. Hansson L, Zanchetti A, Carruthers SG, et al. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. Lancet. 1998;351:1755–62.
- 25. Hansson L, Lindholm LH, Niskanen L, et al. Effect of angiotensinconverting enzyme inhibition compared with conventional therapy on cardiovascular morbidity and mortality in hypertension: the

- Captopril Prevention Project (CAPP) randomised trial. Lancet. 1999:353:611–6.
- Tuomilehto J, Rastenyte D, Birkenhager WH, et al. Effects of calcium-channel blockade in older patients with diabetes and systolic hypertension. N Engl J Med. 1999;340:677–84.
- 27. Steering Committee of the Physicians' Health Study Research Group. Final Report on the aspirin component of the ongoing Physicians' Health Study. N Engl J Med. 1989;321:129–35.
- Antiplatelet Trialists' Collaboration. Collaborative overview of randomized trials of antiplatelet therapy I: prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. BMJ. 1994;308:71-2, 81-106.
- Haire-Joshu D, Glasgow RE, Tibbs TL. Smoking and diabetes. Diabetes Care. 1999;22:1887–98.
- American Diabetes Association. Consensus statement: Role of cardiovascular risk factors in prevention and treatment of macrovascular disease in diabetes. Diabetes Care. 1989;12:573–9.
- American Diabetes Association. Treatment of hypertension in diabetes. Diabetes Care. 1996;19 (suppl 1):S107–S113.
- 32. American Diabetes Association. Management of dyslipidemia in adults with diabetes. Diabetes Care. 1998;21 (suppl 1):S36–S39.
- American Diabetes Association. Aspirin therapy in diabetes. Diabetes Care. 1998;21:S45–S46.
- 34. National Cholesterol Education Program. Second Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II). Publication No. 93-3095 ed. Bethesda, MD: National Institutes of Health; 1993.
- Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. Arch Intern Med. 1997;157: 2413–46.
- National Center for Health Statistics. Public Use Data Documentation: 1991 National Ambulatory Medical Care Survey. Washington, DC: National Center for Health Statistics, US Public Health Service; 1992.
- National Center for Health Statistics. Public Use Data Documentation: 1995 National Ambulatory Medical Care Survey. Washington, DC: National Center for Health Statistics, US Public Health Service; 1996.
- National Center for Health Statistics. Public Use Data Documentation: 1996 National Ambulatory Medical Care Survey. Washington, DC: National Center for Health Statistics, US Public Health Service; 1997.
- 39. Potthoff R, Woodbury M, Manton K. Equivalent sample size and equivalent degrees of freedom: refinements for inference using survey weights under superpopulation models. J Am Stat Soc. 1992;87:382–96.
- DeLozier J. The design and methods of the National Ambulatory Medical Care Survey. Med Group Mgmt. 1974;12:5–7.
- Fleiss JL. Statistical Methods for Rates and Proportions. 2nd ed. New York, NY: Wiley and Sons; 1981.
- Hosmer DW, Lemeshow S. Applied Logistic Regression. New York, NY: Wiley-Interscience; 1989.
- 43. Zeger SL, Liang K-Y. Longitudinal data analysis for discrete and continuous outcomes, Biometrics, 1986;42:121–30.
- 44. SAS Institute Inc. SAS/STAT User's Guide, Version 6. 4th ed. Cary, NC: SAS Institute, Inc; 1989.
- National Diabetes Data Group. Diabetes in America. 2nd ed. Bethesda, Md: National Institutes of Health; 1995. NIH publication 95-1468.
- Meigs JB, Nathan DM, Wilson PWF, Cupples LA, Singer DE. Metabolic risk factors worsen continuously across the spectrum of nondiabetic glucose tolerance: the Framingham Offspring Study. Ann Intern Med. 1998;128:524–33.
- American Diabetes Association. Consensus Statement: Detection and Management of Lipid Disorders in Diabetes. Diabetes Care. 1993:16:828–34.

- 48. American Diabetes Association. Consensus Statement: Treatment of Hypertension in Diabetes. Diabetes Care. 1993;16:1394–401.
- 49. Joint Committee on Detection, Evaluation, and Treatment of High Blood Pressure. The fifth report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure (JNC V). Arch Intern Med. 1993;153:154–83.
- Stern MP, Patterson JK, Haffner SM, Hazuda HP, Mitchell BD. Lack of awareness and treatment of hyperlipidemia in type II diabetes in a community survey. JAMA. 1989;262:360–4.
- Hiss RG, Anderson RM, Hess GE, Stepien CJ, Davis WK. Community diabetes care: a 10-year perspective. Diabetes Care. 1994; 17:1124–34.
- Martin TL, Selby JV, Zhang D. Physician and patient prevention practices in NIDDM in a large urban managed-care organization. Diabetes Care. 1995;18:1124–32.
- Peters AL, Legorreta AP, Ossorio RC, Davidson MB. Quality of outpatient care provided to diabetic patients: a health maintenance organization experience. Diabetes Care. 1996;19:601–6.
- 54. Krumholz HM, Radford MJ, Wang Y, Chen J, Heiat A, Marciniak TA. National use and effectiveness of beta-blockers for the treatment of elderly patients after acute myocardial infarction: National Cooperative Cardiovascular Project. JAMA. 1998;280(7):623–9.
- 55. Haffner SM, Stern MP, Hazuda HP, Mitchell BD, Patterson JK. Cardiovascular risk factors in confirmed prediabetic individuals: does the clock for coronary heart disease start ticking before the onset of diabetes? JAMA. 1990;263:2893–8.
- 56. D'Agostino RB Jr, Mykkanen L, Goff DC, et al. Atherosclerosis in type 2 diabetic and non-diabetic subjects with and without previous myocardial infarction: the Insulin Reistance Atherosclerosis Study (IRAS). Circulation 1998;97:817.
- 57. Haffner SM, Lehto S, Ronnemaa T, Pyorala K, Laakso M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. N Engl J Med. 1998;339(4):229–34.
- 58. Gundersen T, Kjekshus J. Timolol treatment after myocardial infarction in diabetic patients. Diabetes Care. 1983;6:285–90.
- Malmberg K, Herlitz J, Hjalmarson A, Ryden L. Effects of metoprolol on mortality and late infarction in diabetics with suspected acute myocardial infarction: retrospective data from two large studies. Eur Heart J. 1989;10:423–8.
- Olsson G, Wikstrand J, Warnold I, et al. Metoprolol-induced reduction in postinfarction mortality: pooled results from five double-blind randomized trials. Eur Heart J. 1992;13:28–32.
- 61. Kendall MJ, Lynch KP, Hjalmarson A, Kjekshus J.  $\beta$ -blockers and sudden cardiac death. Ann Intern Med. 1995;123:358–67.

- Jonas M, Reicher-Reiss H, Boyko V, et al. Usefulness of beta-blocker therapy in patients with non-insulin-dependent diabetes mellitus and coronary artery disease. Am J Cardiol. 1996;77:1273–7.
- 63. Ravid M, Savin H, Jutrin I, Bental T, Katz B, Lishner M. Long term stabilizing effect of angiotensin converting enzyme inhibition on plasma creatinine and on proteinuria in normotensive type II diabetic patients. Ann Intern Med. 1993;118:577–81.
- 64. Kasiske BL, Kalil RSN, Ma JZ, Liao M, Keane WF. Effect of antihypertensive therapy on the kidney in patients with diabetes: a meta-regression analysis. Ann Intern Med. 1993;118:129–38.
- Nielsen FS, Rossing P, Gall M, Skott P, Smidt UM, Parving H. Long-term effect of lisinopril and atenolol on kidney function in hypertensive NIDDM subjects with diabetic nephropathy. Diabetes. 1997;46:1182–8.
- Klein R, Klein BEK, Moss SE. Prevalence of microalbuminuria in older-onset diabetes. Diabetes Care. 1993;16:1325–30.
- 67. Morris AD, Boyle DIR, McMahon AD, et al. ACE inhibitor use is associated with hospitalization for severe hypoglycemia in patients with diabetes. Diabetes Care. 1997;20:1363–7.
- Kaplan NM, Rosenstock J, Raskin P. A differing view of treatment of hypertension in patients with diabetes mellitus. Arch Intern Med. 1987;147:1160–2.
- Lithell HO. Effect of antihypertensive drugs on insulin, glucose, and lipid metabolism. Diabetes Care. 1991;14:203–9.
- Warram JH, Laffel LMB, Valsania P, Christlieb AR, Krolewski AS.
   Excess mortality associated with diuretic therapy in diabetes mellitus. Arch Intern Med. 1991;151:1350–6.
- Giugliano D, Acampora R, Marfella R, et al. Metabolic and cardiovascular effects of carvedilol and atenolol in non-insulin-dependent diabetes mellitus and hypertension. Ann Intern Med. 1997; 126:955–9.
- Lakshman MR, Reda DJ, Materson BJ, Cushman WC, Freis ED. Diuretics and β-blockers do not have adverse effects at 1 year on plasma lipid and lipoprotein profiles in men with hypertension. Arch Intern Med. 1999;159:551–8.
- 73. Estacio RO, Jeffers BW, Hiatt WR, Biggerstaff SL, Gifford N, Schrier RW. The effect of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin-dependent diabetes and hypertension. N Engl J Med. 1998;338:645–52.
- 74. Tatti P, Pahor M, Byington RP, et al. Outcome results of the Fosinopril Versus Amlopidine Cardiovascular Events Randomized Trial (FACET) in patients with hypertension and NIDDM. Diabetes Care. 1998;21:597–603.
- 75. Cutler J. Which drug for treatment of hypertension? Lancet. 1999;353:604–5.