

Linear Regression

Nipun Batra and the teaching staff

IIT Gandhinagar

August 28, 2025

Table of Contents

1. Setup
2. Normal Equation
3. Basis Expansion
4. Geometric Interpretation
5. Dummy Variables and Multicollinearity
6. Practice and Review

Setup

Linear Regression

- Output is continuous in nature.

Linear Regression

- Output is continuous in nature.
- Examples of linear systems:

Linear Regression

- Output is continuous in nature.
- Examples of linear systems:
 - $F = ma$

Linear Regression

- Output is continuous in nature.
- Examples of linear systems:
 - $F = ma$
 - $v = u + at$


Task at hand

- TASK: Predict $\text{Weight} = f(\text{height})$

Height	Weight
3	29
4	35
5	39
2	20
6	41
7	?
8	?
1	?

The first part of the dataset is the training points. The latter ones are testing points.

Scatter Plot



Matrix representation of the expression

- $weight_1 \approx \theta_0 + \theta_1 \cdot height_1$
- $weight_2 \approx \theta_0 + \theta_1 \cdot height_2$
- $weight_N \approx \theta_0 + \theta_1 \cdot height_N$

Matrix representation of the expression

- $weight_1 \approx \theta_0 + \theta_1 \cdot height_1$
- $weight_2 \approx \theta_0 + \theta_1 \cdot height_2$
- $weight_N \approx \theta_0 + \theta_1 \cdot height_N$

$$weight_i \approx \theta_0 + \theta_1 \cdot height_i$$

Matrix representation of the expression

$$\begin{bmatrix} weight_1 \\ weight_2 \\ \dots \\ weight_N \end{bmatrix} = \begin{bmatrix} 1 & height_1 \\ 1 & height_2 \\ \dots & \dots \\ 1 & height_N \end{bmatrix} \begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix}$$

Matrix representation of the expression

$$\begin{bmatrix} weight_1 \\ weight_2 \\ \dots \\ weight_N \end{bmatrix} = \begin{bmatrix} 1 & height_1 \\ 1 & height_2 \\ \dots & \dots \\ 1 & height_N \end{bmatrix} \begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix}$$

$$\hat{\mathbf{y}}_{n \times 1} = \mathbf{X}_{n \times d} \boldsymbol{\theta}_{d \times 1}$$

Matrix representation of the expression

$$\begin{bmatrix} weight_1 \\ weight_2 \\ \dots \\ weight_N \end{bmatrix} = \begin{bmatrix} 1 & height_1 \\ 1 & height_2 \\ \dots & \dots \\ 1 & height_N \end{bmatrix} \begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix}$$

$$\hat{\mathbf{y}}_{n \times 1} = \mathbf{X}_{n \times d} \boldsymbol{\theta}_{d \times 1}$$

- θ_0 - Bias Term/Intercept Term

Matrix representation of the expression

$$\begin{bmatrix} weight_1 \\ weight_2 \\ \dots \\ weight_N \end{bmatrix} = \begin{bmatrix} 1 & height_1 \\ 1 & height_2 \\ \dots & \dots \\ 1 & height_N \end{bmatrix} \begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix}$$

$$\hat{\mathbf{y}}_{n \times 1} = \mathbf{X}_{n \times d} \boldsymbol{\theta}_{d \times 1}$$

- θ_0 - Bias Term/Intercept Term
- θ_1 - Slope

Extension to multiple dimensions

- In the previous example $y = f(x)$, where x is one-dimensional

Extension to multiple dimensions

- In the previous example $y = f(x)$, where x is one-dimensional
- Now consider examples in multiple dimensions

Extension to multiple dimensions

- In the previous example $y = f(x)$, where x is one-dimensional
- Now consider examples in multiple dimensions
- Example: Predict the water demand of the IITGN campus

Extension to multiple dimensions

- In the previous example $y = f(x)$, where x is one-dimensional
- Now consider examples in multiple dimensions
- Example: Predict the water demand of the IITGN campus
- Mathematical representation:

$$\text{Demand} = f(\# \text{ occupants, Temperature})$$

Extension to multiple dimensions

- In the previous example $y = f(x)$, where x is one-dimensional
- Now consider examples in multiple dimensions
- Example: Predict the water demand of the IITGN campus
- Mathematical representation:

$$\text{Demand} = f(\# \text{ occupants, Temperature})$$

- Linear form:

$$\text{Demand} = \text{Base Demand} + K_1 * \# \text{ occupants} + K_2 * \text{Temperature}$$

Intuition

We hope to:

- Learn f : $Demand = f(\#occupants, Temperature)$
- From training dataset
- To predict the condition for the testing set

Linear Relationship

We have

- $x_i = \begin{bmatrix} \textit{Temperature}_i \\ \# \textit{Occupants}_i \end{bmatrix}$

Linear Relationship

We have

- $x_i = \begin{bmatrix} Temperature_i \\ \#Occupants_i \end{bmatrix}$
- Estimated demand for i^{th} sample is
 $\hat{demand}_i = \theta_0 + \theta_1 Temperature_i + \theta_2 Occupants_i$

Linear Relationship

We have

- $x_i = \begin{bmatrix} Temperature_i \\ \#Occupants_i \end{bmatrix}$
- Estimated demand for i^{th} sample is
 $\hat{demand}_i = \theta_0 + \theta_1 Temperature_i + \theta_2 Occupants_i$
- $\hat{demand}_i = x_i'^T \theta$

Linear Relationship

We have

- $x_i = \begin{bmatrix} Temperature_i \\ \#Occupants_i \end{bmatrix}$
- Estimated demand for i^{th} sample is
 $\hat{demand}_i = \theta_0 + \theta_1 Temperature_i + \theta_2 Occupants_i$
- $\hat{demand}_i = x_i'^T \theta$
- where $\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix}$

Linear Relationship

We have

- $x_i = \begin{bmatrix} Temperature_i \\ \#Occupants_i \end{bmatrix}$
- Estimated demand for i^{th} sample is
 $\hat{demand}_i = \theta_0 + \theta_1 Temperature_i + \theta_2 Occupants_i$
- $\hat{demand}_i = x_i'^T \theta$
- where $\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix}$
- and $x_i' = \begin{bmatrix} 1 \\ Temperature_i \\ \#Occupants_i \end{bmatrix} = \begin{bmatrix} 1 \\ x_i \end{bmatrix}$

Linear Relationship

We have

- $x_i = \begin{bmatrix} Temperature_i \\ \#Occupants_i \end{bmatrix}$
- Estimated demand for i^{th} sample is
 $\hat{demand}_i = \theta_0 + \theta_1 Temperature_i + \theta_2 Occupants_i$
- $\hat{demand}_i = x_i'^T \theta$
- where $\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix}$
- and $x_i' = \begin{bmatrix} 1 \\ Temperature_i \\ \#Occupants_i \end{bmatrix} = \begin{bmatrix} 1 \\ x_i \end{bmatrix}$
- Notice the transpose in the equation! This is because x_i is a column vector

We can expect the following

- Demand increases, if # occupants increases, then θ_2 is likely to be positive

We can expect the following

- Demand increases, if # occupants increases, then θ_2 is likely to be positive
- Demand increases, if temperature increases, then θ_1 is likely to be positive

We can expect the following

- Demand increases, if $\#$ occupants increases, then θ_2 is likely to be positive
- Demand increases, if temperature increases, then θ_1 is likely to be positive
- Base demand is independent of the temperature and the $\#$ occupants, but, likely positive, thus θ_0 is likely positive.

Normal Equation

Generalized Linear Regression Format

- Assuming N samples for training

Generalized Linear Regression Format

- Assuming N samples for training
- # Features = M

Generalized Linear Regression Format

- Assuming N samples for training
- # Features = M

Generalized Linear Regression Format

- Assuming N samples for training
- # Features = M

$$\begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_N \end{bmatrix}_{N \times 1} = \begin{bmatrix} 1 & x_{1,1} & x_{1,2} & \dots & x_{1,M} \\ 1 & x_{2,1} & x_{2,2} & \dots & x_{2,M} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 1 & x_{N,1} & x_{N,2} & \dots & x_{N,M} \end{bmatrix}_{N \times (M+1)} \begin{bmatrix} \theta_0 \\ \theta_1 \\ \vdots \\ \theta_M \end{bmatrix}_{(M+1) \times 1}$$

Generalized Linear Regression Format

- Assuming N samples for training
- # Features = M

$$\begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_N \end{bmatrix}_{N \times 1} = \begin{bmatrix} 1 & x_{1,1} & x_{1,2} & \dots & x_{1,M} \\ 1 & x_{2,1} & x_{2,2} & \dots & x_{2,M} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 1 & x_{N,1} & x_{N,2} & \dots & x_{N,M} \end{bmatrix}_{N \times (M+1)} \begin{bmatrix} \theta_0 \\ \theta_1 \\ \vdots \\ \theta_M \end{bmatrix}_{(M+1) \times 1}$$


$$\hat{\mathbf{y}} = \mathbf{X}\boldsymbol{\theta}$$

Relationships between feature and target variables

- There could be different $\theta_0, \theta_1 \dots \theta_M$. Each of them can represents a relationship.
- Given multiples values of $\theta_0, \theta_1 \dots \theta_M$ how to choose which is the best?
- Let us consider an example in 2d


Relationships between feature and target variables

Out of the three fits, which one do we choose?




Relationships between feature and target variables

We have $\hat{y} = 2 + 1x$ as one relationship.



Relationships between feature and target variables

How far is our estimated \hat{y} from ground truth y ?



Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$

Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$
- **Critical Assumption:** ϵ_i are independent and identically distributed (i.i.d.)

Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$
- **Critical Assumption:** ϵ_i are independent and identically distributed (i.i.d.)
- y_i denotes the ground truth for i^{th} sample

Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$
- **Critical Assumption:** ϵ_i are independent and identically distributed (i.i.d.)
- y_i denotes the ground truth for i^{th} sample
- \hat{y}_i denotes the prediction for i^{th} sample, where $\hat{y}_i = \mathbf{x}_i^\top \boldsymbol{\theta}$

Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$
- **Critical Assumption:** ϵ_i are independent and identically distributed (i.i.d.)
- y_i denotes the ground truth for i^{th} sample
- \hat{y}_i denotes the prediction for i^{th} sample, where $\hat{y}_i = \mathbf{x}_i^\top \boldsymbol{\theta}$
- ϵ_i denotes the error/residual for i^{th} sample

Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$
- **Critical Assumption:** ϵ_i are independent and identically distributed (i.i.d.)
- y_i denotes the ground truth for i^{th} sample
- \hat{y}_i denotes the prediction for i^{th} sample, where $\hat{y}_i = \mathbf{x}_i^\top \boldsymbol{\theta}$
- ϵ_i denotes the error/residual for i^{th} sample
- θ_0, θ_1 : The parameters of the linear regression

Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$
- **Critical Assumption:** ϵ_i are independent and identically distributed (i.i.d.)
- y_i denotes the ground truth for i^{th} sample
- \hat{y}_i denotes the prediction for i^{th} sample, where $\hat{y}_i = \mathbf{x}_i^\top \boldsymbol{\theta}$
- ϵ_i denotes the error/residual for i^{th} sample
- θ_0, θ_1 : The parameters of the linear regression
- $\epsilon_i = y_i - \hat{y}_i$

Error terms

- $y_i = \hat{y}_i + \epsilon_i$ where $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$
- **Critical Assumption:** ϵ_i are independent and identically distributed (i.i.d.)
- y_i denotes the ground truth for i^{th} sample
- \hat{y}_i denotes the prediction for i^{th} sample, where $\hat{y}_i = \mathbf{x}_i^\top \boldsymbol{\theta}$
- ϵ_i denotes the error/residual for i^{th} sample
- θ_0, θ_1 : The parameters of the linear regression
- $\epsilon_i = y_i - \hat{y}_i$
- $\epsilon_i = y_i - (\theta_0 + x_i \cdot \theta_1)$

Good fit

- $|\epsilon_1|, |\epsilon_2|, |\epsilon_3|, \dots$ should be small.

Good fit

- $|\epsilon_1|, |\epsilon_2|, |\epsilon_3|, \dots$ should be small.
- minimize $\epsilon_1^2 + \epsilon_2^2 + \dots + \epsilon_N^2$ - L_2 Norm

Good fit

- $|\epsilon_1|, |\epsilon_2|, |\epsilon_3|, \dots$ should be small.
- minimize $\epsilon_1^2 + \epsilon_2^2 + \dots + \epsilon_N^2$ - L_2 Norm
- minimize $|\epsilon_1| + |\epsilon_2| + \dots + |\epsilon_n|$ - L_1 Norm

Normal Equation

- Model specification:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\epsilon}$$

Normal Equation

- Model specification:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\epsilon}$$

- To Learn: $\boldsymbol{\theta}$

Normal Equation

- Model specification:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\epsilon}$$

- To Learn: $\boldsymbol{\theta}$
- Objective: minimize $\epsilon_1^2 + \epsilon_2^2 + \cdots + \epsilon_N^2$

Normal Equation

$$\boldsymbol{\epsilon} = \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_N \end{bmatrix}$$

Normal Equation

$$\boldsymbol{\epsilon} = \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_N \end{bmatrix}$$

Objective: Minimize $\boldsymbol{\epsilon}^\top \boldsymbol{\epsilon}$

Derivation of Normal Equation

$$\boldsymbol{\epsilon} = \mathbf{y} - \mathbf{X}\boldsymbol{\theta}$$

$$\boldsymbol{\epsilon}^\top \boldsymbol{\epsilon} = (\mathbf{y} - \mathbf{X}\boldsymbol{\theta})^\top (\mathbf{y} - \mathbf{X}\boldsymbol{\theta})$$

$$= \mathbf{y}^\top \mathbf{y} - 2\mathbf{y}^\top \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\theta}^\top \mathbf{X}^\top \mathbf{X}\boldsymbol{\theta}$$

This is what we wish to minimize

Minimizing the objective function

$$\frac{\partial \epsilon^\top \epsilon}{\partial \theta} = 0$$

- $\frac{\partial}{\partial \theta} \mathbf{y}^\top \mathbf{y} = 0$
- $\frac{\partial}{\partial \theta} (-2\mathbf{y}^\top \mathbf{X}\theta) = -2\mathbf{X}^\top \mathbf{y}$
- $\frac{\partial}{\partial \theta} (\theta^\top \mathbf{X}^\top \mathbf{X}\theta) = 2\mathbf{X}^\top \mathbf{X}\theta$

Substitute the values in the top equation

Normal Equation derivation

$$\mathbf{0} = -2\mathbf{X}^\top \mathbf{y} + 2\mathbf{X}^\top \mathbf{X} \boldsymbol{\theta}$$

$$\mathbf{X}^\top \mathbf{y} = \mathbf{X}^\top \mathbf{X} \boldsymbol{\theta}$$


$$\hat{\boldsymbol{\theta}}_{OLS} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$$

Worked out example

x	y
0	0
1	1
2	2
3	3

Given the data above, find θ_0 and θ_1 .

Scatter Plot



Worked out example

$$\mathbf{X} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix}$$

$$\mathbf{X}^\top = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \end{bmatrix}$$

$$\mathbf{X}^\top \mathbf{X} = \begin{bmatrix} 4 & 6 \\ 6 & 14 \end{bmatrix}$$

Given the data above, find θ_0 and θ_1 .

Worked out example

$$(\mathbf{X}^\top \mathbf{X})^{-1} = \frac{1}{20} \begin{bmatrix} 14 & -6 \\ -6 & 4 \end{bmatrix}$$


$$\mathbf{X}^\top \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 14 \end{bmatrix}$$

Worked out example

$$\boldsymbol{\theta} = (\mathbf{X}^\top \mathbf{X})^{-1}(\mathbf{X}^\top \mathbf{y})$$

$$\begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix} = \frac{1}{20} \begin{bmatrix} 14 & -6 \\ -6 & 4 \end{bmatrix} \begin{bmatrix} 6 \\ 14 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Scatter Plot




Effect of outlier

x	y
1	1
2	2
3	3
4	0

Compute the θ_0 and θ_1 .

Scatter Plot



Worked out example

$$\mathbf{X} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

$$\mathbf{X}^\top = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix}$$

$$\mathbf{X}^\top \mathbf{X} = \begin{bmatrix} 4 & 10 \\ 10 & 30 \end{bmatrix}$$

Given the data above, find θ_0 and θ_1 .

Worked out example

$$(\mathbf{X}^\top \mathbf{X})^{-1} = \frac{1}{20} \begin{bmatrix} 30 & -10 \\ -10 & 4 \end{bmatrix}$$


$$\mathbf{X}^\top \mathbf{y} = \begin{bmatrix} 6 \\ 14 \end{bmatrix}$$

Worked out example

$$\boldsymbol{\theta} = (\mathbf{X}^\top \mathbf{X})^{-1}(\mathbf{X}^\top \mathbf{y})$$

$$\begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix} = \begin{bmatrix} 2 \\ (-1/5) \end{bmatrix}$$

Scatter Plot



Basis Expansion

Variable Transformation

Transform the data, by including the higher power terms in the feature space.

t	s
0	0
1	6
3	24
4	36

The above table represents the data before transformation

Variable Transformation

Add the higher degree features to the previous table

t	t^2	s
0	0	0
1	1	6
3	9	24
4	16	36

- The above table represents the data after transformation

Variable Transformation

Add the higher degree features to the previous table

t	t^2	s
0	0	0
1	1	6
3	9	24
4	16	36

- The above table represents the data after transformation
- Now, we can write $\hat{s} = f(t, t^2)$

Variable Transformation

Add the higher degree features to the previous table

t	t^2	s
0	0	0
1	1	6
3	9	24
4	16	36

- The above table represents the data after transformation
- Now, we can write $\hat{s} = f(t, t^2)$
- Other transformations: $\log(x)$, $x_1 \times x_2$

A big caveat: Linear in what?!¹

1. $\hat{s} = \theta_0 + \theta_1 * t$ is linear

¹<https://stats.stackexchange.com/questions/8689/what-does-linear-stand-for-in-linear-regression>

A big caveat: Linear in what?!¹

1. $\hat{s} = \theta_0 + \theta_1 * t$ is linear
2. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2$ linear?

¹<https://stats.stackexchange.com/questions/8689/what-does-linear-stand-for-in-linear-regression>

A big caveat: Linear in what?!¹

1. $\hat{s} = \theta_0 + \theta_1 * t$ is linear
2. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2$ linear?
3. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2 + \theta_3 * \cos(t^3)$ linear?

¹<https://stats.stackexchange.com/questions/8689/what-does-linear-stand-for-in-linear-regression>

A big caveat: Linear in what?!¹

1. $\hat{s} = \theta_0 + \theta_1 * t$ is linear
2. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2$ linear?
3. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2 + \theta_3 * \cos(t^3)$ linear?
4. Is $\hat{s} = \theta_0 + \theta_1 * t + e^{\theta_2} * t$ linear?

¹<https://stats.stackexchange.com/questions/8689/what-does-linear-stand-for-in-linear-regression>

A big caveat: Linear in what?!¹

1. $\hat{s} = \theta_0 + \theta_1 * t$ is linear
2. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2$ linear?
3. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2 + \theta_3 * \cos(t^3)$ linear?
4. Is $\hat{s} = \theta_0 + \theta_1 * t + e^{\theta_2} * t$ linear?
5. All except #4 are linear models!

¹<https://stats.stackexchange.com/questions/8689/what-does-linear-stand-for-in-linear-regression>

A big caveat: Linear in what?!¹

1. $\hat{s} = \theta_0 + \theta_1 * t$ is linear
2. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2$ linear?
3. Is $\hat{s} = \theta_0 + \theta_1 * t + \theta_2 * t^2 + \theta_3 * \cos(t^3)$ linear?
4. Is $\hat{s} = \theta_0 + \theta_1 * t + e^{\theta_2} * t$ linear?
5. All except #4 are linear models!
6. Linear refers to the relationship between the parameters that you are estimating (θ) and the outcome

¹<https://stats.stackexchange.com/questions/8689/what-does-linear-stand-for-in-linear-regression>

Basis Functions

- Linear regression only refers to linear in the parameters
- We can perform an arbitrary nonlinear transformation $\phi(x)$ of the inputs x and then linearly combine the components of this transformation.
- $\phi : \mathbb{R}^D \rightarrow \mathbb{R}^K$ is called the basis function

Basis Functions

Some examples of basis functions:

- Polynomial basis: $\phi(x) = \{1, x, x^2, x^3, \dots\}$
- Fourier basis: $\phi(x) = \{1, \sin(x), \cos(x), \sin(2x), \cos(2x), \dots\}$
- Gaussian basis:
$$\phi(x) = \{1, \exp(-\frac{(x-\mu_1)^2}{2\sigma^2}), \exp(-\frac{(x-\mu_2)^2}{2\sigma^2}), \dots\}$$
- Sigmoid basis: $\phi(x) = \{1, \sigma(x - \mu_1), \sigma(x - \mu_2), \dots\}$ where
$$\sigma(x) = \frac{1}{1+e^{-x}}$$

Notebook: [basis.html](#)

Interactive examples and visualizations of different basis functions

Geometric Interpretation

Linear Combination of Vectors

- Let $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \dots, \mathbf{v}_i$ be vectors in \mathbb{R}^D , where D denotes the dimensions

where $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_i \in \mathbb{R}$

Linear Combination of Vectors

- Let $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \dots, \mathbf{v}_i$ be vectors in \mathbb{R}^D , where D denotes the dimensions
- A linear combination of $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \dots, \mathbf{v}_i$ is of the following form:

$$\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \alpha_3 \mathbf{v}_3 + \dots + \alpha_i \mathbf{v}_i$$

where $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_i \in \mathbb{R}$

Span of vectors

- Let v_1, v_2, \dots, v_i be vectors in \mathbb{R}^D , with D dimensions

Span of vectors

- Let v_1, v_2, \dots, v_i be vectors in \mathbb{R}^D , with D dimensions
- The span of v_1, v_2, \dots, v_i is denoted by $\text{SPAN}\{v_1, v_2, \dots, v_i\}$:

$$\{\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_i v_i \mid \alpha_1, \alpha_2, \dots, \alpha_i \in \mathbb{R}\}$$

Span of vectors

- Let v_1, v_2, \dots, v_i be vectors in \mathbb{R}^D , with D dimensions
- The span of v_1, v_2, \dots, v_i is denoted by $\text{SPAN}\{v_1, v_2, \dots, v_i\}$:

$$\{\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_i v_i \mid \alpha_1, \alpha_2, \dots, \alpha_i \in \mathbb{R}\}$$

- It is the set of all vectors that can be generated by linear combinations of v_1, v_2, \dots, v_i

Span of vectors

- Let v_1, v_2, \dots, v_i be vectors in \mathbb{R}^D , with D dimensions
- The span of v_1, v_2, \dots, v_i is denoted by $\text{SPAN}\{v_1, v_2, \dots, v_i\}$:

$$\{\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_i v_i \mid \alpha_1, \alpha_2, \dots, \alpha_i \in \mathbb{R}\}$$


- It is the set of all vectors that can be generated by linear combinations of v_1, v_2, \dots, v_i
- If we stack the vectors v_1, v_2, \dots, v_i as columns of a matrix V , then the span of v_1, v_2, \dots, v_i is given as $V\alpha$ where $\alpha \in \mathbb{R}^i$

Example


Find the span of $\left(\begin{bmatrix} 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}\right)$

Notebook: [geometric-linear-regression.html](#)

Interactive geometric visualization of vector spans and linear regression



Example




We have $v_3 = v_1 + v_2$

We have $v_4 = v_1 - v_2$

Example

Simulating the above example in python using different values of α_1 and α_2




$\text{Span}((v_1, v_2)) \in \mathcal{R}^2$

Example

Find the span of $\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \end{bmatrix} \right)$


- Can we obtain a point (x, y) s.t. $x = 3y$?



Example

Find the span of $\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \end{bmatrix} \right)$


- Can we obtain a point (x, y) s.t. $x = 3y$?
- No



Example

Find the span of $\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \end{bmatrix} \right)$


- Can we obtain a point (x, y) s.t. $x = 3y$?
- No
- Span of the above set is along the line $y = 2x$



Example

Find the span of $\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix} \right)$


- Visualization:



Example

Find the span of $\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix} \right)$

- Visualization:



- The span is the plane $z = x$ or $x_3 = x_1$

Geometric Interpretation

Consider \mathbf{X} and \mathbf{y} as follows.

$$\mathbf{X} = \begin{pmatrix} 1 & 2 \\ 1 & -2 \\ 1 & 2 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} 8.8957 \\ 0.6130 \\ 1.7761 \end{pmatrix}$$

- We are trying to learn $\boldsymbol{\theta}$ for $\hat{\mathbf{y}} = \mathbf{X}\boldsymbol{\theta}$ such that $\|\mathbf{y} - \hat{\mathbf{y}}\|_2$ is minimised

Geometric Interpretation

Consider \mathbf{X} and \mathbf{y} as follows.

$$\mathbf{X} = \begin{pmatrix} 1 & 2 \\ 1 & -2 \\ 1 & 2 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} 8.8957 \\ 0.6130 \\ 1.7761 \end{pmatrix}$$

- We are trying to learn $\boldsymbol{\theta}$ for $\hat{\mathbf{y}} = \mathbf{X}\boldsymbol{\theta}$ such that $\|\mathbf{y} - \hat{\mathbf{y}}\|_2$ is minimised
- Consider the two columns of \mathbf{X} . Can we write $\mathbf{X}\boldsymbol{\theta}$ as the span of $\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix} \right)$?

Geometric Interpretation

Consider \mathbf{X} and \mathbf{y} as follows.


$$\mathbf{X} = \begin{pmatrix} 1 & 2 \\ 1 & -2 \\ 1 & 2 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} 8.8957 \\ 0.6130 \\ 1.7761 \end{pmatrix}$$

- We are trying to learn $\boldsymbol{\theta}$ for $\hat{\mathbf{y}} = \mathbf{X}\boldsymbol{\theta}$ such that $\|\mathbf{y} - \hat{\mathbf{y}}\|_2$ is minimised
- Consider the two columns of \mathbf{X} . Can we write $\mathbf{X}\boldsymbol{\theta}$ as the span of $\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix} \right)$?
- We wish to find $\hat{\mathbf{y}}$ such that


$$\arg \min_{\hat{\mathbf{y}} \in \text{SPAN}\{\bar{x}_1, \bar{x}_2, \dots, \bar{x}_D\}} \|\mathbf{y} - \hat{\mathbf{y}}\|_2$$

Geometric Interpretation

Span of $\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix} \right)$




Geometric Interpretation




- We seek a $\hat{\mathbf{y}}$ in the span of the columns of \mathbf{X} such that it is closest to \mathbf{y}

Geometric Interpretation




- This happens when $y - \hat{y} \perp x_j \forall j$ or $x_j^\top (y - \hat{y}) = 0$

Geometric Interpretation



- This happens when $y - \hat{y} \perp x_j \forall j$ or $x_j^\top (y - \hat{y}) = 0$
- $\mathbf{X}^\top (y - \mathbf{X}\theta) = 0$

Geometric Interpretation



- This happens when $y - \hat{y} \perp x_j \forall j$ or $x_j^\top (y - \hat{y}) = 0$
- $\mathbf{X}^\top (\mathbf{y} - \mathbf{X}\boldsymbol{\theta}) = 0$
- $\mathbf{X}^\top \mathbf{y} = \mathbf{X}^\top \mathbf{X}\boldsymbol{\theta}$ or $\hat{\boldsymbol{\theta}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$

Dummy Variables and Multicollinearity

Multi-collinearity

- There can be situations where inverse of $\mathbf{X}^\top \mathbf{X}$ is not computable

$$\mathbf{X} = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 2 & 4 \\ 1 & 3 & 6 \end{bmatrix} \quad (1)$$

Multi-collinearity

- There can be situations where inverse of $\mathbf{X}^\top \mathbf{X}$ is not computable
- This condition arises when the $|\mathbf{X}^\top \mathbf{X}| = 0$

$$\mathbf{X} = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 2 & 4 \\ 1 & 3 & 6 \end{bmatrix} \quad (1)$$

Multi-collinearity

- There can be situations where inverse of $\mathbf{X}^\top \mathbf{X}$ is not computable
- This condition arises when the $|\mathbf{X}^\top \mathbf{X}| = 0$

$$\mathbf{X} = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 2 & 4 \\ 1 & 3 & 6 \end{bmatrix} \quad (1)$$

- The matrix \mathbf{X} is not full rank

Multi-collinearity

It arises when one or more predictor variables/features in X can be expressed as a linear combination of others

How to tackle it?

- Regularize

Multi-collinearity

It arises when one or more predictor variables/features in X can be expressed as a linear combination of others

How to tackle it?

- Regularize
- Drop variables

Multi-collinearity

It arises when one or more predictor variables/features in X can be expressed as a linear combination of others

How to tackle it?

- Regularize
- Drop variables
- Avoid dummy variable trap

Dummy variables

Say Pollution in Delhi = P

- Model specification:

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_2 * \text{Wind speed} + \theta_3 * \text{Wind Direction}$$

Dummy variables

Say Pollution in Delhi = P

- Model specification:

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_2 * Wind\ speed + \theta_3 * Wind\ Direction$$

- But, wind direction is a categorical variable

Dummy variables

Say Pollution in Delhi = P

- Model specification:

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_2 * Wind\ speed + \theta_3 * Wind\ Direction$$

- But, wind direction is a categorical variable
- It is denoted as follows $\{N:0, E:1, W:2, S:3\}$

Dummy variables

Say Pollution in Delhi = P

- Model specification:

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_2 * Wind\ speed + \theta_3 * Wind\ Direction$$

- But, wind direction is a categorical variable
- It is denoted as follows $\{N:0, E:1, W:2, S:3\}$
- Can we use the direct encoding?

Dummy variables

Say Pollution in Delhi = P

- Model specification:

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_2 * Wind\ speed + \theta_3 * Wind\ Direction$$

- But, wind direction is a categorical variable
- It is denoted as follows $\{N:0, E:1, W:2, S:3\}$
- Can we use the direct encoding?
- This incorrectly implies that $S > W > E > N$ (a meaningless ordering)

Dummy Variables

N-1 Variable encoding

	Is it N?	Is it E?	Is it W?
N	1	0	0
E	0	1	0
W	0	0	1
S	0	0	0

Dummy Variables

N Variable encoding

	Is it N?	Is it E?	Is it W?	Is it S?
N	1	0	0	0
E	0	1	0	0
W	0	0	1	0
S	0	0	0	1

Dummy Variables

Which is better N variable encoding or N-1 variable encoding?

- The N-1 variable encoding is better because the N variable encoding can cause multi-collinearity

Dummy Variables

Which is better N variable encoding or N-1 variable encoding?

- The N-1 variable encoding is better because the N variable encoding can cause multi-collinearity
- Is it $S = 1 - (\text{Is it N} + \text{Is it W} + \text{Is it E})$

Binary Encoding

N	00
E	01
W	10
S	11

- W and S are related by one bit

Binary Encoding

N	00
E	01
W	10
S	11

- W and S are related by one bit
- This introduces dependencies between them, and this can cause confusion in classifiers

Interpreting Dummy variables

Gender	height
F	...
F	...
F	...
M	...
M	...

Interpreting Dummy variables

Gender	height
F	...
F	...
F	...
M	...
M	...

Encoding

Interpreting Dummy variables

Gender	height
F	...
F	...
F	...
M	...
M	...

Encoding

Is Female	height
1	...
1	...
1	...
0	...
0	...

Interpreting Dummy Variables

Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

- Model: $height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$

Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

- Model: $height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$
- We get $\theta_0 = 5.9$ and $\theta_1 = -0.7$

Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

- Model: $height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$
- We get $\theta_0 = 5.9$ and $\theta_1 = -0.7$
- $\theta_0 = \text{Avg height of Male} = 5.9$

Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

- Model: $height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$
- We get $\theta_0 = 5.9$ and $\theta_1 = -0.7$
- $\theta_0 = \text{Avg height of Male} = 5.9$
- $\theta_0 + \theta_1$ is chosen based (equal to) on 5, 5.2, 5.4 (for three records)

Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

- Model: $height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$
- We get $\theta_0 = 5.9$ and $\theta_1 = -0.7$
- $\theta_0 = \text{Avg height of Male} = 5.9$
- $\theta_0 + \theta_1$ is chosen based (equal to) on 5, 5.2, 5.4 (for three records)
- θ_1 is chosen based on 5-5.9, 5.2-5.9, 5.4-5.9

Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

- Model: $height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$
- We get $\theta_0 = 5.9$ and $\theta_1 = -0.7$
- $\theta_0 = \text{Avg height of Male} = 5.9$
- $\theta_0 + \theta_1$ is chosen based (equal to) on 5, 5.2, 5.4 (for three records)
- θ_1 is chosen based on 5-5.9, 5.2-5.9, 5.4-5.9
- $\theta_1 = \text{Avg. female height } (5+5.2+5.4)/3 - \text{Avg. male height}(5.9)$

Interpreting Dummy Variables

Alternatively, instead of a 0/1 coding scheme, we could create a dummy variable

- Alternative encoding: $x_i = \begin{cases} 1 & \text{if } i \text{ th person is female} \\ -1 & \text{if } i \text{ th person is male} \end{cases}$

Interpreting Dummy Variables

Alternatively, instead of a 0/1 coding scheme, we could create a dummy variable

- Alternative encoding: $x_i = \begin{cases} 1 & \text{if } i \text{ th person is female} \\ -1 & \text{if } i \text{ th person is male} \end{cases}$
- Model becomes:
$$y_i = \theta_0 + \theta_1 x_i + \epsilon_i = \begin{cases} \theta_0 + \theta_1 + \epsilon_i & \text{if } i \text{ th person is female} \\ \theta_0 - \theta_1 + \epsilon_i & \text{if } i \text{ th person is male.} \end{cases}$$

Interpreting Dummy Variables

Alternatively, instead of a 0/1 coding scheme, we could create a dummy variable

- Alternative encoding: $x_i = \begin{cases} 1 & \text{if } i \text{ th person is female} \\ -1 & \text{if } i \text{ th person is male} \end{cases}$
- Model becomes:
$$y_i = \theta_0 + \theta_1 x_i + \epsilon_i = \begin{cases} \theta_0 + \theta_1 + \epsilon_i & \text{if } i \text{ th person is female} \\ \theta_0 - \theta_1 + \epsilon_i & \text{if } i \text{ th person is male.} \end{cases}$$
- Interpretation: θ_0 = average person height, θ_1 = amount that female height is above average and male height is below average

Practice and Review

Pop Quiz: Linear Regression

1. What is the geometric interpretation of least squares?

Pop Quiz: Linear Regression

1. What is the geometric interpretation of least squares?
2. When does the normal equation have a unique solution?

Pop Quiz: Linear Regression

1. What is the geometric interpretation of least squares?
2. When does the normal equation have a unique solution?
3. How do polynomial features help with non-linear relationships?

Pop Quiz: Linear Regression

1. What is the geometric interpretation of least squares?
2. When does the normal equation have a unique solution?
3. How do polynomial features help with non-linear relationships?
4. What are the assumptions behind linear regression?

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other
- **Homoscedasticity:** Error variance is constant across all values of x

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other
- **Homoscedasticity:** Error variance is constant across all values of x
- **Normality:** Errors are normally distributed (for inference)

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other
- **Homoscedasticity:** Error variance is constant across all values of x
- **Normality:** Errors are normally distributed (for inference)
- **No Multicollinearity:** Features are not highly correlated

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other
- **Homoscedasticity:** Error variance is constant across all values of x
- **Normality:** Errors are normally distributed (for inference)
- **No Multicollinearity:** Features are not highly correlated

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other
- **Homoscedasticity:** Error variance is constant across all values of x
- **Normality:** Errors are normally distributed (for inference)
- **No Multicollinearity:** Features are not highly correlated

Violation Consequences:

- Biased coefficient estimates

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other
- **Homoscedasticity:** Error variance is constant across all values of x
- **Normality:** Errors are normally distributed (for inference)
- **No Multicollinearity:** Features are not highly correlated

Violation Consequences:

- Biased coefficient estimates
- Invalid confidence intervals

Critical Assumptions of Linear Regression

Before using linear regression, verify these assumptions:

- **Linearity:** Relationship between x and y is linear
- **Independence:** Observations are independent of each other
- **Homoscedasticity:** Error variance is constant across all values of x
- **Normality:** Errors are normally distributed (for inference)
- **No Multicollinearity:** Features are not highly correlated

Violation Consequences:

- Biased coefficient estimates
- Invalid confidence intervals
- Poor prediction performance

Key Takeaways

- **Linear Model:** Assumes linear relationship between features and target

Key Takeaways

- **Linear Model:** Assumes linear relationship between features and target
- **Least Squares:** Minimizes sum of squared residuals

Key Takeaways

- **Linear Model:** Assumes linear relationship between features and target
- **Least Squares:** Minimizes sum of squared residuals
- **Normal Equation:** Closed-form solution when $\mathbf{X}^\top \mathbf{X}$ is invertible

Key Takeaways

- **Linear Model:** Assumes linear relationship between features and target
- **Least Squares:** Minimizes sum of squared residuals
- **Normal Equation:** Closed-form solution when $\mathbf{X}^\top \mathbf{X}$ is invertible
- **Geometric View:** Projection onto column space of design matrix

Key Takeaways

- **Linear Model:** Assumes linear relationship between features and target
- **Least Squares:** Minimizes sum of squared residuals
- **Normal Equation:** Closed-form solution when $\mathbf{X}^\top \mathbf{X}$ is invertible
- **Geometric View:** Projection onto column space of design matrix
- **Feature Engineering:** Basis expansion enables non-linear modeling

Key Takeaways

- **Linear Model:** Assumes linear relationship between features and target
- **Least Squares:** Minimizes sum of squared residuals
- **Normal Equation:** Closed-form solution when $\mathbf{X}^\top \mathbf{X}$ is invertible
- **Geometric View:** Projection onto column space of design matrix
- **Feature Engineering:** Basis expansion enables non-linear modeling
- **Foundation:** Building block for more complex models