

CH365 Chemical Engineering Thermodynamics

Lesson 40
Review and Wrap-up

Lesson 40 Agenda

- Online Canvas surveys
- Paper ABET end-of-semester surveys
- TEE exam review

Term-End Exam

Rooms 331, 341

17 December 0730-1100 Wednesday

500 points, 5 problems, 100 points each

- Apply Raoult's Law and Rachford-Rice method to solve VLE with and without activity.
- Calculate T_{pc} , P_{pc} , and ω_{mix} for a mixture and Z , V_R , H_R , and S_R and ϕ from Lee-Kesler tables.
- Calculate properties of ideal gases and ideal gas mixtures ($C_{p,mix}$, V_{ig} , H_{ig} , and S_{ig}).
- Combine residuals and ideal properties to get real properties.
- Calculate residual properties V_R , S_R , and H_R from cubic EOS (PR, RK, SRK, vdW).

Term-End Exam Review

Slide 6

Calculating Properties

- Pure components – Z, V, H, S, G, and ϕ (or f)

- Ideal gases $Z = \frac{PV}{RT} = 1$ and $\phi = 1$ (ideal gas)

$$H = H_0^{ig} + R \cdot \int_{T_0}^T \left(\frac{C_{P_{298}}^{ig}}{R} \right) dT + H^R \quad (eq 6.50)$$

$$S = S_0^{ig} + R \cdot \left(\int_{T_0}^T \left(\frac{C_{P_{298}}^{ig}}{T} \right) dT - \ln \left(\frac{P}{P_0} \right) \right) + S^R \quad (eq 6.51)$$

- Real gas Z, V, and number of phases from cubic EOS

$$Z = \frac{PV}{RT} \neq 1 \quad Z = 1 + \beta - q\beta \frac{Z - \beta}{(Z + \epsilon\beta)(Z + \sigma\beta)} \quad (eq 3.52, \text{ real gas})$$

- Residual properties from EOS or Lee-Kessler

- Cubic EOS – Lesson 28, 29, Problem 6.141, WPR3

$$\frac{H^R}{RT} = Z - 1 + T_r \left(\frac{dq}{dT_r} \right) \cdot I \quad \frac{S^R}{R} = \ln(Z - \beta) + \left(q + T_r \frac{dq}{dT_r} \right) \cdot I \quad I = \frac{1}{\sigma - \epsilon} \ln \left(\frac{Z + \sigma\beta}{Z + \epsilon\beta} \right) \quad (p. 488) \quad (eq 13.72)$$

- Lee-Kessler tables – Lesson 30

- Real - combine ideal and residuals

- Fugacity – Lesson 35, Slides 10-13, computing - slides 12-13

$$\ln \phi_i = Z_i - 1 - \ln(Z_i - \beta_i) - q_i I_i \quad \beta_i = \Omega \frac{P_{r_i}}{T_r} \quad (eq 3.50) \quad q_i = \frac{\Psi \alpha}{\Omega T_{r_i}} \quad (eq 3.51)$$

- Mixtures – use average ω , T_C , P_C , C_P

- Weighted averages with y_i

- Ideal gas entropy of mixing

$$\text{adds to entropy} \quad R \sum_{i=1}^3 y_i \ln \frac{1}{y_i} \quad (eq 10.26)$$

Term-End Exam Review

- Flash and activity coefficients – modified Raoult's Law

$$y_i P = x_i P_i^{\text{sat}}$$

$$K_i = \frac{y_i}{x_i} = \frac{P_i^{\text{sat}}}{P}$$

Raoult's Law
K-values
(simple solution,
id vapor phase)

$$K_i = \frac{y_i}{x_i} = \frac{\gamma_i P_i^{\text{sat}}}{P}$$

Modified Raoult's Law
K-values
(real liquid phase, id
vapor phase)

$$K_i = \frac{y_i}{x_i} = \frac{\gamma_i f_i^{\text{sat}}}{f}$$

Modified Raoult's Law
K-values with fugacity
(real liquid phase, real
vapor phase)

Calculating Properties

- Rachford-Rice algorithm – Prob. 13.1
- Fugacity and fugacity coefficients
- Practical applications – CHEMCAD, pipeline diameters – Prob. 6.83

Conceptual Issues

- What does it mean? Be able to define:

- | | | |
|--------------------------|--------------------------|------------------------|
| • Simple fluids | • Pitzer correlation | • Fugacity |
| • Acentric factor | • Residual property | • Bip |
| • Enthalpy | • Excess property | • Activity coefficient |
| • Entropy | • Gibbs Energy | • Ideal solution |
| • Gibbs Energy | • Partial molar property | • Ideal gas |
| • Partial molar property | • Chemical potential | • Phase equilibrium |
| • Chemical potential | • Pitzer correlation | • Clapeyron's equation |

Questions?