

CH365 Chemical Engineering Thermodynamics

Lesson 14

Generalized Correlations for Gases and Liquids

Generalized Correlations for Gases

Slide 2

Lee-Kesler Method

Byung Ik Lee and Michael Kesler,
AIChE Journal, 1975, 21(3), 511-527

Pitzer Correlation (L13)

$$Z = Z^0 + \omega Z^1 \quad (\text{Eq. 3.53})$$

simple fluids

deviation from
simple fluids:

$$\omega Z^1 = Z - Z^0$$

$$Z^0 = 1 + B^0 \frac{P_r}{T_r} \quad Z^1 = B^1 \cdot \frac{P_r}{T_r} \quad (\text{Eq. 3.60})$$

Lesson 13, Slide 5
formulas for B^0 and B^1
eqns. 3.61 and 3.62

Lee-Kesler Modification

$Z^{(r)}$: calculated for n-octane

$$Z = Z^0 + \frac{\omega}{\omega^{(r)}} (Z^{(r)} - Z^0) \quad \text{where} \quad Z^1 = \frac{(Z^{(r)} - Z^0)}{\omega^{(r)}}$$

Lee and Kesler used a modified Benedict-Webb-Rubin EOS:

$$Z = 1 + \frac{B}{V_r} + \frac{C}{V_r^2} + \frac{D}{V_r^5} + \frac{c_4}{T_r^3 V_r^2} \left(\beta + \frac{\gamma}{V_r^2} \right) \exp \left(-\frac{\gamma}{V_r^2} \right)$$

B, C, D are functions of T_r
(published in the paper)
 β, γ, c_4 , etc. are constants

Z calculated twice:

Z^0 : calculated for simple fluids

$Z^{(r)}$: calculated for n-octane

Tables: Appendix – Tables D.1-D.4, pp. 676-692

Example: Find Z for n-octane at $P_r=0.4$, $T_r=0.9$

Copyright © McGraw-Hill Education. Permission required for reproduction or display.

Table D.1: Values of Z^0 Page 677

$P_r =$	0.0100	0.0500	0.1000	0.2000	0.4000	0.6000	0.8000	1.0000
T_r								
0.30	0.0029	0.0145	0.0290	0.0579	0.1158	0.1737	0.2315	0.2892
0.35	0.0026	0.0130	0.0261	0.0522	0.1043	0.1564	0.2084	0.2604
0.40	0.0024	0.0119	0.0239	0.0477	0.0953	0.1429	0.1904	0.2379
0.45	0.0022	0.0110	0.0221	0.0442	0.0882	0.1322	0.1762	0.2200
0.50	0.0021	0.0103	0.0207	0.0413	0.0825	0.1236	0.1647	0.2056
0.55	0.9804	0.0098	0.0195	0.0390	0.0778	0.1166	0.1553	0.1939
0.60	0.9849	0.0093	0.0186	0.0371	0.0741	0.1109	0.1476	0.1842
0.65	0.9881	0.9377	0.0178	0.0356	0.0710	0.1063	0.1415	0.1765
0.70	0.9904	0.9504	0.8958	0.0344	0.0687	0.1027	0.1366	0.1703
0.75	0.9922	0.9598	0.9165	0.0336	0.0670	0.1001	0.1330	0.1656
0.80	0.9935	0.9669	0.9319	0.8539	0.0661	0.0985	0.1307	0.1626
0.85	0.9946	0.9725	0.9436	0.8810	0.0661	0.0983	0.1301	0.1614
0.90	0.9954	0.9768	0.9528	0.9015	0.7800	0.1006	0.1321	0.1630
0.93	0.9959	0.9790	0.9573	0.9115	0.8059	0.6635	0.1359	0.1664
0.95	0.9961	0.9803	0.9600	0.9174	0.8206	0.6967	0.1410	0.1705
0.97	0.9963	0.9815	0.9625	0.9227	0.8338	0.7240	0.5580	0.1779
0.98	0.9965	0.9821	0.9637	0.9253	0.8398	0.7360	0.5887	0.1844
0.99								
1.00								
1.01								
1.02								
1.05								
1.10								
1.15								
1.20								
1.30								
1.40								
1.50								
1.60								
1.70								
1.80								
1.90								
2.00								
2.20								
2.40								
2.60								
2.80								
3.00								
3.50								
4.00								

$Z^0=0.780$

$Z^1=-0.1118$

$\omega=0.400$

$$Z = Z^0 + \omega Z^1$$
$$= 0.780 + (0.400) \cdot (-0.1118)$$
$$= .7353$$

Copyright © McGraw-Hill Education. Permission required for reproduction or display.

Table D.2: Values of Z^1 Page 678

$P_r =$	0.0100	0.0500	0.1000	0.2000	0.4000	0.6000	0.8000	1.0000
T_r								
0.30	-0.0008	-0.0040	-0.0081	-0.0161	-0.0323	-0.0484	-0.0645	-0.0806
0.35	-0.0009	-0.0046	-0.0093	-0.0185	-0.0370	-0.0554	-0.0738	-0.0921
0.40	-0.0010	-0.0048	-0.0095	-0.0190	-0.0380	-0.0570	-0.0758	-0.0946
0.45	-0.0009	-0.0047	-0.0094	-0.0187	-0.0374	-0.0560	-0.0745	-0.0929
0.50	-0.0009	-0.0045	-0.0090	-0.0181	-0.0360	-0.0539	-0.0716	-0.0893
0.55	-0.0314	-0.0043	-0.0086	-0.0172	-0.0343	-0.0513	-0.0682	-0.0849
0.60	-0.0205	-0.0041	-0.0082	-0.0164	-0.0326	-0.0487	-0.0646	-0.0803
0.65	-0.0137	-0.0772	-0.0078	-0.0156	-0.0309	-0.0461	-0.0611	-0.0759
0.70	-0.0093	-0.0507	-0.1161	-0.0148	-0.0294	-0.0438	-0.0579	-0.0718
0.75	-0.0064	-0.0339	-0.0744	-0.0143	-0.0282	-0.0417	-0.0550	-0.0681
0.80	-0.0044	-0.0228	-0.0487	-0.1160	-0.0272	-0.0401	-0.0526	-0.0648
0.85	-0.0029	-0.0152	-0.0319	-0.0715	-0.0268	-0.0391	-0.0509	-0.0622
0.90	-0.0019	-0.0099	-0.0205	-0.0442	-0.1118	-0.0396	-0.0503	-0.0604
0.93	-0.0015	-0.0075	-0.0154	-0.0326	-0.0763	-0.1662	-0.0514	-0.0602
0.95	-0.0012	-0.0062	-0.0126	-0.0262	-0.0589	-0.1110	-0.0540	-0.0607
0.97	-0.0010	-0.0050	-0.0101	-0.0208	-0.0450	-0.0770	-0.1647	-0.0623
0.98	-0.0009	-0.0044	-0.0090	-0.0184	-0.0390	-0.0641	-0.1100	-0.0641
0.99	-0.0008	-0.0039	-0.0079	-0.0161	-0.0335	-0.0531	-0.0796	-0.0680
1.00	-0.0007	-0.0034	-0.0069	-0.0140	-0.0285	-0.0435	-0.0588	-0.0879
1.01	-0.0006	-0.0030	-0.0060	-0.0120	-0.0240	-0.0351	-0.0429	-0.0223
1.02	-0.0005	-0.0026	-0.0051	-0.0102	-0.0198	-0.0277	-0.0303	-0.0062
1.05	-0.0003	-0.0015	-0.0029	-0.0054	-0.0092	-0.0097	-0.0032	0.0220
1.10	0.0000	0.0000	0.0001	0.0007	0.0038	0.0106	0.0236	0.0476
1.15	0.0002	0.0011	0.0023	0.0052	0.0127	0.0237	0.0396	0.0625
1.20	0.0004	0.0019	0.0039	0.0084	0.0190	0.0326	0.0499	0.0719
1.30	0.0006	0.0030	0.0061	0.0125	0.0267	0.0429	0.0612	0.0819
1.40	0.0007	0.0036	0.0072	0.0147	0.0306	0.0477	0.0661	0.0857
1.50	0.0008	0.0039	0.0078	0.0158	0.0323	0.0497	0.0677	0.0864
1.60	0.0008	0.0040	0.0080	0.0162	0.0330	0.0501	0.0677	0.0855
1.70	0.0008	0.0040	0.0081	0.0163	0.0329	0.0497	0.0667	0.0838
1.80	0.0008	0.0040	0.0081	0.0162	0.0325	0.0488	0.0652	0.0814
1.90	0.0008	0.0040	0.0079	0.0159	0.0318	0.0477	0.0635	0.0792
2.00	0.0008	0.0039	0.0078	0.0155	0.0310	0.0464	0.0617	0.0767
2.20	0.0007	0.0037	0.0074	0.0147	0.0293	0.0437	0.0579	0.0719
2.40	0.0007	0.0035	0.0070	0.0139	0.0276	0.0411	0.0544	0.0675
2.60	0.0007	0.0033	0.0066	0.0131	0.0260	0.0387	0.0512	0.0634
2.80	0.0006	0.0031	0.0062	0.0124	0.0245	0.0365	0.0483	0.0598
3.00	0.0006	0.0029	0.0059	0.0117	0.0232	0.0345	0.0456	0.0565
3.50	0.0005	0.0026	0.0052	0.0103	0.0204	0.0303	0.0401	0.0497
4.00	0.0005	0.0023	0.0046	0.0091	0.0182	0.0270	0.0357	0.0443

Generalized Correlations for Liquids Slide 4

Rackett: $V^{\text{sat}} = V_C Z_C^{(1-T_r)^{2/7}}$ (Eq. 3.68)

$$Z^{\text{sat}} = \frac{P_r}{T_r} Z_C^{[1+(1-T_r)^{2/7}]} \quad (\text{Eq. 3.69})$$




Figure 3.15: Generalized density correlation for liquids.

Lydersen, Greenkorn, and Hougen: $\rho_r \equiv \frac{\rho}{\rho_C} = \frac{V_C}{V} \quad (\text{Eq. 3.70})$

Example 3.14

For ammonia at 310 K, estimate the molar volume ~~density~~ of (a) the saturated liquid and (b) the liquid at 100 bar.

(Liquids)

Example 3.14 , part b, continued

Rackett: $V^{\text{sat}} = V_C Z_C^{(1-T_r)^{2/7}}$ (Eq. 3.68) $Z^{\text{sat}} = \frac{P_r}{T_r} Z_C^{[1+(1-T_r)^{2/7}]}$ (Eq. 3.69)




Figure 3.15. Generalized density correlation for liquids.

Lydersen, Greenkorn, and Hougen:

$$\rho_r \equiv \frac{\rho}{\rho_C} = \frac{V_C}{V} \quad (\text{Eq. 3.70})$$

$$\rho_r = 2.43$$

solution deviates
from book;
authors read
reduced density
as 2.38 on p. 117

$$T_r = 0.7641$$

$$P_r = 0.8865$$

Example 3.14, continued

For ammonia at 310 K, estimate the molar volume ~~density~~ of (a) the saturated liquid and (b) the liquid at 100 bar.

Questions

Homework

Problem 3.58

To a good approximation, what is the molar volume of ethanol vapor at 480 deg C and 6000 kPa? How does this result compare with the ideal gas?

Answer the problem in four parts:

(a) Lee-Kesler method.

(b) SRK equation.

(c) Ideal gas equation.

For comparison: Compare LK and SRK to IG. If either is less than IG, explain why using knowledge of IG behavior.

Online Interpolator Tool for Lee-Kesler Tables:

https://www.ajdesigner.com/phpinterpolation/bilinear_interpolation_equation.php