

CH365 Chemical Engineering Thermodynamics

Lesson 13 Cubic Equations of State

Corresponding States

Reduced T: $T_r \equiv \frac{T}{T_C}$

Reduced P: $P_r \equiv \frac{P}{P_C}$

All fluids, when compared at the same reduced temperature and reduced pressure, have approximately the same compressibility factor, and all deviate from ideal-gas behavior to about the same degree.

Two-parameter models in terms of T_C and P_C works for simple fluids (Ar, Kr, Xe)

Three-parameter models are greatly improved: use T_C and P_C , and “ ω ”

ω is the “acentric factor”

$$\omega \equiv -1.0 - \log(P_r^{\text{sat}})_{T_r=0.7}$$

Eq. 3.52

Tables in App. B.1

- Introduced by Pitzer and coworkers, 1958
- Related to molecular structure, “centricity”
- Defined with respect to vapor pressure
- Difference between log of reduced vapor pressure of “simple” fluid and more complex fluids at T_r of 0.7

Measure of molecular asymmetry or deviation from molecules.

Used as a property correction.

Example: n-Octane




Figure 3.10: Approximate temperature dependence of the reduced vapor pressure.

3-Parameter Theorem of Corresponding States

Characteristic Properties of Pure Species, Table B.1, pages 663-665

| | Molar Mass | ω | T_c K | P_c bar | Z_c | V_c $\text{cm}^3\text{mol}^{-1}$ | T_n K |
|---------------|------------|----------|------------|--------------|-------|---------------------------------------|------------|
| Methane | 16.043 | 0.012 | 190.6 | 45.99 | 0.286 | 98.6 | 111.4 |
| Ethane | 30.070 | 0.100 | 305.3 | 48.72 | 0.279 | 145.5 | 184.6 |
| Propane | 44.097 | 0.152 | 369.8 | 42.48 | 0.276 | 200.0 | 231.1 |
| n-Butane | 58.123 | 0.200 | 425.1 | 37.96 | 0.274 | 255. | 272.7 |
| ⋮ | ⋮ | ⋮ | ⋮ | ⋮ | ⋮ | ⋮ | ⋮ |
| n-Octane | 114.231 | 0.400 | 568.7 | 24.90 | 0.256 | 486. | 398.8 |
| ⋮ | ⋮ | ⋮ | ⋮ | ⋮ | ⋮ | ⋮ | ⋮ |
| Nitric Acid | 63.013 | 0.714 | 520.0 | 68.90 | 0.231 | 145. | 356.2 |
| Sulfuric Acid | 98.080 | ... | 924.0 | 64.00 | 0.147 | 177.0 | 610.0 |

All fluids having the same value of ω , when compared at the same T_r and P_r , have about the same value of Z , and all deviate from ideal-gas behavior to about the same degree.

Pitzer Correlations

For next lesson; needed for Problem 3.44(b)

2nd Virial Coefficient

$$Z = 1 + \frac{BP}{RT} = 1 + \hat{B} \frac{P_r}{T_r} \quad \text{where} \quad \hat{B} = \frac{BP_c}{RT_c} \quad \begin{matrix} \text{(Eq. 3.57 and 3.36)} \\ \text{(Eq. 3.58)} \end{matrix}$$

Pitzer and coworkers

$$\hat{B} = B^0 + \omega B^1 \quad \text{(Eq. 3.59)}$$

$$B^0 = 0.083 - \frac{0.422}{T_r^{1.6}} \quad \text{(Eq. 3.61)}$$

$$B^1 = 0.139 - \frac{0.172}{T_r^{4.2}} \quad \text{(Eq. 3.62)}$$

3rd Virial Coefficient

$$Z = 1 + B\rho + C\rho^2 = 1 + \hat{B} \frac{P_r}{T_r Z} + \hat{C} \left(\frac{P_r}{T_r Z} \right)^2 \quad \text{where} \quad \hat{C} \equiv \frac{CP_c^2}{R^2 T_c^2} \quad \begin{matrix} \text{(Eq. 3.63 and 3.38)} \\ \text{(Eq. 3.64)} \end{matrix}$$

$$\hat{C} = C^0 + \omega C^1 \quad \text{(Eq. 3.65)}$$

$$C^0 = 0.01407 + \frac{0.02432}{T_r} - \frac{0.00313}{T_r^{10.5}} \quad \text{(Eq. 3.66)}$$

$$C^1 = -0.02676 + \frac{0.05539}{T_r^{2.7}} - \frac{0.00242}{T_r^{10.5}} \quad \text{(Eq. 3.67)}$$

General Cubic Equations of State

$$P = \frac{RT}{V-b} - \frac{a}{(V+\varepsilon b)(V+\sigma b)}$$

(Page 97 Eq. 3.41)

Problem 3.44

$$a = \Psi \frac{\alpha R^2 T_C^2}{P_C} \quad (\text{Eq. 3.45})$$

$$\alpha = \alpha(T, T_C, \omega) \quad \omega \text{ comes in through } \alpha.$$

α , Ψ , ε , σ , and Ω are defined in Table 3.1

$$T_r = \frac{T}{T_C} \quad (\text{reduced temperature})$$

T_C is the critical temperature, Table B.1

P_C is the critical pressure, Table B.1

R is the gas constant, Table A.2

$$b = \Omega \frac{RT_C}{P_C} \quad (\text{Eq. 3.44})$$

van der Waals

$$\alpha = 1, \varepsilon = 0 \text{ and } \sigma = 0, \text{ T3.1}$$

$$\Psi = \frac{27}{64}, \Omega = \frac{1}{8}, \alpha = 1, \text{ T3.1}$$


$$P = \frac{RT}{V-b} - \frac{a}{V^2}$$

$$a = \frac{27 R^2 T_C^2}{64 P_C}$$

$$b = \frac{RT_C}{8P_C}$$

n-Butane Isotherms from Redlich-Kwong EOS

Slide 7



Parameters for Cubic Equations

RK, SRK and PR equations of state were specifically developed for vapor-liquid calculations

| Equation of State | α | σ | ε | Ω | Ψ | Z_c |
|-------------------|-----------------------|----------------|----------------|----------|---------|---------|
| vdW | 1 | 0 | 0 | 0.12500 | 0.42188 | 0.37500 |
| RK | α_{RK} | 1 | 0 | 0.08664 | 0.42748 | 0.33333 |
| SRK | α_{SRK} | 1 | 0 | 0.08664 | 0.42748 | 0.33333 |
| PR | α_{PR} | $1 + \sqrt{2}$ | $1 - \sqrt{2}$ | 0.07780 | 0.45724 | 0.30740 |

Smith, van Ness, Abbott, and Swihart, **Table 3.1, page 100**

$$\alpha_{\text{RK}} = 1/\sqrt{T_r} = T_r^{-1/2}$$

$$\alpha_{\text{SRK}} = \left[1 + (0.480 + 1.574 \omega - 0.176 \omega^2)(1 - \sqrt{T_r}) \right]^2$$

$$\alpha_{\text{PR}} = \left[1 + (0.37464 + 1.54226 \omega - 0.26992 \omega^2)(1 - \sqrt{T_r}) \right]^2$$

vdW – van der Waals – 1873

RK – Redlich-Kwong – 1949

SRK – Soave-Redlich-Kwong – 1972

PR – Peng-Robinson – 1976

Example 3.9

Given that the vapor pressure of n-butane at 350 K is 9.4573 bar, find the molar volumes of (a) saturated vapor and (b) saturated liquid n-butane at these conditions as given by the Redlich-Kwong equation.

Eq. 3.41 gives three roots. Largest is vapor. Smallest is liquid.

(Eq. 3.48)

$$Z = 1 + \beta - q\beta \frac{Z - \beta}{(Z + \varepsilon\beta)(Z + \sigma\beta)}$$

$$\beta = \Omega \frac{P_r}{T_r}$$

(Eq. 3.46 combined with 3.44)


$$q = \frac{\Psi\alpha(T_r, \omega)}{\Omega T_r}$$

(Eq. 3.47 with 3.44)


$$P = \frac{RT}{V - b} - \frac{a}{(V + \varepsilon b)(V + \sigma b)}$$

(Eq. 3.41)

Claimed to be
better for vapor



Claimed to be
better for liquid



$$Z = \beta + (Z + \varepsilon\beta)(Z + \sigma\beta) \frac{1 + \beta - Z}{q\beta}$$

(Eq. 3.49)

Easy to demonstrate with “FindRoot” in Mathematica (numerical; requires initial guess)

Questions

Homework

Problem 3.44

Calculate Z and V for ethylene at 25 deg C and 12 bar by the following equations:

(a) The truncated virial equation (Eq. 3.38) with the following experimental values of virial coefficients:

$$Z = 1 + \frac{B}{V} + \frac{C}{V^2} \quad (3.38)$$

$$B = -140. \text{ cm}^3 \text{ mol}^{-1} \quad \text{and} \quad C = 7,200 \text{ cm}^6 \text{ mol}^{-2}$$

(b) The truncated virial equation (Eq. 3.36), with a value of B from the generalized Pitzer correlation (Eqs. 3.58-3.62)

(c) The Redlich/Kwong equation.

(d) The Soave/Redlich/Kwong equation

(e) The Peng/Robinson equation.

CHEMCAD Thermodynamics

Which EOS is best?

- Selection of the appropriate thermodynamic method is key to producing accurate simulations
- PR is the most widely used thermodynamic package as it applies to all applications involving hydrocarbons
- Special packages should be used when simulation involves non-hydrocarbon components: TEG, amines, sour water, etc.
- Methanol for hydrate prevention has special fit of BIPs in PR equation of state
- In refinery models, review oil characterization before suspecting thermodynamics

Same comments for Aspen/HYSYS