
CH365 Chemical Engineering Thermodynamics

Lesson 27
Fundamental Property Relations
(for Homogeneous Phases)

Motivation

BLUF – Gibbs generating functions are needed for “residual properties”

CHEMCAD Thermodynamic Settings Window



when water is a pure component, CHEMCAD defaults to IAPWS-IF97

Gibbs energy generating function for water:

$$\frac{g(p,T)}{RT} = \gamma(\pi,\tau) = \sum_{i=1}^{34} n_i (7.1-\pi)^{I_i} (\tau - 1.222)^{J_i} \qquad (7)$$

where $\pi = p/p^*$ and $\tau = T^*/T$ with $p^* = 16.53$ MPa and $T^* = 1386$ K; R is given by Eq. (1).

$$R = 0.461\,526 \text{ kJ kg}^{-1}\text{ K}^{-1} \qquad (1)$$

IAPWS R7-97(2012)

The International Association for the Properties of Water and Steam

Lucerne, Switzerland
August 2007

Revised Release on the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam
(The revision only relates to the extension of region 5 to 50 MPa)

©2007 International Association for the Properties of Water and Steam
Publication in whole or in part is allowed in all countries provided that attribution is given to the International Association for the Properties of Water and Steam

Table 3. Relations of thermodynamic properties to the dimensionless Gibbs free energy γ and its derivatives [a] when using Eq. (7)

Property	Relation
Specific volume $v = (\partial g/\partial p)_T$ eq. 6-38	$v(\pi,\tau)\frac{p}{RT} = \pi\gamma_\pi$
Specific internal energy $u = g - T(\partial g/\partial T)_p - p(\partial g/\partial p)_T$	$\frac{u(\pi,\tau)}{RT} = \tau\gamma_\tau - \pi\gamma_\pi$
Specific entropy $s = -(\partial g/\partial T)_p$	$\frac{s(\pi,\tau)}{R} = \tau\gamma_\tau - \gamma$
Specific enthalpy $h = g - T(\partial g/\partial T)_p$ eq. 6-39	$\frac{h(\pi,\tau)}{RT} = \tau\gamma_\tau$

James Clerk Maxwell (1831-1879)



James and Katherine Maxwell
Wikipedia, “James Clerk Maxwell.”



Maxwell’s dog Toby, in Stoddart’s sculpture on George Street in Edinburgh, “Five Statues of Edinburgh Dogs,”

- Albert Einstein described Maxwell's work as the "most profound and the most fruitful that physics has experienced since the time of Newton."
- Maxwell's equations have been called the "second great unification in physics" after Newton’s theory of gravitation. (*A Treatise on Electricity and Magnetism* in 1873.)
- Between 1859 and 1866, developed the theory of velocity distribution of gas molecules, which led to the formulation of the Maxwell–Boltzmann distribution.
- "On Governors,” *Proceedings of the Royal Society*, vol. 16 (1867–1868), first paper on modern control theory.
- In 1871 he published the relations between 2nd derivatives of thermodynamic potentials that we discuss today.

Chapter 5 Look-back

- Entropy from postulates

  - Carnot's Equations
  - Ideal gas PV cycle

L22 slide 5

$$dS^t = \frac{dQ_{rev}}{T} \Leftrightarrow \Delta S^t = \int \frac{dQ_{rev}}{T} \Leftrightarrow S = \frac{Q}{T} \quad \text{(Eq. 5.1)}$$

- Derived entropy change for ideal gas

Combination of 1st law and enthalpy, L23 slide 5

$$\frac{\Delta S}{R} = \int_{T_0}^{T} \frac{C_P^{ig}}{R} \frac{dT}{T} - \ln \frac{P}{P_0} \quad \text{(Eq. 5.10)}$$

- Derived lost work in open systems

Combination of 1st and 2nd laws, L24 slide16

$$\dot{W}_{lost} = T_\sigma \dot{S}_G \quad \text{(Eq. 5.29)}$$

- Developed the concept of "ideal" work

Combination of 1st and 2nd laws, L24 slide14

$$W_{ideal} = \Delta H - T_\sigma \Delta S \quad \text{(Eq. 5.22)}$$

Chapter 6 Look-ahead

- Phase rule – tells us how many intensive variables but not how to calculate them.
- We will develop the fundamental relationships and Maxwell relations - basis of the mathematical structure of modern classical thermodynamics
- We will derive equations that allow calculation of enthalpy and entropy "residuals."
- We will examine "generalized correlations."

Today's Agenda

- Property Relations. (followed closely – red headings)
- Maxwell Relations.
- Enthalpy & Entropy as Functions of T & P.
- Internal Energy as a Function of P.
- Ideal Gas
- Internal Energy & Entropy as Functions of T & V.
- Gibbs Generating Function.

Fundamental Property Relation for U

(Eq. 2.6, 1st Law, closed system, reversible)

$$d(nU) = dQ + dW$$

$$dQ_{rev} = Td(nS) \quad \text{(Eq. 5.1)} \qquad dW_{rev} = -Pd(nV) \quad \text{(Eq. 1.3)}$$

- Not restricted to reversible processes
- **System must be closed and change must be between equilibrium states.**

$$d(nU) = Td(nS) - Pd(nV) \quad \text{(Eq. 6.1)}$$

$$dU = TdS - PdV \quad \text{(Eq. 6.8)}$$

n=1 or constant comp.

- Includes all primary properties: U, T, S, P, and V

Additional properties arise by definition:

Enthalpy: $H \equiv U + PV$ (Eqns. 2.10 and 6.2)

Helmholtz Energy: $A \equiv U - TS$ (Eq. 6.3)

Gibbs Energy: $G \equiv H - TS$ (Eq. 6.4)

Fundamental Property Relation for H

(Eq. 2.6, 1st Law, closed system, reversible)

$$d(nU) = dQ + dW$$

$$dQ_{rev} = Td(nS) \quad \text{(Eq. 5.1)} \qquad dW_{rev} = -Pd(nV) \quad \text{(Eq. 1.3)}$$

$$d(nU) = Td(nS) - Pd(nV) \quad \text{(Eq. 6.1)}$$

Enthalpy: $H \equiv U + PV$ (Eq. 6.2)

$$nH = nU + P(nV)$$

$$d(nH) = d(nU + P(nV)) \Rightarrow d(nH) = d(nU) + Pd(nV) + (nV)dP$$

$$d(nH) = Td(nS) - Pd(nV) + Pd(nV) + (nV)dP$$

$$d(nH) = Td(nS) + (nV)dP \quad \text{(Eq. 6.5)}$$

- Includes H, T, S, P, and V

Distributive Property:

$$\frac{d}{dx}(u+v) = \frac{du}{dx} + \frac{dv}{dx}$$

$$d(u+v) = du + dv$$

Product Rule:

$$\frac{d}{dx}(u \cdot v) = \frac{du}{dx} \cdot v + u \cdot \frac{dv}{dx}$$

$$d(u \cdot v) = du \cdot v + u \cdot dv$$

$dH = TdS + VdP$ (Eq. 6.9)

n=1 or constant comp.

Summary of Fundamental Property Relations

General equations for a homogenous fluid of constant composition.

$dU = TdS - PdV$ ✓ (Eq.6.8)	$dH = TdS + VdP$ ✓ (Eq. 6.9)
$dA = -PdV - SdT$? (Eq. 6.10)	$dG = VdP - SdT$? (Eq. 6.11)

(Problem 6.1 - first homework problem)

(derive 6.10 and 6.11)

Symmetry of 2nd Derivatives

$$f = f(x,y) \quad \Rightarrow \quad df \equiv \left(\frac{\partial f}{\partial x}\right)_y dx + \left(\frac{\partial f}{\partial y}\right)_x dy$$

$$df = M\,dx + N\,dy \qquad \text{(Eq. 6.12)}$$

$$M \equiv \left(\frac{\partial f}{\partial x}\right)_y \qquad N \equiv \left(\frac{\partial f}{\partial y}\right)_x$$

$$\left(\frac{\partial M}{\partial y}\right)_x = \left(\frac{\partial}{\partial y}\left(\frac{\partial f}{\partial x}\right)_y\right)_x = \frac{\partial^2 f}{\partial y \partial x}$$

$$\left(\frac{\partial N}{\partial x}\right)_y = \left(\frac{\partial}{\partial x}\left(\frac{\partial f}{\partial y}\right)_x\right)_y = \frac{\partial^2 f}{\partial x \partial y}$$

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 f}{\partial x \partial y}$$

(order doesn't matter (symmetry of 2nd derivatives); Clairuat's theorem from calculus; a.k.a. Schwarz's theorem, Young's theorem)

$$\left(\frac{\partial M}{\partial y}\right)_x = \left(\frac{\partial N}{\partial x}\right)_y \qquad \text{(Eq. 6.13)}$$

(Basis of Maxwell's equations)

Maxwell's 1st Equation

$$U = U(S,V) \Rightarrow dU = \left(\frac{\partial U}{\partial S}\right)_V dS + \left(\frac{\partial U}{\partial V}\right)_S dV$$

S and V are "special" to U because they appear in the fundamental property relation.

$$dU = T\,dS - P\,dV$$

(fundamental property relation Eq. 6.8, slide 8)

$$T \equiv \left(\frac{\partial U}{\partial S}\right)_V \qquad P \equiv -\left(\frac{\partial U}{\partial V}\right)_S$$

$$\left(\frac{\partial T}{\partial V}\right)_S = \left(\frac{\partial}{\partial V}\left(\frac{\partial U}{\partial S}\right)_V\right)_S = \frac{\partial^2 U}{\partial V \partial S}$$

$$\left(\frac{\partial P}{\partial S}\right)_V = \left(\frac{\partial}{\partial S}\left(\frac{\partial U}{\partial V}\right)_S\right)_V = -\frac{\partial^2 U}{\partial S \partial V}$$

$$\frac{\partial^2 U}{\partial V \partial S} = \frac{\partial^2 U}{\partial S \partial V}$$

$$\therefore \left(\frac{\partial T}{\partial V}\right)_S = -\left(\frac{\partial P}{\partial S}\right)_V \quad \text{(Eq. 6.14)}$$

(First of the four Maxwell equations)

Maxwell's 4th Equation

$$G = G(P,T) \Rightarrow dG = \left(\frac{\partial G}{\partial P}\right)_T dP + \left(\frac{\partial G}{\partial T}\right)_P dT$$

P and T are "special" to G because they appear in the fundamental property relation.

$$dG = V\,dP - S\,dT$$

(fundamental property relation Eq. 6.11, slide 8)

$$V \equiv \left(\frac{\partial G}{\partial P}\right)_T \qquad S \equiv -\left(\frac{\partial G}{\partial T}\right)_P$$

$$\left(\frac{\partial V}{\partial T}\right)_P = \left(\frac{\partial}{\partial T}\left(\frac{\partial G}{\partial P}\right)_T\right)_P = \frac{\partial^2 G}{\partial T \partial P} \qquad \left(\frac{\partial S}{\partial P}\right)_T = -\left(\frac{\partial}{\partial P}\left(\frac{\partial G}{\partial T}\right)_P\right)_T = -\frac{\partial^2 G}{\partial P \partial T}$$

$$\frac{\partial^2 G}{\partial T \partial P} = \frac{\partial^2 G}{\partial P \partial T}$$

$$\therefore \left(\frac{\partial V}{\partial T}\right)_P = -\left(\frac{\partial S}{\partial P}\right)_T \quad \text{(Eq. 6.17)}$$

(Fourth of the four Maxwell equations)

Summary of Maxwell's Equations

These equations are the basis for a large number of equations relating thermodynamic properties.

$\left(\frac{\partial T}{\partial V}\right)_S = \left(\frac{\partial P}{\partial S}\right)_V$ ✓ (Eq. 6.14)	$\left(\frac{\partial T}{\partial P}\right)_S = \left(\frac{\partial V}{\partial S}\right)_P$? (Eq. 6.15)
$\left(\frac{\partial P}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$? (Eq. 6.16)	$\left(\frac{\partial V}{\partial T}\right)_P = -\left(\frac{\partial S}{\partial P}\right)_T$ ✓ (Eq. 6.17)

(Problem 6.4 - second homework problem)

(derive 6.15 and 6.16)

Enthalpy and Entropy as Functions of T and P

The most useful property relations for enthalpy and entropy are functions of T and P

$$H = H(T,P)$$

To write the total differential for this function (H), we need partial derivatives: $\left(\frac{\partial H}{\partial P}\right)_T$ and $\left(\frac{\partial H}{\partial T}\right)_P$

$$dH = \left(\frac{\partial H}{\partial T}\right)_P dT + \left(\frac{\partial H}{\partial P}\right)_T dP$$

$$S = S(T,P)$$

To write the total differential for this function (S), we need partial derivatives: $\left(\frac{\partial S}{\partial P}\right)_T$ and $\left(\frac{\partial S}{\partial T}\right)_P$

$$dS = \left(\frac{\partial S}{\partial T}\right)_P dT + \left(\frac{\partial S}{\partial P}\right)_T dP$$

Enthalpy as a Function of T and P

$$H = H(T,P)$$

$$dH = \left(\frac{\partial H}{\partial T}\right)_P dT + \left(\frac{\partial H}{\partial P}\right)_T dP$$

(Eq. 2.19) $$\left(\frac{\partial H}{\partial T}\right)_P = C_P$$

$$dH = TdS + VdP \quad \text{(Eq. 6.9, slide 8)}$$

$$\frac{dH}{dP} = T\frac{dS}{dP} + V \quad \text{(divide 6.9 by dP)}$$

$$\left(\frac{dH}{dP}\right)_T = T\left(\frac{dS}{dP}\right)_T + V \quad \text{(constant T)}$$

$$\left(\frac{\partial S}{\partial P}\right)_T = -\left(\frac{\partial V}{\partial T}\right)_P$$ (Maxwell Equation) (Eqn. 6.17) (from slide 12)

$$\left(\frac{dH}{dP}\right)_T = V - T\left(\frac{\partial V}{\partial T}\right)_P \quad \text{(Eq. 6.20)}$$

$$dH = C_P dT + \left[V - T\left(\frac{\partial V}{\partial T}\right)_P\right] dP \quad \text{(Eq. 6.21)}$$

(V is a function of T and P)

Entropy as a Function of T and P

$$S = S(T,P)$$

$$dS = \left(\frac{\partial S}{\partial T}\right)_P dT + \left(\frac{\partial S}{\partial P}\right)_T dP$$

(Maxwell Equation)

$$\left(\frac{\partial S}{\partial P}\right)_T = -\left(\frac{\partial V}{\partial T}\right)_P$$ (Eq. 6.17) (slide 12)

$$dH = TdS + VdP$$ (Eq. 6.9) (slide 8)

$$\frac{dH}{dT} = T\frac{dS}{dT} + V\frac{dP}{dT}$$ (divide 6.9 by dT)

$$\left(\frac{\partial H}{\partial T}\right)_P = T\left(\frac{\partial S}{\partial T}\right)_P$$ (constant P)

$$\left(\frac{\partial H}{\partial T}\right)_P = C_P$$ (Eq. 2.20)

$$C_P = T\left(\frac{\partial S}{\partial T}\right)_P$$

$$\frac{C_P}{T} = \left(\frac{\partial S}{\partial T}\right)_P$$ (Eq. 6.18)

(V is a function of T and P)

$$dS = \frac{C_P}{T}dT - \left(\frac{\partial V}{\partial T}\right)_P dP$$ (Eq. 6.22)

(leads directly to formal derivation of eq. 5.10)

Internal Energy as a Function of P

(T constant - implied)

$$U = U(P)$$

$$dU = \left(\frac{\partial U}{\partial P}\right)_T dP$$

$$U = H - PV$$

$$dU = dH - PdV - VdP$$

$$\frac{dU}{dP} = \frac{dH}{dP} - \frac{PdV}{dP} - \frac{VdP}{dP} \qquad \text{(divide dP)}$$

$$\left(\frac{dU}{dP}\right)_T = \left(\frac{dH}{dP}\right)_T - P\left(\frac{dV}{dP}\right)_T - V \qquad \text{(constant T)}$$

$$\left(\frac{dH}{dP}\right)_T = V - T\left(\frac{\partial V}{\partial T}\right)_P \qquad \text{(Eq. 6.20, Slide 14)}$$

$$\left(\frac{dU}{dP}\right)_T = \cancel{V} - T\left(\frac{\partial V}{\partial T}\right)_P - P\left(\frac{dV}{dP}\right)_T - \cancel{V}$$

$$\left(\frac{\partial U}{\partial P}\right)_T = -T\left(\frac{\partial V}{\partial T}\right)_P - P\left(\frac{\partial V}{\partial P}\right)_T \qquad \text{(page 215, not numbered)}$$

$$dU = \left[-T\left(\frac{\partial V}{\partial T}\right)_P - P\left(\frac{\partial V}{\partial P}\right)_T\right] dP \qquad \text{(Equation not given in text)}$$

Ideal Gas State

$$dH = C_P dT + \left[V - T\left(\frac{\partial V}{\partial T}\right)_P \right] dP$$ (Eq. 6.21, slide 14)

$$P \cdot V^{ig} = R \cdot T \Rightarrow V^{ig} = \frac{R \cdot T}{P} \Rightarrow \left(\frac{\partial V^{ig}}{\partial T}\right)_P = \frac{R}{P}$$

$$dH = C_P dT + \left[V - T\frac{R}{P} \right] dP = C_P dT + [V - V] dP = C_P dT$$

$$dH^{ig} = C_P^{ig} dT$$ (Eq. 6.23; Eq. 2.21 p.43, and 3.14a p. 79)

(See L 11, slide 4)

$$dS = \frac{C_P}{T} dT - \left(\frac{\partial V}{\partial T}\right)_P dP$$ (Eq. 6.22, slide 14)

$$dS^{ig} = \frac{C_P^{ig}}{T} dT - \frac{R}{P} dP$$ (Eq. 6.24; leads to Eq. 5.10, p. 188)

(See L 23, slide 9)

(restatements of equations for ideal gases derived in Sections 3.3 and 5.5)

Internal Energy and Entropy as Functions of T and V

Sometimes T and V are more convenient than T and P.
The most useful property relations are for U and S.

$$U = U(T,V)$$

Need: $\left(\frac{dU}{dT}\right)_V$ and $\left(\frac{\partial U}{\partial V}\right)_T$

$$dU = \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

$$S = S(T,V)$$

Need: $\left(\frac{\partial S}{\partial T}\right)_V$ and $\left(\frac{\partial S}{\partial V}\right)_T$

$$dS = \left(\frac{\partial S}{\partial T}\right)_V dT + \left(\frac{\partial S}{\partial V}\right)_T dV$$

Internal Energy and Entropy as Functions of T and V

$$U = U(T,V)$$

$$dU = TdS - PdV \qquad \text{(Eq. 6.8)}$$

$$\frac{dU}{dT} = T\frac{dS}{dT} - P\frac{dV}{dT} \qquad \text{(divide by dT)}$$

$$\left(\frac{dU}{dT}\right)_V = T\left(\frac{dS}{dT}\right)_V \qquad \text{(constant V)}$$

$$C_V \equiv \left(\frac{dU}{dT}\right)_V \qquad \text{(Eq. 2.15)}$$

$$\left(\frac{dS}{dT}\right)_V = \frac{C_V}{T} \qquad \text{(Eq. 6.30)}$$

$$dU = TdS - PdV \qquad \text{(Eq. 6.8)}$$

$$\frac{dU}{dV} = T\frac{dS}{dV} - P\frac{dV}{dV} \qquad \text{(divide by dV)}$$

$$\left(\frac{dU}{dV}\right)_T = T\left(\frac{dS}{dV}\right)_T - P \qquad \text{(constant T)}$$

$$\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial P}{\partial T}\right)_V \qquad \text{(Maxwell Equation)}$$

(Eq. 6.16)

$$\left(\frac{dU}{dV}\right)_T = T\left(\frac{dP}{dT}\right)_V - P \qquad \text{(Eq. 6.31)}$$

Internal Energy and Entropy as Functions of T and V

$$U = U(T,V)$$

$$dU = \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

$$C_V \equiv \left(\frac{dU}{dT}\right)_V \quad \text{(Eq. 2.15)}$$

$$\left(\frac{dU}{dV}\right)_T = T\left(\frac{dP}{dT}\right)_V - P \quad \text{(Eq. 6.31)}$$

$$dU = C_V dT + \left[T \cdot \left(\frac{\partial P}{\partial T}\right)_V - P\right] dV \quad \text{(Eq. 6.32)}$$

$$S = S(T,V)$$

$$dS = \left(\frac{\partial S}{\partial T}\right)_V dT + \left(\frac{\partial S}{\partial V}\right)_T dV$$

$$\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial P}{\partial T}\right)_V \quad \text{(Maxwell Equation)} \quad \text{(Eq. 6.16)}$$

$$\left(\frac{dS}{dT}\right)_V = \frac{C_V}{T} \quad \text{(Eq. 6.30)}$$

$$dS = \frac{C_V}{T} dT + \left(\frac{\partial P}{\partial T}\right)_V dV \quad \text{(Eq. 6.33)}$$

Gibbs Energy Generating Functions I

Equations 6.7-6.10 imply that each of the thermodynamic properties is related to a special pair of variables. These are the *canonical* variables.

$$dG = VdP - SdT \qquad \text{(Eq. 6.11)}$$

$G = G(T,P)$ $\quad A = A(T,V)$ $\quad U = U(S,V)$ $\quad H = H(S,P)$

special

$G \equiv H - TS$

$$d\left(\frac{G}{RT}\right) \equiv \frac{1}{RT}dG - \frac{G}{RT^2}dT$$

$$d\left(\frac{G}{RT}\right) = \frac{VdP - SdT}{RT} - \frac{H - TS}{RT^2}dT$$

$$d\left(\frac{G}{RT}\right) = \frac{V}{RT}dP - \frac{S}{RT}dT - \frac{H}{RT^2}dT + \frac{TS}{RT^2}dT$$

$$d\left(\frac{G}{RT}\right) = \frac{V}{RT}dP - \frac{H}{RT^2}dT \qquad \text{(Eq. 6.37)}$$

Gibbs Energy Generating Functions II

$$d\left(\frac{G}{RT}\right) = \frac{V}{RT}dP - \frac{H}{RT^2}dT \qquad \text{(Eq. 6.37)}$$

$$\frac{G}{RT} = f(T,P) \Rightarrow df \equiv \left(\frac{\partial(G/RT)}{\partial P}\right)_T dP + \left(\frac{\partial(G/RT)}{\partial T}\right)_P dT \qquad \text{(Slide 9)}$$

$$\therefore \frac{V}{RT} = \left(\frac{\partial(G/RT)}{\partial P}\right)_T \quad \text{and} \quad -\frac{H}{RT^2} = \left(\frac{\partial(G/RT)}{\partial T}\right)_P$$

(Eq. 6.38)

Carefully note that both terms V/RT and $-H/RT^2$ are functions of G/RT – Significant Result!

$$\frac{H}{RT} = -T\cdot\left(\frac{\partial(G/RT)}{\partial T}\right)_P$$

(Eq. 6.39)

$$G \equiv H - TS \qquad H \equiv U + PV \qquad A \equiv U - TS$$

$$\frac{S}{R} = \frac{H}{RT} - \frac{G}{RT} \qquad \frac{U}{RT} = \frac{H}{RT} - \frac{PV}{RT} \qquad \frac{A}{RT} = \frac{U}{RT} - \frac{S}{R}$$

Equation 6.11 leads to equations for all thermodynamic properties

Questions?