

Full Photometry Procedure

Abi Chown

Abstract

The procedure outlined in this document takes you through the full photometry pipeline, starting from the calibrated FITS images, all the way through to plotting a light curve for a variable star. The parameters used here are not universal so should not be applied to every situation. They work on Spitzer Data from channels 1 and 2, that is wavelengths of $3.6\mu m$ and $4.5\mu m$. The IRAC (Infrared Array Camera) on board Spitzer has a pixel size of $1.2''$. The “raw” images that are the starting point of the pipeline are Level 2 BCD data so have already been flat-fielded etc.

1 File Setup

The files downloaded from the Spitzer Heritage Archive are given not very useful names, namely the folders are called ‘r’ followed by a string of numbers corresponding to the AOR key. Inside each of these directories are many subdirectories sorting the files into file type and channel they correspond to.

To make the file organisation easier to understand with more intuitive file names, I have organised my files in a hierarchical way as follows: first, the data is sorted into the galaxy the target star belongs to. Then by whether the image is a BCD or PBCD image. Then by the name of the target star. Finally, it is sorted into the correct epoch for which the image was taken. For short, Galaxy \rightarrow BCD/PBCD \rightarrow Target star \rightarrow Epoch. This structure allows one to navigate easily through the data. In order to get my data into this format, I have created the script `file_setup.py`. First, this script creates a list of all the files in the original unsorted directory that need to be sorted. Two functions, `find_bcd_a_home` and `find_pbcd_a_home` then sort all the BCD and PBCD files respectively into their new organisation system. They do this by finding relevant keywords from the FITS headers such as target name, epoch number, galaxy and dither position. These keywords are not only used to sort the file into its correct place, but they also change the name of the file to something more comprehensible, given by `target_channel_epoch_dither_cbcd.fits` for BCD images and `target_channel_epoch_dither_munc.fits` where `munc` can be replaced by `maic` or `mcov`.

Now that the files are in a more logical structure, each file is then converted to data counts using `convert_to_counts.py`. The FITS images are in units of MJy/sr and so Equation 1 is applied, where $flux_{conv}$ and $exptime$ are obtained from the image header.

$$flux_{DN} = \frac{flux_{MJy/sr} \times exptime}{flux_{conv}} \quad (1)$$

The filenames are then amended to reflect this change of unit, namely `_dn` is added between the `cbcd` and the `.fits`.

2 The Star List

An example of the file which provides the pipeline with information on the stars is given here:

GALAXY	STAR	PERIOD	RA	DEC	CHANNEL
LMC	HV00872	29.820	73.7765	-67.4749	1
LMC	HV00878	23.306	74.4624	-69.9583	1
LMC	HV00879	36.840	74.5232	-69.4544	1

The input file needs to be formatted in this way for the pipeline to work.

3 Aperture Photometry

Now that the files are set up in a logical way and we have our list of stars, we can start the actual photometry.

First, we need to perform initial aperture photometry on all the **individual** BCD images. To do this, use `aper_phot.py` with the star list from Section 2 as input (`sysargv[1]`).

The script `aper_phot.py` then uses this text file to perform aperture photometry for all the dithers for each epoch specified in the file. It also copies the `DAOPHOT.opt` and `PHOTO.opt` files to the current working directory. These are necessary for performing photometry with DAOPHOT.

4 Medianed Image & PSF model

5 PSF Photometry

6 ALLFRAME

7 Calibration Steps

8 GLOESS light curves