Supplemental Information

Figure S1: Population forecasts in response to varying removal efforts. Size distributions show the crab abundance in each size class, N_{size} , at the end of the year after overwinter mortality when A. 0 traps, B. 28 traps, C. 112 traps, D. 560 traps, E. 840 traps, F. 1400 traps, and G. 2800 traps were applied evenly over a trapping season of 14 biweeks. Solid line indicates the median size-structured abundance across simulation replicates, and the shaded area indicates ± 1 standard deviation across simulation replicates. Colors indicate trap type used (i.e., in panel B, the purple line shows the resulting size distribution after a trapping effort of 28 Minnow traps).




Figure S2: Population forecasts in response to varying removal efforts, relative to no removal. Size distributions show the ratio of crab abundance in each size class, N_{size} , when no removal occurred relative to a removal effort >0. Ratios are calculated based on size-structured abundance at the end of the year after overwinter mortality when A. 112 traps, B. 560 traps, and C. 2800 traps were applied evenly over a trapping season of 14 biweeks. A ratio of one means that the size-structured abundance after no removal equals the size-structured abundance after application of X traps removes decreases the size-structured abundance, relative to no removal. A ratio greater than one means that the application of X traps increases the size-structured abundance, relative to no removal. Solid line indicates the median size-structured abundance across simulation replicates, and the shaded area indicates ± 1 standard deviation across simulation replicates. Colors indicate trap type used (i.e., in panel B, the purple line shows the resulting size distribution ratio after a trapping effort of 0 traps, relative to 560 Minnow traps).

