

Quantifying the cost of cultural bias in conservation decision making

Abigail G. Keller^{1,*}, Daniel K. Okamoto^{2, a}, Kristin P. Davis^{1, 3, 4, b}, Michael C. Runge^{5, c},
and Carl Boettiger^{1, 4, d}

¹Department of Environment Science, Policy, and Management, University of California, Berkeley, Berkeley, California, USA

²Department of Integrative Biology, University of California, Berkeley, Berkeley, California, USA

³Stone Center for Environmental Stewardship

⁴Eric & Wendy Schmidt Center for Data Science and Environment, University of California, Berkeley

⁵U.S. Geological Survey, Eastern Ecological Science Center at the Patuxent Research Refuge, Laurel, Maryland, USA

*Corresponding author: Abigail G. Keller, agkeller@berkeley.edu, 919-961-5684, 130 Mulford Hall, Berkeley, CA 94720-3114

^adokamoto@berkeley.edu; ^bkpddavis@berkeley.edu; ^cmrunge@usgs.gov; ^dcboettig@berkeley.edu

Key Words

Columbia River Basin; decision theory; ecological management; multi-species functional response; Pacific lamprey; uncertainty quantification; value of information

Article Information

Article type: Letter

Word count, Abstract: 150

Word count, Manuscript: 2977

Number of references: 31

Number of figures: 5

Number of tables: 0

Abstract

Ecological data and the construction of ecological models can reflect cultural bias, dominant systems of knowledge production, and dominant societal values. These human biases limit understanding of the natural world and can impact the effectiveness of conservation decisions. Using methods from decision theory, we demonstrate one way to measure how cultural bias propagates from data to performance on management objectives. We apply this framework in the context of decision making in the Columbia River Basin, where uncertainty about Pacific lamprey – an ecologically-important fish historically overlooked by Euro-American society – impedes decision making. We show that multiple functional forms of the dynamical system are statistically indistinguishable, yet result in very different optimal management actions because of differences that lie outside the range of observed data. By quantifying the cost of uncertainty in units relevant for a decision maker, we demonstrate how legacies of cultural bias in environmental management inhibit conservation decision making.

Introduction

Building an ecological model is an active exercise of bringing about the world (Schlüter et al., 2025). Yet this world-making process is limited, as models often reflect what is known and the biases and perspectives of those who deem information relevant, rather than what is unknown or ignored (Figure 1). The presence or absence of parameters and processes in a model can encode dominant systems of knowledge production, dominant societal values, and cultural bias (Amano & Sutherland, 2013; Silver et al., 2022). The data used to fit these models are also not neutral and often reflect human systems as much as they reflect the natural world. For example, global biodiversity observations reflect macro-economic spatial patterns (Amano & Sutherland, 2013), and human perceptions of species charisma shape citizen science reporting rates (Stoudt et al., 2022). These human biases can generate statistical uncertainty in ecological inference: observing only a subset of environmental space can limit understanding of a full ecological process (Ellis-Soto et al., 2024) (Figure 1). For a conservation decision maker, however, statistical uncertainty is not inherently problematic; understanding the cost of cultural bias requires translation to performance on real-world management objectives (Boettiger, 2022).

Management of Pacific lamprey (*Entosphenus tridentatus*) in the Columbia River Basin (CRB) in the United States provides a useful setting for understanding how cultural bias and dominant societal values propagate from knowledge production to conservation decision making (Box 1). Columbia River Indigenous peoples have long recognized the Pacific lamprey’s cultural and ecological value, yet for most of the last century, Euro-American management and society have placed little value on the Pacific lamprey due to negative perceptions of the species as pests and association with invasive lamprey in the Laurentian Great Lakes (Close et al., 2002). Lamprey abundance in the CRB has precipitously declined over the last century because the species was largely ignored by U.S. federal, state, and land-use management agencies (Close et al., 2002).

Pacific lamprey decline and historic neglect challenge conservation decision making in the Columbia River Basin today. Over the past few decades, the federally endangered, migratory Pacific salmon (*Oncorhynchus* spp.) has faced increased predation pressure by federally-protected California sea lions (*Zalophus californianus*) traveling into the Columbia River (Wargo Rub et al., 2019). This conflict between two protected species has strained decision making around how to support both species, ultimately precipitating an amendment to the Marine Mammal Protection Act to allow legal euthanasia of these sea lions (Marine Mammal Protection Act § 120, 16 U.S.C. § 1389, 2018). The Pacific lamprey could play an important role in buffering this conflict and relieving predation pressure on salmon, as the lamprey has been documented to be a high-value prey item for marine mammals (Columbia River Inter-Tribal Fish Commission, 2011). Yet, the extent


to which Pacific lamprey modulates salmonid survival is uncertain because the legacy of nearly a century of disregard is reflected in data about salmon-lamprey-sea lion interactions (Box 1).

In this study, we apply a decision theoretic lens to quantify the degree to which uncertainty related to the dynamics of an ecologically foundational species - ignored due to cultural bias - impedes decision making in the CRB. We first quantify parametric uncertainty in the multi-species functional response (MSFR), or the functional form describing the rate of sea lion predation as a function of available fish prey (Rosenbaum et al., 2024). This model is developed to consider how systems of knowledge production have generated asymmetries in the observation-generating processes between fish species (Carlen et al., 2024) (Figure 2). We then build a decision model to quantify the value of resolving uncertainty related to the lamprey's role in modulating salmon survival as both a parasite and predation offset for salmon (Figures 1-2). This value of resolving uncertainty - Value of Information - can be considered in the inverse as the cost of a knowledge gap generated as an outcome of cultural bias. Through this Pacific lamprey case study, we quantify one cost of cultural bias in conservation decision making, moving from legacies of neglect reflected in data to performance on a decision objective.

Box 1. Pacific lamprey, Chinook salmon, and California sea lions in the Columbia River Basin

Construction of hydroelectric dams along the Columbia River in the northwestern United States has dramatically altered flow regimes and fish passage, particularly for migratory Pacific salmonids of high cultural and economic value protected under the Endangered Species Act. While millions of dollars have been invested in freshwater habitat restoration and fish ladder construction, in the past few decades these fish have faced a new source of mortality: pinnipeds including California sea lions that have learned to travel to natural pinch points like dams and fish ladders to consume these fish by the thousands. The number of pinnipeds at these predation hotspots is increasing due to recovery efforts related to the Marine Mammal Protection Act (MMPA) and ocean warming events that have shifted the geographic range of marine mammals in the California Current (Wargo Rub et al., 2019). To mitigate the effects of sea lion predation on salmon stocks at risk of extinction, the MMPA was amended in 2018 to authorize states and tribes to lethally remove sea lions, yet these controversial euthanasia practices have strained decision-making processes intended to maximize salmonid returns to the Columbia River Basin (CRB) (Marine Mammal Protection Act § 120, 16 U.S.C.72 § 1389, 2018).

An underacknowledged piece of uncertainty that may impede decisions around policies protecting both salmon and sea lions is the role of Pacific lamprey in modulating salmonid survival and salmonid-sea lion interactions. The Pacific lamprey is an ancient, foundational species in the CRB, alongside which nearly all extant species in the CRB have co-evolved (Close, 2002). The Pacific lamprey serves as a predation buffer for upstream migrating adult salmon from marine mammals, as the species is extraordinarily rich in fats and relatively easy to capture (CRITFC, 2011; Roffe and Matte, 1984). The Pacific lamprey may also decrease salmonid survival in the marine environment, as the species plays a parasitic role in its adult stage, though the impact of parasitic attack wounds on salmonid fitness and survival is not well characterized (Beamish 1980; Weitkamp et al., 2015).



Despite its critical ecological role in the CRB and its high cultural value for Columbia Basin Indigenous peoples (Close et al., 2002; 2004), the Pacific lamprey has faced a precipitous decline in the last century, stemming in part from Euro-American cultural bias and association with invasive lamprey in the Laurentian Great Lakes.

"[L]amprey are considered pests or trash fish by the dominant society, the result of which is the development of eradication programs or at best, a management policy of benign neglect" (Close 2004).

While attention toward Pacific lamprey decline has increased in the last decade — largely due to advocacy, research, and knowledge generated by the Columbia River Inter-Tribal Fish Commission — the legacy of cultural bias against the Pacific lamprey persists in data. Since Pacific lamprey abundance is low relative to historic levels, understanding density-dependence in the

lamprey-salmonid-sea lion functional response is difficult, as only local, rather than global, properties of this functional response are observed (Figure 2; Figure S1). Additionally, most data on lamprey are collected incidental to monitoring of salmonids, where the design and efficiency of the data collection effort is not always adequate for lampreys, resulting in amplified observation error for Pacific lamprey (Kostow, 2002; Tidwell et al., 2023).

Methods

We first describe a stylized representation of a Chinook salmon and Pacific lamprey dynamical system (Box 1). Uncertainty in estuarine predation is quantified using a Bayesian multi-species functional response (MSFR) model with sea lion diet data and fish passage data at Bonneville dam. We then build a decision model to quantify how uncertainty impedes decision making.

Multi-species dynamical system

Here we describe a dynamical system model throughout the lifecycle of Chinook salmon and Pacific lamprey, including predation by California sea lions in the estuary, population growth in the freshwater environment, and mortality in the marine environment. Since California sea lions typically migrate to Bonneville dam in the spring, we focus on the spring run of Chinook salmon (Figure S1). The parameter values used to describe freshwater population growth and marine mortality were guided by historical observations and are described in Table S1. The MSFR parameters were directly estimated with data, and the modeling framework is described in the Supplemental Information.

Estuary predation

The expected number of Pacific lamprey and Chinook salmon consumed by California sea lions is described using a multi-species functional response (MSFR) that allows the attack rate to be dependent on prey density (Rosenbaum et al., 2024). Here, $\tilde{F}_i$ represents the expected number of prey i consumed per predator integrated across one day, a_i represents the density-dependent attack rate of species i , h_k represents the handling time for species $k = 1 \dots m$, and E_i represents the abundance of species i entering the estuary.

$$\tilde{F}_i = \frac{a_i E_i}{1 + \sum_{k=1}^m a_k h_k E_k} \quad (1)$$

The density-dependent attack rate, a_i , is a function of coefficient, b_i , and exponent, q . When $q = 0$, the MSFR follows a Holling type 2 functional response, and when $q < 0$, the MSFR follows a Holling type 3 functional response (Figure S2).

$$a_i = b_i E_i^q \quad (2)$$

The relationship between the abundance of fish returning to freshwater to spawn, R_i , and adults entering the estuary, E_i , of species $i \in \{L, S\}$ is therefore a function of the expected number of prey, $\tilde{F}_i$, and the expected number of sea lion days (i.e., daily abundance in estuary $\times$ days), P :

$$R_i = E_i - \tilde{F}_i \times P \quad (3)$$

The MSFR parameters were estimated with California sea lion gastro-intestinal diet data and fish count data at Bonneville dam (Tables S2-S5). Since the Bonneville dam counting systems were originally designed to count Pacific salmon and not Pacific lamprey returning to freshwater, the Bayesian model formulation was developed to account for asymmetric measurement error between the two fish species (Figure 2; Supporting

Information).

Freshwater population growth

The relationship between 1) the abundance of fish returning to the Columbia River Basin to spawn, R_i and 2) the abundance of juveniles/smolt exiting Bonneville dam downstream, J_i , is described with the Beverton-Holt stock-recruit relationship, where α is a production parameter, and K is the smolt or juvenile carrying capacity. The production parameter, α , combines both freshwater mortality and the population intrinsic rate of growth.

We use the following growth equations to describe the relationship between the abundance of returning fish, R_i , and abundance of juveniles, J_i , for fish species i :

$$J_i = R_i \times \frac{\alpha_i}{(1 + \beta_i R_i)} \quad (4)$$

$$\beta_i = \frac{\alpha_i}{K_i} \quad (5)$$

Marine mortality

In the marine environment, Chinook salmon and Pacific lamprey exhibit a host-parasite relationship (Clemens et al., 2017). The relationship between the abundance of lamprey juveniles entering the ocean, J_L , and the abundance of lamprey adults returning to the estuary, E_L , is described by the following equation, where survival increases as a function of salmon density, J_S , at a rate described by D_L :

$$E_L = J_L \times \phi \times (1 - e^{-D_L \times J_S}) \quad (6)$$

The relationship between the abundance of salmon smolts entering the ocean, J_S , and the abundance of salmon adults returning to the estuary, E_S , is described by the following equation, where survival decreases as a function of lamprey density, J_L , at a rate scaled by D_S :

$$E_S = J_S \times \phi \times (1 - e^{-D_S/J_L}) \quad (7)$$

Here, density-independent ocean survival, ϕ , is shared between both species.

Decision model

We built a decision model to quantify one cost of lamprey-related uncertainty to a decision maker. The Pacific lamprey has both inherent value and spiritual and cultural value to humans, including as a First

Food for Columbia River Basin tribes (Close et al., 2002; Columbia River Inter-Tribal Fish Commission, 2011). However, for the purposes of this decision problem, we primarily considered the value of Pacific lamprey through its ecological role as a predation offset for salmonids to reflect a decision maker whose values align with the dominant value system in the region that prioritizes Pacific salmon. We therefore framed the decision problem as finding the action, a^* , that maximizes utility, U , where utility is defined as the equilibrium abundance of adult Chinook salmon returning to spawn in freshwater, $\hat{R}_S$.

The set of actions, $a \in A$, under consideration in the decision problem include changes in lamprey production, α_L . Since α_L can be understood as the product of adult freshwater survival and intrinsic rate of population growth, an action of increasing α_L could refer to increasing adult lamprey survival (e.g., improving dam passage) and/or increasing lamprey reproduction (e.g., artificial lamprey propagation). While many uncertainties exist in this system, we considered two sources of parametric uncertainty (Figure 2): 1) functional response uncertainty, F (Figure 3) and 2) parasitism uncertainty, P , describing uncertainty in the rate at which salmon ocean survival declines as a function of lamprey abundance.

We calculated the utility of each action, a , using the dynamical system model (Equations 1-7) under all combinations of functional response and parasitism uncertainty, F and P . Here, $U(a, f, p)$ is the utility associated with taking action a assuming $f \in F$ and $p \in P$. The set F corresponds to 1000 randomly selected samples from the MSFR model posterior distribution, and the set P corresponds to three values of D_S (Equation 7) with equal prior probability (Figure 2, Table S6). The utility (i.e., equilibrium salmon return abundance, $\hat{R}_S$) was calculated through simulation, starting at an arbitrary abundance of salmon and lamprey returns, R_S and R_L , and finding the mean salmon return abundance, $\hat{R}_S$, after removing transient dynamics.

Value of Information

We then used a Value of Information analysis to calculate the value of lamprey information. The expected value of perfect information (EVPI), is the difference between the expected value of an optimal action after new information has been collected and the expected value of an optimal action before new information has been collected:

$$\text{EVPI} = E_{F,P}[\max_a U(a, f, p)] - \max_a E_{F,P}[U(a, f, p)] \quad (8)$$

The first term in Equation 8 represents the expected utility once all uncertainty has been resolved, because the optimal action is chosen after perfectly knowing the multi-species functional response relationship and how parasitism strength scales with lamprey density. The second term in Equation 8 is the bet-

hedging strategy, or the expected value associated with the action that maximizes utility over all sources of uncertainty. The difference between these two terms, EVPI, is therefore defined as the expected increase in utility if lamprey uncertainty is resolved.

Results

Quantified multi-species functional response uncertainty

To quantify uncertainty in the California sea lion multi-species functional response (MSFR) (Equations 1-2), we developed a Bayesian model that relates sea lion prey consumption data from gastro-intestinal diet analysis to the observed count of fish passing through Bonneville dam. Across most time points, the abundance of Pacific lamprey was less than 10% of the abundance of Chinook salmon (Figure 3A). Yet despite this low observed abundance, Pacific lamprey were frequently over-represented in the sea lion's diet (Figure 3A). The relatively low abundance of lamprey in the prey community limited observation of global properties of the multi-species functional response, creating high parametric uncertainty in model regions where lamprey make up more than 25% of the salmon-lamprey prey community (Figure 3B-C). This observation of local properties of the density-dependent MSFR prevented distinction between a global preference for lamprey, where lamprey are favored at all densities, and negative switching, where lamprey switch from being over-represented to under-represented in the diet of sea lions as lamprey density increases (Figure 3B-C). These differing functional forms represent realistic density-dependent dynamics (Hellström et al., 2014; Rindorf et al., 2006), yet are statistically indistinguishable due to differences beyond the region of observed data (Figure 3B).

Decision model

The optimal action, a^* , that maximized utility (i.e., equilibrium abundance of adult Chinook salmon returning to the Columbia River Basin) varied across different expressions of functional response and parasitism uncertainty (Figure 4A; Figure S5). The utility associated with the optimal action, $\max_a U(a, f, p)$, also varied across different expressions of functional response and parasitism uncertainty (Figure 4B; Figure S5). In general, expressions of MSFR uncertainty associated with a global preference for lamprey (Figure 3C, blue line) corresponded to higher maximized expected utility, $\max_a E_P[U(a, f, p)]$ (Figure 4A, blue line; Figure S6). Whereas expressions of MSFR uncertainty associated with negative switching (Figure 3C, green line) corresponded to lower maximized expected utility (Figure 4A, green line; Figure S6). Additionally, the bet-hedging strategy, or the action that maximizes utility over all uncertainty ($\max_a E_{F,P}[U(a, f, p)]$), tended to

have a higher expected utility than the action that maximized utility after functional response uncertainty had been resolved ($\max_a E_P[U(a, f, p)]$) (Figure 4A; Figure S6).

Cost of lamprey uncertainty to a decision maker

In the context of this decision problem, the value of lamprey information was quantified as the expected improvement on the management objective (maximize abundance of adult Chinook salmon returning to the Columbia River Basin) if functional response uncertainty, F , and parasitism uncertainty, P , were reduced (Runge et al., 2011). The utility associated with the bet-hedging strategy ($\max_a E_{F,P}[U(a, f, p)]$) was 125,343 fish, whereas the expected utility after all uncertainty had been resolved was 132,511 fish (Figure 4B). The expected value of perfect information, EVPI, was therefore 7167 fish (Figure 4B). Framed another way, the expected increase in Chinook salmon returning to the Columbia River Basin was 7167 fish, if lamprey uncertainty - generated as an outcome of cultural bias - was not impeding the choice of action.

Discussion

Evaluating the outcomes of conservation actions can be limited by model uncertainty that arises from cultural bias. To explore this challenge, we develop a decision model representing the dynamics and management of three interacting species in the Columbia River Basin, where management has historically focused on high-value salmon fisheries and recent predation pressure by sea lions while overlooking a culturally and ecologically important species - the Pacific lamprey. We find that the historical bias against Pacific lamprey by dominant, Euro-American society has led to asymmetric measurement error and observation limited to local properties of the system with lamprey at low relative abundance. As a result of this historical bias, we show that multiple functional forms for the rate of sea lion predation on these two fish species fit the observed data equally well, yet differ in resultant management policies (Runge & Johnson, 2002). Compared to an action designed to perform optimally despite uncertainty, resolving uncertainty related to the rates of California sea lion predation and Pacific lamprey parasitism would increase the expected number of Chinook salmon returning to the Columbia River by almost 6%. While the ecological community broadly acknowledges the ways in which humans and systems of power shape knowledge disparities and assumptions in ecological modeling (Chapman et al., 2024; Silver et al., 2022), our work takes an important step towards quantifying how legacies of cultural bias in environmental management impede today's conservation decisions.

Reducing unknown unknowns

Quantifying the cost of cultural bias to a decision maker through a Value of Information (VOI) analysis is only possible if a spark brings to light previously ignored sources of uncertainty. In the case of the Pacific lamprey, Native American tribes began championing the importance of the species in the early 1990s, recognizing the declining numbers of lamprey and reflecting on the role of dominant Euro-American societal values in contributing to the species' decline (Clemens et al., 2017; Close et al., 2002, 2004). There is, however, not a VOI calculus for uncertainty that is not articulated, reflecting the philosophical conundrum of understanding the cost of “unknown unknowns” in scenarios where the truth lies far outside of the articulated hypotheses (Wintle et al., 2010).

Moving from “unknown unknowns” to quantifying the cost of “known unknowns” can involve interrogating systems of knowledge production through long-established ideas from the fields of environmental justice and science and technology studies. Structural uncertainty in ecological modeling is often not random, as science is not neutral to power and uneven knowledge can arise systematically from political-economic structures and processes (Bloor, 2004; Silver et al., 2022). Here, we have investigated how uneven distribution of power has shaped conservation investment and research in the Columbia River Basin, quantified statistical uncertainty as a result of these intertwined environmental and social processes, and evaluated the impact of this uncertainty on management decisions. Considering knowledge production as a social process in statistical ecology and decision making will be crucial for understanding how legacies of human structures can be present or create absences in ecological model development.

Limitations of an utilitarian understanding of information value

In this analysis, we applied a Western science approach to assess potential impacts of cultural bias in conservation decision making, illustrating this issue by using the currency of the current system of power. We also only focused our analysis on how dominant observation systems can create uncertainties, rather than interrogating the consequences of favoring dominant observation systems at the expense of other perspectives and ways of observing. This approach has both benefits and limitations: while Western scientific data and models can be instruments of domination and exclusion, they can invite contestation due to their legibility and transparency (Peluso, 1995; Scoville et al., 2025).

While we translated signatures of cultural bias reflected in ecological data to a measurement of cost in units relevant to a decision maker (i.e., expected number of salmon), this utilitarian representation reflects a narrow understanding of information value. We chose to evaluate the cost of lamprey uncertainty with respect to one decision objective - maximize the expected annual abundance of adult Chinook salmon returning to

the CRB - among many possible decision objectives. This deliberate choice to frame the value of lamprey through the lens of its ecosystem service for salmon was chosen to represent how an historic, dominant value system that prioritizes Pacific salmon can impede decision making, even if the value system remains unchanged. The cost of cultural bias, however, would change with a different decision objective or a different understanding of cost. Human bias in decision making affects the distribution of costs and benefits of decisions across different groups (Okamoto et al., 2020). Including an objective related to fairness or equity would allow measurement of how uncertainty about Pacific lamprey impedes knowledge of who bears risk as an outcome of a decision (Lockwood et al., 2010). Cultural bias-derived uncertainty can also impede decision making in other ways, including generating conflict and diminishing trust among decision makers, thereby inhibiting the adaptive governance of the social-ecological system (Folke et al., 2005).

Conclusion

By combining uncertainty quantification with decision theory, we have demonstrated one approach to measuring the cost of overlooking the Pacific lamprey in decision making in an important regional fishery. The Bayesian approach to uncertainty quantification offers a flexible framework for representing uneven data-generating processes, and the Value of Information analysis facilitates systematic exploration of this uncertainty in a decision-making context. By unifying these approaches, we quantify the extent to which uncertainty about Pacific lamprey impedes decision making about Chinook salmon, which can be understood as the cost of cultural bias, measured in units relevant to a decision maker.

Data Availability

All data and code for the analysis can be found at https://github.com/abigailkeller/cultural_bias_decisions.

Acknowledgements

Thank you to Marissa Paulling, whose passion for Columbia River fish and advocacy for the Pacific lamprey inspired this work. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, under Award Number DE-SC0024386 and in part upon work supported by the National Science Foundation under Grant No. DBI-1942280.

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the Department of Energy but do represent the views of the US Geological Survey. This draft manuscript is distributed solely for purposes of scientific peer review. Its content is deliberative and predecisional, so it must not be disclosed or released by reviewers. Because the manuscript has not yet been approved for publication by the U.S. Geological Survey (USGS), it does not represent any official USGS finding or policy.

References

- Amano, T., & Sutherland, W. J. (2013). Four barriers to the global understanding of biodiversity conservation: Wealth, language, geographical location and security. *Proceedings of the Royal Society B: Biological Sciences*, 280(1756), 20122649.
- Beamish, R. J. (1980). Adult biology of the river lamprey (*Lampetra ayresi*) and the Pacific lamprey (*Lampetra tridentata*) from the Pacific coast of Canada. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(11), 1906–1923.
- Bloor, D. (2004). Sociology of scientific knowledge. In *Handbook of epistemology* (pp. 919–962). Springer.
- Boettiger, C. (2022). The forecast trap. *Ecology Letters*, 25(7), 1655–1664.
- Carlen, E. J., Estien, C. O., Caspi, T., Perkins, D., Goldstein, B. R., Kreling, S. E., Hentati, Y., Williams, T. D., Stanton, L. A., Des Roches, S., et al. (2024). A framework for contextualizing social-ecological biases in contributory science data. *People and Nature*, 6(2), 377–390.
- Chapman, M., Goldstein, B. R., Schell, C. J., Brashares, J. S., Carter, N. H., Ellis-Soto, D., Faxon, H. O., Goldstein, J. E., Halpern, B. S., Longdon, J., et al. (2024). Biodiversity monitoring for a just planetary future. *Science*, 383(6678), 34–36.

- Clemens, B. J., Beamish, R. J., Coates, K. C., Docker, M. F., Dunham, J. B., Gray, A. E., Hess, J. E., Jolley, J. C., Lampman, R. T., McIlraith, B. J., et al. (2017). Conservation challenges and research needs for Pacific lamprey in the Columbia river basin. *Fisheries*, 42(5), 268–280.
- Close, D. A., Fitzpatrick, M. S., & Li, H. W. (2002). The ecological and cultural importance of a species at risk of extinction, Pacific lamprey. *Fisheries*, 27(7), 19–25.
- Close, D. A., Jackson, A., Conner, B., & Li, H. (2004). Traditional ecological knowledge of Pacific lamprey (*Entosphenus tridentatus*) in northeastern Oregon and southeastern Washington from indigenous peoples of the Confederated Tribes of the Umatilla Indian Reservation. *Journal of Northwest Anthropology*, 38(2), 141–162.
- Columbia River Inter-Tribal Fish Commission. (2011). *Tribal Pacific Lamprey Restoration Plan for the Columbia River Basin* (tech. rep.).
- Ellis-Soto, D., Chapman, M., & Koltz, A. M. (2024). Addressing data disparities is critical for biodiversity assessments. *Trends in Ecology & Evolution*.
- Folke, C., Hahn, T., Olsson, P., & Norberg, J. (2005). Adaptive governance of social-ecological systems. *Annu. Rev. Environ. Resour.*, 30(1), 441–473.
- Hellström, P., Nyström, J., & Angerbjörn, A. (2014). Functional responses of the rough-legged buzzard in a multi-prey system. *Oecologia*, 174(4), 1241–1254.
- Kostow, K. (2002). *Oregon lampreys: Natural history, status, and problem analysis*. Oregon Department of Fish; Wildlife [Fish Division].
- Lockwood, M., Davidson, J., Curtis, A., Stratford, E., & Griffith, R. (2010). Governance principles for natural resource management. *Society and natural resources*, 23(10), 986–1001.
- Marine Mammal Protection Act § 120, 16 U.S.C. § 1389. (2018).
- Okamoto, D. K., Poe, M. R., Francis, T. B., Punt, A. E., Levin, P. S., Shelton, A. O., Armitage, D. R., Cleary, J. S., Dressell, S. C., Jones, R., et al. (2020). Attending to spatial social–ecological sensitivities to improve trade-off analysis in natural resource management. *Fish and Fisheries*, 21(1), 1–12.
- Peluso, N. L. (1995). Whose woods are these? Counter-mapping forest territories in Kalimantan, Indonesia. *Antipode*, 27(4), 383–406.
- Rindorf, A., Gislason, H., & Lewy, P. (2006). Prey switching of cod and whiting in the North sea. *Marine Ecology Progress Series*, 325, 243–253.
- Roffe, T. J., & Mate, B. R. (1984). Abundances and feeding habits of pinnipeds in the Rogue River, Oregon. *The Journal of Wildlife Management*, 1262–1274.

- Rosenbaum, B., Li, J., Hirt, M. R., Ryser, R., & Brose, U. (2024). Towards understanding interactions in a complex world: Design and analysis of multi-species functional response experiments. *Methods in Ecology and Evolution*, 15(9), 1704–1719.
- Runge, M. C., Converse, S. J., & Lyons, J. E. (2011). Which uncertainty? Using expert elicitation and expected value of information to design an adaptive program. *Biological Conservation*, 144(4), 1214–1223.
- Runge, M. C., & Johnson, F. A. (2002). The importance of functional form in optimal control solutions of problems in population dynamics. *Ecology*, 83(5), 1357–1371.
- Schlüter, M., Hertz, T., Klein, A., & Wijermans, N. (2025). Disentangling the entangled in productive ways: Modelling social–ecological systems from a process-relational perspective. *Sustainability Science*, 20(3), 793–815.
- Scoville, C., Amironesei, R., Xu, L., Chapman, M., Record, N. R., & Boettiger, C. (2025). From maps to models: Participation and contestability in the dynamic management of natural resources. *Geo: Geography and Environment*, 12(2), e70028.
- Silver, J. J., Okamoto, D. K., Armitage, D., Alexander, S. M., Atleo, C., Burt, J. M., Jones, R., Lee, L. C., Muhl, E.-K., Salomon, A. K., et al. (2022). Fish, people, and systems of power: Understanding and disrupting feedback between colonialism and fisheries science. *The American Naturalist*, 200(1), 168–180.
- Stoudt, S., Goldstein, B. R., & de Valpine, P. (2022). Identifying engaging bird species and traits with community science observations. *Proceedings of the National Academy of Sciences*, 119(16), e2110156119.
- Tidwell, K. S., Braun, M. W., & van der Leeuw, B. K. (2023). Evaluation of Pinniped Predation on Adult Salmonids and Other Fish in the Bonneville Dam Tailrace, 2022 (Technical Report). U.S. Army Corps of Engineers, Portland District, Fisheries Field Unit. Cascade Locks, OR.
- Wargo Rub, A. M., Som, N. A., Henderson, M. J., Sandford, B. P., Van Doornik, D. M., Teel, D. J., Tennis, M. J., Langness, O. P., van der Leeuw, B. K., & Huff, D. D. (2019). Changes in adult Chinook salmon (*Oncorhynchus tshawytscha*) survival within the lower Columbia River amid increasing pinniped abundance. *Canadian Journal of Fisheries and Aquatic Sciences*, 76(10), 1862–1873.
- Weitkamp, L. A., Hinton, S. A., & Bentley, P. J. (2015). Seasonal abundance, size, and host selection of Western River (*Lampetra ayresii*) and Pacific (*Entosphenus tridentatus*) lampreys in the Columbia River estuary. *Fishery Bulletin*, 113(2), 213–226.
- Wintle, B. A., Runge, M. C., & Bekessy, S. A. (2010). Allocating monitoring effort in the face of unknown unknowns. *Ecology letters*, 13(11), 1325–1337.

Figures




Figure 1: What is the cost of looking only under a lamp-post for a scientific solution, simply because that's where the light is? This focus as result of human bias can lead to unsampled, data-poor regions of environmental space, generating uncertainty in ecological processes that can propagate to uncertainty in the decision space. Here we present a methodological framework combining uncertainty quantification and decision theory that can be used to quantify the cost of cultural bias to a decision maker.




Figure 2: Two sources of parametric uncertainty considered in the decision problem: **A.** Functional response uncertainty, including the density-dependent rates of predation and prey switching, derived from 1) uneven observation error of the true abundance of fish passing through the dam (S^P and L^P for salmon and lamprey, respectively) and 2) observation of local properties of the density-dependent multi-species functional response (i.e., observation space limited to low Pacific lamprey estuarine abundance, E_L). **B.** Parasitism uncertainty, or an unknown rate at which salmon ocean survival (proportion of juvenile salmon returning to the estuary, E_S/J_S) declines with juvenile lamprey abundance (J_L). The parameter D_S describes this rate, and values are selected to cover the universe of possibilities, from a commensal lamprey-salmon marine relationship ($D_S = 9e7$), to a scenario where salmon ocean survival declines to 0.04 at high lamprey densities ($D_S = 3e6$).




Figure 3: Multi-species functional response, measuring the rate at which California sea lions consume prey as a function of the composition of the prey community. **A.** Relationship between the proportion of each fish species in the predator diet, measured with gastro-intestinal analysis of euthanized California sea lions, and proportion of each fish species in the prey community, based on the mean posterior estimate of available prey, L_A and S_A for salmon and lamprey, respectively. The credibility intervals for the estimates of available prey, L_A and S_A , are shown in Figure S4. **B.** Conceptual multi-species functional responses demonstrating how two multi-species functional response curves can have similar local properties in the observed parametric region shown in panel A, yet different global properties due to a large, unobserved parametric region. **C.** Estimated multi-species functional response, where each black line represents a posterior sample describing the predicted relationship between the expected proportion of each species in the sea lion diet and the proportion of each species in the prey community. Four posterior samples are highlighted in color and are used in Figure 4. The dashed line indicates the 1:1 line, where sea lions consume prey in equal proportion to their abundance in the prey community.




Figure 4: Quantifying the value of reducing lamprey-related uncertainty. **A.** Relationship between the expected utility and action for each multi-species functional response (MSFR) posterior sample highlighted in Figure 3C. Asterisk indicates that the expected utility, $E_P[U(a, f, p)]$, is calculated as the expectation over parasitism uncertainty, P (Figure 2). The actions are the lamprey production, α_L , relative to salmon production, α_S , and can be understood as actions that either increase freshwater lamprey adult survival or the intrinsic rate of lamprey population growth. The points indicate the action that maximizes expected utility (i.e., $\max_a E_P[U(a, f, p)]$) for each expression of functional response uncertainty. The black line indicates the expected utility over both parasitism and functional response uncertainty (i.e., $E_{F,P}[U(a, f, p)]$), and the black point indicates the action that maximizes utility over all uncertainty, or the bet-hedging strategy (i.e., $\max_a E_{F,P}[U(a, f, p)]$). **B.** Violin plot of the maximized expected utility of all posterior samples, calculated as the expectation across parasitism uncertainty (i.e., $\max_a E_P[U(a, f, p)]$). Colored points indicate the maximized expected utility of the selected posterior samples from panel A. The black point indicates the maximized expected utility across both parasitism and functional response uncertainty, or the utility associated with the bet-hedging strategy, $\max_a E_{F,P}[U(a, f, p)]$. The purple dashed line indicates the expected utility after all uncertainty has been resolved, $E_{F,P}[\max_a U(a, f, p)]$. The expected value of functional response and parasitism information (i.e., expected value of perfect information, EVPI) is therefore the difference between $E_{F,P}[\max_a U(a, f, p)]$ and $\max_a E_{F,P}[U(a, f, p)]$. Purple shading signifies reduction of uncertainty.