



JOHNS HOPKINS
BLOOMBERG SCHOOL
of PUBLIC HEALTH

Lecture 5

The classical linear regression model


The material in this video is subject to the copyright of the owners of the material and is being provided for educational purposes under rules of fair use for registered students in this course only. No additional copies of the copyrighted work may be made or distributed.

Review of key concepts from Lecture 3 and 4



► Simple linear regression model

► $ARM = B0 + B1 (age - 6) + e$, $e \sim N(0, \sigma^2)$, independent



systematic component \rightarrow random component / error

$B0$ = average ARM at age = 6 months


$B1$ = expected difference in ARM comparing two children who differ in age by 1 month



Review of key concepts from Lecture 3 and 4

- Sex adjusted relationship between ARM and age

- $ARM = B_0 + B_1 (age - 6) + B_2 \text{ Female} + e, e \sim N(0, \sigma^2), \text{ independent}$



$$B_0 + B_1 (age - 6) + B_2$$

$$(B_0 + B_2) + B_1 (age - 6)$$


B_0 = Average ARM among male children 6 months of age

B_1 = difference in mean ARM comparing children of the same gender but whom differ by 1 month of age

B_2 = Difference in average ARM comparing female to males at any Age

Review of key concepts from Lecture 3 and 4

- ▶ Height adjusted relationship between ARM and age
- ▶ $ARM = B_0 + B_1 (age - 6) + B_2 (HT - 62) + e$, $e \sim N(0, \sigma^2)$, independent




B_1 = difference in the mean ARM comparing children who differ in age by 1 month but whom have the same HT.

Review of key concepts from Lecture 3 and 4

- Effect modification: Is the ARM vs. age relationship the same or different by sex

- $ARM = B_0 + B_1 (age - 6) + B_2 \text{ Female} + B_3 (age - 6) \text{ Female} + e, e \sim N(0, \sigma^2)$, independent



$B_3 =$ difference in monthly growth of ARM comparing females to males

Multiple Linear Regression Model

- ▶ Y is a random variable representing the outcome of interest in the population
- ▶ The explanatory variables, X_1, X_2, \dots, X_p are fixed/known (not random or measured with error)
- ▶ Sample of size n is observed, data are: $(y_i, X_{1i}, X_{2i}, \dots, X_{pi})$

$$\Rightarrow Y_i = \underbrace{\mu_i(\beta, X_i)}_{\text{systematic component}} + \underbrace{\varepsilon_i}_{\text{random component}}$$

outcome random variable

- ▶ X is the design matrix or table combining all the explanatory variables $[1s, X_1, X_2, \dots, X_p]$
- ▶ X_i is the row of the design matrix corresponding to subject i \hookrightarrow column vector of length n
 $(1, X_{1i}, X_{2i}, \dots, X_{pi})$

Multiple Linear Regression Model

$$Y_i = \mu_i(\beta, X_i) + \varepsilon_i$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} \quad p+1 \text{ rows} \times 1$$

- Systematic component:

$$\mu_i(\beta, X_i) = \beta_0 + \beta_1 X_{i1} + \beta_2 X_{i2} + \dots + \beta_p X_{ip}$$


- ε_i is the random components: $\varepsilon_i \sim N(0, \sigma^2)$, $\text{Cor}(\varepsilon_i, \varepsilon_j) = 0$ for all $i \neq j$

- The least squares solution finds the values of β that minimize:

$$\sum_{i=1}^n (y_i - \mu_i(\beta, X_i))^2$$

Find β s.t.

the residual
sums of squares
are minimized



Least squares solution: simple linear regression

$$Y_i = \beta_0 + \beta_1 X_i + \varepsilon_i$$

Least squares solution is:

$$\hat{\beta}_1 = \frac{\sum (Y_i - \bar{Y})(X_i - \bar{X})}{\sum (X_i - \bar{X})^2} \quad \left. \vphantom{\hat{\beta}_1} \right\} \frac{\text{Covariance between } Y \text{ and } X}{\text{Variance of } X}$$

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X}$$

If we were to
standardize Y and X ,
 $\text{mean}(Y) = \text{mean}(X) = 0$
 $\text{SD}(Y) = \text{SD}(X) = 1$
 $\hat{\beta}_1 = r$ Pearson
correlation
coefficient

Maximum likelihood inference in MLR

- Start with the MLR:

$$\Rightarrow \underbrace{Y_i = \beta_0 + \beta_1 X_{i1} + \dots + \beta_p X_{ip}}_{\text{Data: } (y_i, X_i)} + \varepsilon_i, \quad \varepsilon_i \sim \mathcal{N}(0, \sigma^2) \text{ independent}$$

- Other notation:

Y_i = Random variable, y_i = observation

$\begin{bmatrix} Y_1 \\ \vdots \\ Y_n \end{bmatrix}$ Y , \mathbf{y} = vectors (lists) of independent random variables or observations

X_i = row vector containing explanatory variables for subject i

$$\hat{Y}_i = \text{RV} = \hat{\beta}_0 + \hat{\beta}_1 X_{i1} + \dots + \hat{\beta}_p X_{ip}$$

$$R_i = \text{RV} = Y_i - \hat{Y}_i \text{ residual}$$



Likelihood function definition

► **Model:** under the assumption $\epsilon_i \sim N(0, \sigma^2)$, $Y_i = \beta'X_i$, $X_i = \text{fixed}$
 $Y_i \sim N(\mu_i(\beta, X_i), \sigma^2)$

► **Probability density function:**

$$f(\underline{y} | \underline{\mu}(\beta, X), \sigma^2) = \prod_{i=1}^n f(y_i | \mu_i(\beta, X_i), \sigma^2)$$

pdf is a function of \underline{y} with $\mu_i(\beta, X_i)$ and σ^2 fixed

► **Likelihood function:** $L(\mu(\beta, X), \sigma^2 | \underline{y}) = \prod_{i=1}^n L(\mu_i(\beta, X_i), \sigma^2 | y_i)$


likelihood function is viewed as a
function of $\mu_i(\beta, X_i)$ and σ^2 for fixed \underline{y}

Identify the values of β and σ^2 that maximize
 L given fixed \underline{y}

Maximum likelihood estimation under gaussian residuals

- Likelihood function, rely on the normality assumption and independence

$$\begin{aligned} L(\beta, \sigma^2 | y) &= \prod_{i=1}^n L(u_i(\beta, x_i), \sigma^2 | y_i) \\ &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y_i - u_i(\beta, x_i))^2\right) \\ &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y_i - \beta_0 - \beta_1 x_{i1} + \beta_2 x_{i2} - \dots - \beta_p x_{ip})^2\right) \end{aligned}$$



Maximum likelihood estimation under gaussian residuals

► Log Likelihood Function

$$l(\beta, \sigma^2 | \underline{y}) = \text{Log } L(\beta, \sigma^2 | \underline{y})$$

$$\Rightarrow = \sum_{i=1}^n \left(-\frac{1}{2} \log(2\pi) - \log(\sigma) - \frac{1}{2\sigma^2} (y_i - \mu_i(\beta, x_i))^2 \right)$$

Find $\hat{\beta}$ and $\hat{\sigma}^2$ that maximize $l(\beta, \sigma^2 | \underline{y})$
by differentiating with respect to β and σ^2 ,
setting these derivatives = 0 and solving for
 β and σ^2

Maximum likelihood estimation under gaussian residuals

► Solution for β_j $j=0, \dots, p$

$$l(\beta, \sigma^2 | y) = \sum_{i=1}^n \left(\underbrace{-\frac{1}{2} \log(2\pi) - \log(\sigma)}_{\text{constant}} - \frac{1}{2\sigma^2} (y_i - \mu_i(\beta, X_i))^2 \right)$$

wrt β $l(\beta, \sigma^2 | y) \propto \sum_{i=1}^n \frac{-1}{2\sigma^2} (y_i - \mu_i(\beta, X_i))^2$

Score equation for β_j

$$U_{\beta_j}(\beta | \sigma^2) = \frac{\partial}{\partial \beta_j} l(\beta, \sigma^2 | y)$$

$$= \frac{\partial}{\partial \beta_j} \sum_{i=1}^n \frac{-1}{2\sigma^2} (y_i - (\beta_0 + \beta_1 X_{i1} + \dots + \beta_p X_{ip}))^2$$

$$= -\frac{2}{2\sigma^2} \sum_{i=1}^n (y_i - \mu_i(\beta, X_i)) (-X_{ij})$$


↳ a score equation for each $\beta \Rightarrow p+1$ score equations
 $\Rightarrow 0$ solve for β_j $p+1$ unknowns

Maximum likelihood estimation under gaussian residuals

► Solution for β_j

$$U_{\beta} = \sum_{i=1}^n (y_i - \mu_i(\beta, x_i)) \begin{bmatrix} 1 \\ x_{1i} \\ x_{2i} \\ \vdots \\ x_{pi} \end{bmatrix} = 0$$

$p+1 \times 1$



Maximum likelihood estimation under gaussian residuals

► Solution for σ^2

Given the MLEs $\hat{\beta}$, derive score equation for σ^2

$$U_{\sigma^2}(\hat{\beta}) = \sum_{i=1}^n \left(-\frac{1}{2\sigma^2} + \frac{1}{2(\sigma^2)^2} (y_i - \mu_i(\hat{\beta}, X_i))^2 \right)$$

$\Rightarrow 0$ and solve for σ^2


$$\Rightarrow \hat{\sigma}_{MLE}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \mu_i(\hat{\beta}, X_i))^2$$

$$E(\hat{\sigma}_{MLE}^2) = \frac{n - (p+1)}{n} \sigma^2$$


In practice, we use

$$\hat{\sigma}^2 = \frac{1}{n - (p+1)} \sum (y_i - \mu_i(\hat{\beta}, X_i))^2$$


MLEs for simple linear regression



MLEs for simple linear regression



MLEs for simple linear regression



Take away messages



Take away messages



Next time....

- ▶ Vector / Matrix representation of MLR
- ▶ Geometry of least squares
- ▶ Distribution of MLEs for regression parameters

