



JOHNS HOPKINS
BLOOMBERG SCHOOL
of PUBLIC HEALTH

Lecture 13

More with Linear Mixed Models

The material in this video is subject to the copyright of the owners of the material and is being provided for educational purposes under rules of fair use for registered students in this course only. No additional copies of the copyrighted work may be made or distributed.

Subject specific or random effects models

- ▶ Consider the data generating structure within the NEPAL1 and NEPAL2 simulated datasets:
 - ▶ Children are enrolled between 1 and 5 months of age
 - ▶ Children are followed over time and growth in weight is recorded every 4 months for a total of 5 assessments (enrollment + 4 follow-ups)
- ▶ For each child, we can think of the child's growth:

$$Y_{ij} = \beta_{0i} + \beta_{1i}age_{ij} + \beta_{2i}(age_{ij} - 6)^+ + e_{ij}$$



Subject specific or random effects models

- ▶ The β describe characteristics of the specific children and we assume that these characteristics can vary from child to child, specifically,

$$\begin{bmatrix} \beta_{0i} \\ \beta_{1i} \\ \beta_{2i} \end{bmatrix} = \begin{bmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} b_{0i} \\ b_{1i} \\ b_{2i} \end{bmatrix}$$

$$\beta_i = \beta + b_i, b_i \sim MVN(0, D), D = \begin{bmatrix} \tau_0^2 & \tau_{01} & \tau_{02} \\ \tau_{01} & \tau_1^2 & \tau_{12} \\ \tau_{02} & \tau_{12} & \tau_2^2 \end{bmatrix}$$



Visualization



General Model

We can rewrite the model above as:

$$Y_{ij} = (\beta_0 + b_{0i}) + (\beta_1 + b_{1i})age_{ij} + (\beta_2 + b_{2i})(age_{ij} - 6)^+ + e_{ij}$$

In vector notation,

$$Y_{ij} = \begin{bmatrix} 1 \\ age_{ij} \\ (age_{ij} - 6)^+ \end{bmatrix}' \begin{bmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} 1 \\ age_{ij} \\ (age_{ij} - 6)^+ \end{bmatrix}' \begin{bmatrix} b_{0i} \\ b_{1i} \\ b_{2i} \end{bmatrix} + e_{ij}$$

Even more generally,

$$Y_{ij} = X_{ij}'\beta + Z_{ij}'b_i + e_{ij}$$

where $b_i \sim MVN(0, D)$, e_{ij} iid $N(0, \sigma^2)$ and b_i and e_{ij} are independent!



Means and Variances

- In the random effects model, we express the mean function for an individual subject as:

$$E(Y_{ij}|X_{ij}, b_i) = X_{ij}\beta + Z_{ij}b_i$$

- We can express the population mean (i.e. the average over all subjects) as:

$$E(Y_{ij}|X_{ij}) = E[E(Y_{ij}|X_{ij}, b_i)] = E[X_{ij}\beta + Z_{ij}b_i] = X_{ij}\beta$$

- We can derive the variance of Y_{ij} as

$$Var(Y_{ij}|X_{ij}) = E_{b_i}[Var(Y_{ij}|X_{ij}, b_i)] + Var_{b_i}[E(Y_{ij}|X_{ij}, b_i)]$$

$$Var(Y_{ij}|X_{ij}) = E_{b_i}[\sigma^2] + Var_{b_i}[X_{ij}'\beta + Z_{ij}'b_i]$$

$$Var(Y_{ij}|X_{ij}) = \sigma^2 + Z_{ij}'DZ_{ij}$$



Correlation

- ▶ Assume a random intercept only model:

- ▶ $Y_{ij} = \beta_{0i} + \beta_1 age_{ij} + \beta_2 (age_{ij} - 6)^+ + e_{ij}, \beta_{0i} \sim N(\beta_0, \tau_0^2), e_{ij} \sim N(0, \sigma^2), Cov(\beta_{0i}, e_{ij}) = 0$
- ▶ $Y_{ij} = \beta_0 + b_{0i} + \beta_1 age_{ij} + \beta_2 (age_{ij} - 6)^+ + e_{ij}, b_{0i} \sim N(0, \tau_0^2), e_{ij} \sim N(0, \sigma^2), Cov(b_{0i}, e_{ij}) = 0$
- ▶ $Corr(Y_{ij}, Y_{ik}) = \frac{\tau_0^2}{\tau_0^2 + \sigma^2}$

- ▶ Assume a random intercept and random slope for age model:

- ▶ $Y_{ij} = \beta_0 + b_{0i} + (\beta_1 + b_{1i}) age_{ij} + \beta_2 (age_{ij} - 6)^+ + e_{ij}, \text{ where}$

$$b_{0i} \sim N(0, \tau_0^2), b_{1i} \sim N(0, \tau_1^2), Cov(b_{0i}, b_{1i}) = \tau_{01}, e_{ij} \sim N(0, \sigma^2), Cov(b_{0i}, e_{ij}) = 0, Cov(b_{1i}, e_{ij}) = 0$$

- ▶ $Corr(Y_{ij}, Y_{ik}) =$



Revisit NEPAL₁ analysis AND do another example, NEPAL₂

