



JOHNS HOPKINS
BLOOMBERG SCHOOL
of PUBLIC HEALTH

Lecture 5

The classical linear regression model


The material in this video is subject to the copyright of the owners of the material and is being provided for educational purposes under rules of fair use for registered students in this course only. No additional copies of the copyrighted work may be made or distributed.

Review of key concepts from Lecture 3 and 4

► Simple linear regression model

► $ARM = B_0 + B_1(\text{age} - 6) + e$, $e \sim N(0, \sigma^2)$, independent

B_0 \rightarrow systematic component \rightarrow random component



B_0 = average ARM among 6 month old children

B_1 = difference in average ARM comparing children who vary in age by 1-month


Review of key concepts from Lecture 3 and 4

► Sex adjusted relationship between ARM and age

► $\text{ARM} = B_0 + B_1(\text{age} - 6) + B_2 \text{Female} + e$, $e \sim N(0, \sigma^2)$, independent

male : $\text{ARM} = B_0 + B_1(\text{age} - 6) + e$

female : $\text{ARM} = B_0 + B_1(\text{age} - 6) + B_2 + e$
 $= (B_0 + B_2) + B_1(\text{age} - 6) + e$



B_2 : Difference in mean ARM comparing females to males who are the same age

Controlling for
= adjusting for


B_0 = Average ARM among male children who are 6 months of age

B_1 = difference in mean ARM comparing two children who have the same sex but differ in age by 1 month

Review of key concepts from Lecture 3 and 4

► Height adjusted relationship between ARM and age

- $ARM = B_0 + B_1 (\text{age} - \underline{6}) + B_2 (\text{HT} - \underline{62}) + e$, $e \sim N(0, \sigma^2)$, independent




Review of key concepts from Lecture 3 and 4

- Effect modification: Is the ARM vs. age relationship the same or different by sex

- $ARM = B_0 + B_1 (age - 6) + B_2 \text{ Female} + B_3 (age - 6) \text{ Female} + e$, $e \sim N(0, \sigma^2)$, independent

male: $ARM = B_0 + B_1 (age - 6) + e$

female: $ARM = (\beta_0 + B_2) + (B_1 + B_3) (age - 6) + e$



$E(\$) = B_0 + B_1(\text{older}) + B_2(\text{low income}) + B_3(\text{older}) \times (\text{low income})$

Category	Effect
younger higher income	B_0
older only	B_1
low income only	B_2
Both	B_3

Multiple Linear Regression Model

- ▶ Y is a random variable representing the outcome of interest in the population
 $n_y = \text{observations}$
- ▶ The explanatory variables, X_1, X_2, \dots, X_p are fixed/known (not random or measured with error)
- ▶ Sample of size n is observed, data are: *$(y_i, X_{1i}, X_{2i}, \dots, X_{pi})$*

$$\rightarrow \underline{Y_i} = \underbrace{\mu_i(\beta, X_i)}_{\substack{\text{systematic} \\ E(Y_i | X_i)}} + \underbrace{\varepsilon_i}_{\text{random component}}$$

- ▶ X is the design matrix
- ▶ X_i is the row of the design matrix corresponding to subject i

$$X = \begin{bmatrix} 1 & X_{11} & X_{21} & \dots & X_{p1} \\ 1 & X_{12} & X_{22} & \dots & X_{p2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & X_{1n} & X_{2n} & \dots & X_{pn} \end{bmatrix}$$

$X_n \leftarrow$

Multiple Linear Regression Model

$$Y_i = \mu_i(\beta, X_i) + \varepsilon_i$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix}_{p+1 \times 1}$$

- ▶ Systematic component:

- ▶ $\mu_i(\beta, X_i) = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_p X_{pi}$

- ▶ ε_i is the random components: $\varepsilon_i \sim N(0, \sigma^2)$, $\text{Cov}(\varepsilon_i, \varepsilon_j) = 0$ for all $i \neq j$
- ▶ The least squares solution finds the values of β that minimize:

$$\sum_{i=1}^n (y_i - \mu_i(\beta, X_i))^2$$
$$\sum_{i=1}^n (y_i - \beta_0 - \beta_1 X_{1i} - \beta_2 X_{2i} - \dots - \beta_p X_{pi})^2$$

Least squares solution: simple linear regression

$$SLR = p = 1 \quad Y_i = \beta_0 + \beta_1 X_i + \varepsilon_i$$

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (y_i - \bar{y})(x_i - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

covariance
between y and x
variance in x

Standardize y and x
to have mean 0
and $SD = 1$

$\hat{\beta}_1 = r$ Pearson
correlation
coefficient

Maximum likelihood inference in MLR

- Start with the MLR:

$$Y_i = \beta_0 + \beta_1 X_{1i} + \dots + \beta_p X_{pi} + \varepsilon_i$$

$\varepsilon_i \sim N(0, \sigma^2)$ and independent

Data: (y_i, X_i) for $i = 1, \dots, n$

- Other notation:

$$Y_i = RV \quad y_i = \text{observation}$$

$$Y = \text{vector of RVs} \quad y = \text{vector of observations}$$

$$\tilde{X}_i = i\text{th row of design matrix } X$$

$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_{1i} + \dots + \hat{\beta}_p X_{pi}$$

$$\hat{\varepsilon}_i = Y_i - \hat{Y}_i$$

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}$$

Likelihood function definition

- ▶ **Model:** systematic component $\mu_i(\beta, X_i)$ plus random component $\varepsilon_i \sim N(0, \sigma^2)$
$$Y_i \sim N(\mu_i(\beta, X_i), \sigma^2)$$
- ▶ **Probability density function:**
$$f(y | \mu(\beta, X), \sigma^2) = \prod_{i=1}^n f(y_i | \mu_i(\beta, X_i), \sigma^2)$$

↳ a function of y given fixed mean $\mu_i(\beta, X_i)$ and variance σ^2
- ▶ **Likelihood function:**
$$L(\mu(\beta, X), \sigma^2 | y) = \prod_{i=1}^n L(\mu_i(\beta, X_i), \sigma^2 | y_i)$$

↳ a function of $\mu_i(\beta, X_i)$ and σ^2 given the observed data y_i

Maximum likelihood estimation under gaussian residuals

► Likelihood function $y_i \sim \mathcal{N}(\mu_i(\beta, X_i), \sigma^2)$

$$\begin{aligned} L(\beta, \sigma^2 | y) &= \prod_{i=1}^n L(\mu_i(\beta, X_i), \sigma^2 | y_i) \\ &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y_i - \underbrace{\mu_i(\beta, X_i)})^2\right) \\ &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2} (y_i - \beta_0 - \beta_1 X_{i1} - \dots - \beta_p X_{ip})^2\right) \end{aligned}$$



Maximum likelihood estimation under gaussian residuals

► Log Likelihood Function

$$\begin{aligned} \ell(\beta, \sigma^2 | y) &= \log L(\beta, \sigma^2 | y) \\ &= \sum_{i=1}^n \left(-\frac{1}{2} \log(2\pi) - \log \sigma - \frac{1}{2\sigma^2} (y_i - \mu_i(\beta, X_i))^2 \right) \end{aligned}$$

= To find $\hat{\beta}$ and $\hat{\sigma}^2$ that maximize $\ell(\beta, \sigma^2 | y)$
we take the derivative wrt β and σ^2 ,
set the derivatives = 0 and solve for
 β and σ^2

Maximum likelihood estimation under gaussian residuals

► Solution for β_j

$j = 0, \dots, p$

$$l(\beta, \sigma^2 | y) = \sum_{i=1}^n \left(-\frac{1}{2} \log(2\pi) - \log \sigma - \frac{1}{2\sigma^2} (y_i - \mu_i(\beta, X_i))^2 \right)$$

$$l(\beta, \sigma^2 | y) \propto \sum_{i=1}^n \frac{1}{2\sigma^2} (y_i - \mu_i(\beta, X_i))^2$$

as a function of β

Define a score equation for β_j

$$U_{\beta_j}(\beta | \sigma^2) = \frac{\partial}{\partial \beta_j} l(\beta, \sigma^2 | y)$$

$$= \frac{\partial}{\partial \beta_j} \sum_{i=1}^n -\frac{1}{2\sigma^2} (y_i - \beta_0 - \beta_1 X_{i1} - \dots - \beta_p X_{ip})^2$$

= 0 and
solve for β_j

$$= -\frac{1}{2\sigma^2} \sum_{i=1}^n 2 \times (y_i - \mu_i(\beta, X_i)) (-X_{ij})$$

Maximum likelihood estimation under gaussian residuals

- ▶ Solution for β_j




Maximum likelihood estimation under gaussian residuals


- ▶ Solution for σ^2




MLEs for simple linear regression



MLEs for simple linear regression



MLEs for simple linear regression



Take away messages



Take away messages



Next time....

- ▶ Vector / Matrix representation of MLR
- ▶ Geometry of least squares
- ▶ Distribution of MLEs for regression parameters

