

# Effects of artificial warming on the structural, physiological, and biochemical changes of maize (*Zea mays* L.) leaves in northern China

Yunpu Zheng · Ming Xu · Ruichang Shen ·  
Shuai Qiu

Received: 5 March 2013 / Revised: 17 May 2013 / Accepted: 22 May 2013  
© Franciszek Górski Institute of Plant Physiology, Polish Academy of Sciences, Kraków 2013

**Abstract** We examined the effects of artificial warming on physiological, biochemical, and structural changes in leaves of maize plants (*Zea mays* L.) with a field warming experiment in the North China Plain. Stomatal characters, leaf anatomy and ultrastructure, gas exchange, and carbohydrate and mineral nutrition concentrations were examined using light microscopy, electron microscopy, portable photosynthesis system (Licor-6400), and inductively coupled plasma atomic emission spectroscopy. We found that artificial warming (about 2 °C) increased both the stomatal index and stomatal size, and thus increased net photosynthesis rate ( $A$ ), stomatal conductance ( $g_s$ ), and transpiration rate ( $E$ ). Artificial warming also significantly increased the profile area of chloroplast and mitochondria, but decreased leaf width and thickness, mesophyll thickness, and mesophyll cell size (mainly palisade cell size). In addition, artificial warming also significantly increased the foliar

C:N ratio and soluble sugar contents (glucose, fructose, and sucrose), but not the mineral nutrients and starch contents. Our findings suggest that future global warming may affect the maize growth and production in northern China due to the direct warming effects on the structures (anatomy and ultrastructure), biochemical properties and gas exchanges of the maize leaves.

**Keywords** Elevated temperature · Gas exchange · Image analysis · Leaf anatomy · Mineral nutrition · Carbohydrates · Stomatal characteristics

## Introduction

It is widely evidenced that the concentrations of greenhouse gases (GHG), such as CH<sub>4</sub>, CO<sub>2</sub> and N<sub>2</sub>O, in the atmosphere have dramatically increased since the beginning of the industrial revolution mainly due to fossil fuel combustion and land use change (IPCC 2007). Meanwhile, the increased GHG concentration has led to a rapid rise in global temperature over the past decades (Gunderson et al. 2000; Wigley and Raper 2001). Moreover, it is estimated that the mean global surface temperature will continue to increase by 1.1–6.4 °C over this century depending on GHG emissions (IPCC 2007). This projected global warming is expected to cause shifts in community species composition and distribution (Epstein et al. 2004; Klein et al. 2004; Walker et al. 2006; Parolo and Rossi 2008), changes in ecosystem structure and function (Peñuelas et al. 2004; Biasi et al. 2008a, b; Jägerbrand et al. 2009), and even local/global extinctions of plant species in terrestrial ecosystems (Parmesan and Yohe 2003; Franco et al. 2006; Malcolm et al. 2006). Global warming may also generate profound alterations in plant structure and

Communicated by U. Feller.

Y. Zheng (✉) · M. Xu (✉) · R. Shen · S. Qiu  
Key Laboratory of Ecosystem Network Observation  
and Modeling, Institute of Geographic Sciences and Natural  
Resources Research, Chinese Academy of Sciences,  
11A Datun Road, Beijing 100101, China  
e-mail: zhengyunpu\_000@sina.com

M. Xu  
e-mail: mingxu@crssa.rutgers.edu

Y. Zheng · R. Shen · S. Qiu  
Graduate School, Chinese Academy of Sciences,  
Beijing 100039, China

Y. Zheng · M. Xu  
Department of Ecology, Evolution and Natural Resources,  
Center for Remote Sensing and Spatial Analysis, Rutgers  
University, 14 College Farm Road, New Brunswick,  
NJ 08901, USA

function (Kudo and Suzuki 2003; Llorens et al. 2004a; Yin et al. 2008; Niu et al. 2008; Zhao and Liu 2009), and thus alter biomass accumulation and allocation (Prieto et al. 2009a; Lin et al. 2010; Wang et al. 2012), and/or net primary production of terrestrial plants (Alward et al. 1999; Peñuelas et al. 2007; Lin et al. 2012; Hou et al. 2012).

In addition to the elevated mean global surface temperature, global precipitation may decline and become unevenly distributed both temporally and spatially (IPCC 2007; Chun et al. 2011). Drought stress caused by a decline in precipitation is also a major factor limiting plants growth and yield. The two abiotic stresses drought and warming are becoming increasing concerns because drought and warming often occur simultaneously in the summer in many areas all over the world (Jiang and Huang 2000, 2001). Previous studies reported that drought and warming alone or in combination affect leaf structure and function such as photosynthesis and respiration (Rachmilevitch et al. 2006), and thus the growth and yield of plants under future climate change. Moreover, most plant species have an optimal growth temperature and only grow in a certain temperature range (Jin et al. 2011). Thus, some species are likely to adapt to warmer temperatures by changing their growth and development or by shifting their ranges, provided that the optimum temperatures are not exceeded. In contrast, some species may fail to adapt to this global change and may even become extinct if the air temperature is too high (Thomas et al. 2004; Colwell et al. 2008; van Mantgem et al. 2009). Meanwhile, many recent studies have found that global warming may come with more frequent extreme climate events such as heat waves resulted in more days with extreme high temperatures (Meehl and Tebaldi 2004; Schär et al. 2004; Tebaldi et al. 2006; Fischer and Schär 2010). It should be noted that there is usually a growth temperature threshold for each development stage of plants/crops, and thus plants/crops grow in a temperature exceeding their growth temperature threshold that may result in more damages than exposure to longer warmer period.

Global warming usually affects plant functions through direct effects on metabolic rates, such as photosynthesis and respiration (Tjoelker et al. 1999; Zha et al. 2001; Llorens et al. 2004b; Niu et al. 2008; Han et al. 2009; Prieto et al. 2009b; Albert et al. 2011), and indirect effects on soil nitrogen mineralization and availability (Rustad et al. 2001; Peñuelas et al. 2004; Sardans et al. 2008a, b), soil moisture (Wan et al. 2005; Holsten et al. 2009), and growing season length (Menzel and Fabian 1999; Walther et al. 2002; Cleland et al. 2006). Most of the impacts of global warming on plant functions are resulted from the changes in plant structure, especially in leaves (Niinemets et al. 2007; Jin et al. 2011; Xu et al. 2012; Smith et al. 2012). Many studies found that leaf thickness and mesophyll volume are strongly correlated with leaf area-based

carbon assimilation (Higuchi et al. 1999; Niinemets et al. 2007; Xu et al. 2012) and leaf mass per area (LMA) (Gorsuch et al. 2010). Previous studies found that plants grown at higher temperatures had thinner leaves, which was mainly caused by reductions in the thickness of epidermal, palisade, and spongy layers, and an associated decrease in the size of mesophyll cells (Hartikainen et al. 2009; Gorsuch et al. 2010; Jin et al. 2011). Earlier studies also found that the number and size of chloroplasts are highly correlated with leaf photosynthetic rates (Jin et al. 2011; Xu et al. 2012). For example, in *Arabidopsis thaliana*, it was observed that the number of chloroplast per cell remained unchanged under 2.5 °C warming, but decreased by 22 % under 5 °C warming (relative to a control day/night temperature of 23/18 °C); Meanwhile the size of chloroplasts also decreased with warmer growth temperature (Jin et al. 2011).

The concentration of TNC reflects the balance between net photosynthetic carbon uptake and irreversible investments for structures (plant growth) or loss of carbon. Most studies have found that global warming generally decreases the TNC concentration in plant leaves (Zha et al. 2001; Tingey et al. 2003; Jin et al. 2011; Wang et al. 2012; Smith et al. 2012). In contrast, several studies found that global warming may increase (Djanaguiraman et al. 2011) or have little effect on the TNC concentration in plant leaves (Tjoelker et al. 1999; Xu et al. 2012). For example, Tjoelker et al. (1999) reported that warming had no effect on the carbohydrate concentration in four boreal tree species including *Betula papyrifera*, *Larix laricina*, *Pinus Banksiana*, and *Picea mariana*, but increased the carbohydrate concentration in *Populus tremuloides*. Previous studies have found that global warming increases leaf C:N ratio (Tolvanen and Henry 2001; Olszyk et al. 2003; Biasi et al. 2008a, b; Yang et al. 2011).

Maize (*Zea mays* L.), a widely distributed C<sub>4</sub> plant species, is one of the most important crops in many regions and thus the maize production determines the safety of future food supply all over the world. It is important to note that the global warming will alter evapotranspiration and water use efficiency (WUE) of crop (Thomas 2008; Mo et al. 2009) and thus affect crop physiological process and growth, which lead to alteration of crop yield (Mo et al. 2009; Liu et al. 2010; Guo et al. 2010). Previous studies have found that global warming decreases the maize yield in many areas of the world (Lobell and Asner 2003). The North China Plain (NCP) is one of the most important regions for crop production in China, accounting for approximately 69 % of wheat (*Triticum aestivum* L.) and 35 % of maize (*Zea mays* L.) grain yields of the whole country (Liu et al. 2010). Recent modeling results have shown that global warming may reduce the maize production by 22.8 % for a 5 °C increase and 13.2 % for a

2 °C increase in the NCP (Liu et al. 2010). However, the underlying mechanisms of climate change on the maize production are far from understood, especially the leaf structure, physiology and biochemistry. The objectives of this study are to examine warming effects on: (1) leaf anatomy and ultrastructure, (2) leaf gas exchange, and (3) carbohydrate and nutrient concentration of maize leaves through a field warming experiment with infrared heaters in northern China.

## Materials and methods

### Site description

We conducted this study in the Yucheng Artificial Station operated by the Chinese Academy of Sciences (36°40′–37°12′N, 116°22′–116°45′E; an elevation of 28 m). The station is located in the lower reach of the Yellow River in the NCP. The study area features a semi-arid climate with average annual precipitation of 610 mm and temperature of 13.1 °C. Approximately 70 % of annual precipitation occurs between June and September. The soil could be classified as calcareous fluvisols in the FAO-Uneson system with 66 % silt; 22 % clay; and 12 % sand. The chemical properties of the soil are as follows: pH 8.5, organic matter 1.47 g/kg, total N 0.9 g/kg, total P ( $P_2O_5$ ) 0.2 % and total K ( $K_2O$ ) 2.26 %. Winter wheat (*T. aestivum* L.) and summer maize (*Z. mays* L.) double cropping is predominant in the NCP.

### Warming experiment

The warming experiment features six  $3 \times 4$  m plots with three of the plots serving as treatments and the others as control plots. The warmed plots have been heated continuously since 4 February 2010 using infrared radiators (165 × 1.5 cm, MSR-2420, Kalglo Electronics Inc., Bethlehem, PA) suspended 2.25 m above the ground. In the control plot, one “dummy” heater with the same shape and size as the infrared radiator was also suspended 2.25 m above the ground to simulate the shading effects of the heater. The distance between the control and the warmed plot was approximately 5 m to avoid heating the control plot by the infrared radiator. Air temperature and soil temperature were hourly monitored with PT 100 thermocouples (Unism Technologies Incorporated, Beijing, China) at 2.4 m above and at the depth of 5 cm in the soil. The warming, on average, has increased air, soil, and canopy temperature by  $1.42 \pm 0.18/1.77 \pm 0.24$  (day/night),  $1.68 \pm 0.9/2.04 \pm 0.16$  (day/night),  $2.08 \pm 0.72$  (day), respectively, in comparison with the control during the maize growth period from 24 June to 7 October 2011. Soil moisture in the top 0–10 cm soil layer was also

recorded with a FDS100 soil moisture sensor (Unism Technologies Incorporated, Beijing, China). During the maize growing season, the mean soil moisture (%volume) in the warmed plots was slightly lower than that in the control plots,  $25.04 \pm 0.52$  vs.  $26.02 \pm 0.86$  %, none of which were likely to limit the maize growth.

### Field sampling

Seeds of maize (*Zea mays* L.) were exposed to a dark, cold, and wet treatment at 4 °C for 2 days before planting to promote uniform germination. Thereafter, they were sown in the field soil in the control and warmed plots in 24 June 2011 and both the maize seedlings in the control and warmed plots grew above the soil surface in 1 July 2011. Maize plants were irrigated with below-ground water to avoid drought stress during the growing period from 24 June 2011 to 7 October 2011. Given that the ear leaf is the most important leaf for determining the crop yield, we sampled the fully expanded ear leaves of maize at the heading stage in 24 August 2011, 60 days after sowing, for the following investigations.

### Gas exchange and stomatal characteristics

Three fully expanded leaves were selected from each of five plants grown in ambient temperature and elevated temperature to measure the net  $CO_2$  assimilation rate ( $A$ ), stomatal conductance ( $g_s$ ), transpiration rate ( $E$ ), intercellular  $CO_2$  concentration ( $C_i$ ), and leaf dark respiration ( $R_d$ ) using an Li-6400 Portable Photosynthesis System (LI-COR Inc. Lincoln, Nebraska, USA). The WUE was determined by the value of  $A$  and  $E$ . The measurements were conducted with leaf temperature at 30 °C, PAR at  $1,500 \mu mol m^{-2} s^{-1}$ ,  $CO_2$  concentration at  $380 \mu mol mol^{-1}$ , and cuvette vapor pressure deficit (VPD) at 2.0 kPa.

We randomly selected five fully expanded ear leaves at the heading stage in each of the warming and control plots on 24 August 2011. To determine the stomatal and epidermal cell characteristics, we sampled impressions from the tip, middle, and base sections of the adaxial and abaxial leaf surfaces using colorless nail polish. The adaxial and abaxial epidermis of the leaves were cleaned first by a degreased cotton ball and then carefully smeared with nail varnish from the mid-area between the central vein and the leaf edge for about half an hour. The thin film (approximately  $5 \times 15$  mm) was peeled off from the leaf surface and mounted on a glass slide. Then the thin film was immediately covered with a cover slip and pressured lightly with a fine-point tweezers. The imprints were observed and photographed in the laboratory with a microscope (DM2500, Leica Corp, Germany) equipped with a digital camera (DFC 300-FX, Leica Corp,

Germany). Thirty separate fields of 0.16 mm were analyzed in each section of the sampled leaves. Stomata and epidermal cells at the tip, middle, and base sections on each surface were counted and combined for calculating stomatal density (SD), epidermal cell density, and stomatal index (SI) (the ratio of stomata to epidermal cells) on the adaxial surface and abaxial surface, respectively. Moreover, we also randomly selected six digital photographs from each leaf section of the adaxial and abaxial surfaces to measure the stomatal aperture length, width, and area using Image J quantification software (NIH, USA). Then the data from the tip, middle, and base sections of both leaf surfaces were combined and averaged for estimating the warming effects on stomatal aperture size.

For visualizing and comparing the differences of stomata between the warming and control plots under scanning electron microscopy (SEM), we snapped three pieces ( $2 \times 2$  mm) from the middle section of each leaf with a fixative solution consisting of 2.5 % (v/v) glutaraldehyde (0.1 M phosphate buffer, pH 7.0). Samples were stored at 4 °C and transported to the laboratory as soon as possible. Then the samples were washed six times with the same buffer and postfixed in 1 % (v/v) osmium tetroxide for 3 h at room temperature. After being washed with the same buffer, leaf tissues were passed through an ethanol dehydration series. Then the samples were critical point-dried, mounted on stubs, and coated with gold in a high-vacuum evaporation unit. Samples were examined and photographed at 10 kV under a Quanta 200 scanning electron microscope (FEI Corp, USA).

#### Anatomical measurements and ultrastructure observation

We used the same ear leaves for further anatomical observations and measurements. The internal anatomy of the selected leaves was assessed on leaf sections harvested from the middle of the leaf and prepared for light microscope observation as described by Sage and Williams (1995). Images of leaf cross-sections were used to estimate leaf anatomical features using Image J software (NIH, USA). Leaf mesophyll thickness was measured between epidermal layers at five points in each cross-section and the interveinal distance was measured between two adjacent vascular bundles (Pengelly et al. 2010). The percentage of the leaf cross-section covered by M or BS tissue was determined by laying a grid of 200 random points over cross-sections of images and calculating the proportion of points falling on M or BS cells (Parkhurst 1982; McKown and Dengler 2007; Riyadh et al. 2007).

For examining the ultrastructure of cellular organelles, we took the advantage of the transmission electron microscopes. Five fully expanded ear leaves of maize

plants were dissected and immediately fixed in 2.5 % (v/v) glutaraldehyde (0.1 M phosphate buffer, pH 7.0) for 2 h at 4 °C. Then the samples were washed four times with the same buffer and postfixed in 1 % osmium tetroxide for 3 h. After being washed with the same buffer, leaf tissues were passed through an ethanol dehydration series, and then infiltrated and embedded in Spurr's resin. Sections were cut using an LKB-V ultramicrotome (LKB, Bromma, Sweden). Thin sections were stained with uranyl acetate and lead citrate, then observed and photographed under a transmission electron microscope (JEOL Ltd, Tokyo, Japan).

#### Biochemical composition analysis

The sampled leaves were oven-dried to constant weight at 75 °C, and ground to fine powder with a ball mill (MM2, Fa. Retsch, Haan, Germany). Samples were assayed for non-structural carbohydrates according to Hendrix (1993) and Way and Sage (2008). Glucose, fructose, sucrose, and starch concentrations were determined spectrophotometrically (UV-1750, Shimadzu Corp., Tokyo, Japan), using a glucose kit (GAHK-20, Sigma, St Louis, MO, USA). Phosphoglucose isomerase (P5381-1 KU, Sigma) was used to convert fructose to glucose, and invertase (I-4504, Sigma) was used to convert sucrose to glucose. Total carbon (C) and nitrogen (N) contents in leaves were determined using an elemental analyzer (Vario Max CN; Elementar Corp., Germany). The contents of potassium (K), calcium (Ca), magnesium (Mg), phosphorus (P), sodium (Na), and Iron (Fe) were determined using an ICP-OES (Optima 5300DV, PerkinElmer Corp., USA) (Luo-mala et al. 2005). All the chemical analyses were repeated five times and expressed on a percentage dry matter basis.

#### Statistical analysis

The differences of the physiological, anatomical, and biochemical variables between the warmed and control plots were tested with the Student's *t* test ( $P < 0.05$ ) using software SPSS 13.0 (SPSS Inc., Chicago, IL, USA).

## Results

#### Stomatal characters and gas exchange

We found no significant difference in stomatal density on adaxial and abaxial leaf surfaces between ambient temperature and elevated temperature, whereas stomatal index significantly increased c. 20 and 6 % on the adaxial and abaxial surfaces, respectively, (Table 1). Only a slight increase in SD was observed under elevated temperature

**Table 1** Stomatal and physiological properties of maize leaves grown at ambient temperature (control) or elevated temperature (warming)

Parameter	Control	Warming	% Increase	<i>P</i> value
<b>Stomatal data</b>				
Stomatal density (adaxial number mm <sup>-2</sup> )	56 ± 6	58 ± 9	3.6	0.377
Stomatal density (abaxial number mm <sup>-2</sup> )	77 ± 7	81 ± 6	5.2	0.059
Stomatal index on the adaxial surface (%)	13.8 ± 0.1	16.5 ± 0.3	19.6	0.004
Stomatal index on the abaxial surface (%)	19.4 ± 0.5	20.6 ± 0.8	6.2	0.013
Stomatal aperture length (μm)	36.2 ± 2.1	29.7 ± 3.1	-18.0	<0.0001
Stomatal aperture width (μm)	3.5 ± 0.5	4.4 ± 0.8	25.7	<0.0001
Stomatal aperture area (μm <sup>2</sup> )	110 ± 34	142 ± 38	29.1	0.003
Stomatal aperture area per unit leaf area (mm <sup>2</sup> ) <sup>a</sup>	0.7 ± 0.2	1.0 ± 0.3	39.4	<0.0001
<b>Physiological data<sup>b</sup></b>				
<i>A</i> (μmol m <sup>-2</sup> s <sup>-1</sup> )	12.8 ± 4.4	19.5 ± 4.8	52.3	0.019
<i>g<sub>s</sub></i> (mmol m <sup>-2</sup> s <sup>-1</sup> )	174 ± 78	457 ± 164	163	0.001
<i>E</i> (mmol m <sup>-2</sup> s <sup>-1</sup> )	3.1 ± 1.1	5.6 ± 2.0	80.6	0.017
<i>C<sub>i</sub></i> (μmol mol <sup>-1</sup> )	210 ± 75	262 ± 39	24.8	0.130
<i>R<sub>d</sub></i> (μmol m <sup>-2</sup> s <sup>-1</sup> )	3.7 ± 0.5	2.8 ± 0.4	-24.3	0.006
WUE (mmol mol <sup>-1</sup> )	4.7 ± 2.5	4.0 ± 1.7	14.9	0.521

Values given are mean ± standard deviation for five fully expanded ear leaves sampled from five maize plants (*n* = 5 for physiological data; *n* = 400 for epidermal data). Mean values were compared by Student's *t* test at *P* < 0.05 level

*A* net CO<sub>2</sub> assimilation rate, *g<sub>s</sub>* stomatal conductance, *E* transpiration rate, *C<sub>i</sub>* intercellular CO<sub>2</sub> concentration, *R<sub>d</sub>* dark respiration, WUE water use efficiency

<sup>a</sup> The stomatal aperture area per unit leaf area is calculated as mean stomatal density × stomatal aperture area per stoma

<sup>b</sup> The physiological data were determined at 1,500 μmol m<sup>-2</sup>s<sup>-1</sup> PAR, 2.0 kPa VPD, 380 μmol mol<sup>-1</sup> CO<sub>2</sub>, and 25 °C leaf temperature

compared with ambient temperature (Table 1). Compared with ambient temperature, the mean stomatal aperture length was decreased by 18 %, but the stomatal aperture width and area as well as the stomatal aperture area per unit leaf area at elevated temperature were significantly increased by c. 26, 29, and 39 %, respectively, (Table 1; Fig. 1).

Artificial warming not only changed leaf stomatal features but also affected leaf physiological characteristics of maize plants. Our results showed that warming significantly increased *A*, *g<sub>s</sub>* and *E* (*P* < 0.05), but decreased *R<sub>d</sub>* (*P* < 0.05; Table 1). In contrast, warming had little effect on WUE (Table 1).

#### Anatomy and ultrastructure


We found that artificial warming significantly decreased both the width and thickness of maize leaves (*P* < 0.05), but it barely affected leaf length (Table 2). Leaf mesophyll thickness, as defined in Fig. 2, was c. 10 % less in elevated temperature than ambient temperature. As no differences were observed in the average number of cell layers between the two groups of plants, the difference in leaf mesophyll thickness was partially due to changes in cell size rather than the number of cell layers (Table 2). Warming significantly reduced the distance between

vascular bundles (defined as the interveinal distance in Fig. 2) by c. 10 % and the area of vascular bundle by c. 30 %, resulting in more and smaller vascular bodies as evidenced in Fig. 2. Relative to ambient temperature, the cell size of bundle sheath (BS) was not markedly affected by artificial warming (Table 2; Fig. 2), whereas warming caused a decrease of 22 % in the cells size of mesophyll (M), which was mainly due to the significant decreases in cell area of palisade mesophyll (PM) since the cell area of spongy mesophyll (SM) was not affected by artificial warming (Table 2). However, we found no significant difference in the M-to-BS cell area ratio between the elevated temperature and ambient temperature (Table 2). Similarly, although warming significantly enhanced the % M by c. 13 %, both % BS and the M-to-BS tissue ratio were not significantly changed (Table 2). Moreover, no difference was observed in sub-stomatal cavity area between the ambient and elevated temperature (Table 2), but warming significantly reduced the cell wall thickness by c. 30 % (Table 2; Fig. 3c, e).

Relative to ambient temperature, elevated temperature significantly increased the average length and width of chloroplast by 46 and 50 %, respectively, resulting in a dramatically c. 140 % enhance in the chloroplast profile area (Table 2; Fig. 3b, e). However, warming had little effect on the number of chloroplasts per mesophyll (Fig. 3a, d) and the



**Fig. 1** Scanning electron micrographs (SEM) showed the stomatal and epidermal cell characteristics of maize leaves grown at ambient temperature (**a, b**) and elevated temperature (**c, d**). Note that shorter and wider stomatal apertures were observed on the leaves of maize grown at elevated temperature than those of their counterparts grown at ambient temperature. Bars 10  $\mu\text{m}$  (**a, c**) and 40  $\mu\text{m}$  (**b, d**)



plastoglobuli per chloroplast (Fig. 3b, e; Table 2). In addition, warming also statistically enhanced the mitochondrial size by c. 53 % (Table 2; Fig. 3c, f).

#### Carbohydrate and mineral nutrient concentration

We found that artificial warming profoundly increased the concentration of TNC mainly due to the increase of the concentration of soluble sugar because warming had little effect on the concentration of starch (Table 3). Compared to ambient temperature, the concentration of soluble sugar at elevated temperature was significantly enhanced by c. 44 %, resulting from the dramatic increase in glucose of 25 %, fructose of 53 %, and sucrose of 83 %, respectively, ( $P < 0.05$ ; Table 3). As a result, artificial warming significantly increased the TNC concentration by c. 30 %, although the concentration of starch not affected by warming (Table 3).

Our results showed no significant difference on the concentrations of all the foliar nutrients between the two temperatures on a dry weight (DW) basis, whereas artificial warming significantly increased the C:N ratio in maize leaves ( $P < 0.05$ ). In addition, artificial warming had no effect on the average concentrations of N, P, K, Ca, and Mg (Table 4). Interestingly, we also found that artificial warming substantially enhanced the C:N ratio by c. 10 % although the concentrations of C and N were not significantly affected by elevated temperature (Table 4).

#### Discussion

The stomatal responses to artificial warming may be variable and species dependent (Ferris et al. 1996; Reddy et al. 1998; Kouwenberg et al. 2007). Many previous studies have found that warming affected stomatal density but did

**Table 2** Morphological and anatomical features of maize leaves grown at ambient temperature (control) or elevated temperature (warming)

Parameter	Control	Warming	% Increase	<i>P</i> value
<b>Morphological data</b>				
Leaf length (cm)	78.4 ± 3.1	79.9 ± 2.2	1.9	0.256
Leaf width (cm)	10.2 ± 0.7	9.8 ± 0.6	−3.9	0.017
Leaf thickness (μm)	167 ± 11	150 ± 25	−10.2	<0.0001
Leaf mass per unit area (g m <sup>−2</sup> )	14.7 ± 1.1	14.5 ± 1.2	−1.4	0.768
<b>Anatomical data</b>				
Mesophyll thickness (μm)	122 ± 11	110 ± 20	−9.8	0.001
Number of cell layer	8.1 ± 0.7	8.6 ± 0.8	6.2	0.068
Interveinal distance (μm)	122 ± 19	109 ± 29	−10.7	0.027
Vascular bundle area (μm <sup>2</sup> )	4,946 ± 1,251	3,538 ± 1,568	−28.5	0.001
Sub-stomatal cavity area (μm <sup>2</sup> )	742 ± 271	761 ± 344	2.6	0.851
Palisade mesophyll cell area (μm <sup>2</sup> )	374 ± 81	246 ± 153	−52.0	<0.0001
Spongy mesophyll cell area (μm <sup>2</sup> )	226 ± 88	216 ± 85	−4.4	0.525
Mesophyll cell size (μm <sup>2</sup> )	297 ± 57	231 ± 91	−22.2	<0.0001
Bundle sheath cell size (μm <sup>2</sup> )	487 ± 225	459 ± 170	−5.8	0.446
Mesophyll:bundle sheath (cell area ratio)	0.79 ± 0.42	0.64 ± 0.37	−20.0	0.093
%Mesophyll	51.9 ± 7.4	45.2 ± 7.7	−12.9	0.010
%Bundle sheath	28.1 ± 5.3	27.8 ± 5.8	−1.1	0.872
Mesophyll:bundle sheath (tissue ratio)	1.9 ± 0.5	1.7 ± 0.6	−10.5	0.270
Cell wall thickness (μm)	0.6 ± 0.1	0.4 ± 0.1	−31.5	0.031
<b>Chloroplast and mitochondria data</b>				
Chloroplast length (μm)	6.1 ± 1.2	8.9 ± 0.9	45.9	<0.0001
Chloroplast width (μm)	1.4 ± 0.2	2.1 ± 0.2	50.0	<0.0001
Area of chloroplast profile (μm <sup>2</sup> )	6.3 ± 1.1	15.2 ± 1.4	141.3	<0.0001
Number of chloroplasts per cell	8.8 ± 2.6	11.8 ± 2.4	22.7	0.240
Number of plastoglobuli per chloroplast	8.6 ± 1.1	9.9 ± 1.8	15.3	0.151
Mitochondrial area (μm <sup>2</sup> )	0.17 ± 0.04	0.26 ± 0.06	52.9	0.01


Values given are mean ± standard deviation. Mean values were compared by Student's *t* test at *P* < 0.05 level. The leaf mesophyll thickness was measured between epidermal layers at eight points in each cross-section and the interveinal distance was measured between two adjacent vascular bundles. The length of chloroplast is the longest dimension, and the width of chloroplast is the widest dimension

not influence stomatal index (Beerling and Chaloner 1993; Reddy et al. 1998; Xu and Zhou 2005; Xu et al. 2009a, b). However, several studies showed that warming had no effect on stomatal density but affected stomatal index due to the changes in epidermal cell density (Beerling and Chaloner 1993; Ferris et al. 1996). Moreover, no changes in both stomatal density and stomatal index were observed under warming conditions in some other studies (Apple et al. 2000; Hovenden 2001; Kouwenberg et al. 2007; Fraser et al. 2009). In our study, the artificial warming significantly increased stomatal index, but had no effect on stomatal density on both the adaxial and abaxial leaf surfaces of maize plants, indicating that the increased stomatal index mainly resulted from the reduction in epidermal cell density, given that the stomatal density was not affected. These results suggested that artificial warming may have directly affected the differentiation and expansion of epidermal cells, but had little effect on the initiation of the

number of stomata during ontogenesis. Moreover, we also found that stomatal aperture area was statistically increased by elevated temperature. These results suggested that the stomata of maize leaves in response to global warming may stimulate not only stomatal index but also stomatal aperture area. In addition, leaf stomatal conductance ( $g_s$ ) and transpiration rate ( $E$ ) are mostly dependent on stomatal aperture size and stomatal frequency including stomatal density and stomatal index (Buckley et al. 1997; Hetherington and Woodward 2003; Franks and Beerling 2009; Franks et al. 2009). Our results also showed that artificial warming substantially increased  $g_s$  and  $E$  by 163 and 81 % possibly due to the increases of stomatal index and stomatal aperture size of maize leaves.

Previous studies have not found consistent results on leaf photosynthetic rates in response to warming. Several studies reported no changes in leaf gas exchange rates (Wookey et al. 1994; Nijs et al. 1996; Loik et al. 2000;





**Fig. 2** Light micrographs of cross-section through leaves of maize grown at ambient temperature (**a–c**) and elevated temperature (**d, e**). Note that more and small bundle sheaths were observed in leaves of plants grown in elevated temperature. Moreover, elevated


temperature decreased leaf thickness. *BS* bundle sheath, *SC* substomatal cavity, *PM* palisade mesophyll, *SM* spongy mesophyll. Bar 100  $\mu$ m (**e**)

Starr et al. 2000; Llorens et al. 2004a, b), whereas others found decreases (Callaway et al. 1994; Roden and Ball 1996; Djanaguiraman et al. 2011) or increases in leaf photosynthetic rates (Chapin and Shaver 1996; Apple et al. 2000; Yin et al. 2008; Han et al. 2009; Prieto et al. 2009a; Jin et al. 2011). However, comparisons among studies are difficult because of the different characteristics of warming treatments among the experiments and the different temperature sensitivities and optimal temperatures for photosynthesis among species and ecotypes (Shaw et al. 2000; Niu et al. 2008). Moreover, leaf photosynthetic rates in response to warming are often strongly affected by other interacting factors such as nutrient availability, the internal plant water status and the water vapor pressure difference between the leaf and the surrounding air (Chaves 1991; Llorens et al. 2004a). Our results that artificial warming stimulated the net photosynthesis rates of maize plants by 52 % support conclusions from previous reports that global

warming may increase plant photosynthesis and growth (Huxman et al. 1998; Niu et al. 2008; Yang et al. 2011). Indeed, we found that the enhancement of leaf photosynthesis might be explained by the significant increases in the chloroplast length, chloroplast width, and chloroplast profile area though the number of chloroplasts per cell was not affected. This is also confirmed by our results that warming significantly increased leaf stomatal conductance (Jin et al. 2011). These results suggested that the increased maize photosynthetic rates in the warming plots were attributed to the increased chloroplast profile area and stomatal conductance. It should be noted that the maize plants in the current study were well watered and fertilized, thus they had the optimal growth conditions. However, it is possible that the photosynthetic response to increased temperature may be negative under the nitrogen or other nutrients deficit conditions. In addition, artificial warming resulted in the maize plants characterizing higher photosynthesis,



**Fig. 3** Transmission electron micrographs showed chloroplast, mitochondria, and cell wall ultrastructure of maize leaves grown under ambient temperature (**a–c**) or elevated temperature (**d–f**). Note that larger chloroplasts and mitochondria were observed in leaves of plants grown under elevated temperature. In addition, elevated temperature decreased the thickness of cell wall. *Ch* chloroplast, *Mi* mitochondria, *CW* cell wall, *Pl* plastoglobuli. Bar 2  $\mu\text{m}$  (**a**, **d**) or 1  $\mu\text{m}$  (**b**, **c**, and **e–f**)



bigger stomatal aperture size and stomatal index, and higher leaf transpiration, which may be disadvantageous for maize plants when the periods of drought occurred simultaneously with elevated temperature.

Many of the most productive crops such as maize use the  $C_4$  photosynthetic pathway, which offers  $C_4$  plants the potential to achieve higher rates of leaf photosynthesis and

more efficient use of water and nitrogen than  $C_3$  plants (Osborne and Freckleton 2009; Taylor et al. 2012). In comparison with  $C_3$  plants, the higher photosynthetic capacity of  $C_4$  plants is mainly due to their unique mode of  $\text{CO}_2$  assimilation, featuring strict compartmentation of photosynthetic enzymes into two distinct cell types, mesophyll and bundle sheath (Wang et al. 2009). In

**Table 3** Content of carbohydrates in leaves of maize grown at ambient temperature (control) or elevated temperature (warming)

Carbohydrates (mg g <sup>-1</sup> DW)	Control	Warming	% Increase	<i>P</i> value
Glucose	16.2 ± 2.7	20.2 ± 1.8	24.9	0.048
Fructose	11.4 ± 1.8	17.4 ± 0.9	52.8	0.003
Sucrose	5.1 ± 1.1	9.4 ± 1.9	83.0	0.009
Soluble sugars	32.7 ± 4.6	47.0 ± 1.4	43.7	0.001
Starch	37.5 ± 4.3	45.0 ± 3.6	20.0	0.080
Total non-structural carbohydrates	69.9 ± 5.2	91.3 ± 4.5	30.8	0.006

Values given are mean ± standard deviation. The ear leaves of maize plants were sampled and mixed for the determination of carbohydrates. Mean values ( $n = 5$  samples, with five plants per sample) were compared by Student's *t* test. Total non-structural carbohydrates = soluble sugars + starch

DW dry weight

**Table 4** Nutrient concentration in leaves of maize grown at ambient temperature (control) or elevated temperature (warming)

Element (mg g <sup>-1</sup> DW)	Control	Warming	% Increase	<i>P</i> value
C	436 ± 3	439 ± 3	0.8	0.179
N	28.5 ± 1.6	26.4 ± 0.7	-7.4	0.058
P	2.9 ± 0.4	2.5 ± 0.1	-13.8	0.110
K	12.7 ± 2.2	12.0 ± 1.3	-5.5	0.618
Ca	6.5 ± 0.8	5.4 ± 1.8	-17.0	0.249
Mg	6.9 ± 1.6	5.1 ± 1.6	-26.8	0.077
C:N	15.4 ± 0.7	16.6 ± 0.3	7.8	0.028

Values given are mean ± standard deviation. The ear leaves of maize plants were sampled and mixed for the determination of carbohydrates. Mean values ( $n = 5$  samples, with five plants per sample) were compared by Student's *t* test

DW dry weight

addition, the higher photosynthetic capacity of C<sub>4</sub> plants can also be attributed to the higher CO<sub>2</sub> concentration and the limitation of photorespiration in the leaves of C<sub>4</sub> plants. In this study, we also found that warming significantly increased net CO<sub>2</sub> assimilation rates (52.3 %), which may result from the increases in the chloroplast size, stomatal index, and stomatal conductance. However, artificial warming increased the soluble sugar concentration in maize leaves, which may be a signal to decrease photosynthesis. Moreover, our results that artificial warming decreased the palisade mesophyll cell area by 52 % also suggested a reduction of photosynthesis of maize plants. Therefore, an opposite trends in photosynthesis and growth may be observed with warming experiments continued for several vegetative seasons. In addition, the enhancement in C<sub>4</sub> photosynthesis efficiency may increase the above-

ground biomass accumulation of C<sub>4</sub> plants (Luo et al. 2009). For example, Luo et al. (2009) showed that artificial warming stimulates above-ground biomass accumulation through enhancing C<sub>4</sub> dominance in a North America tallgrass prairie. However, a recent study from the same warming experiment as the current study found that artificial warming increased the above ground biomass but had little effect on the yield of winter wheat. Therefore, our results suggested that future global warming may affect the contribution of agroecosystems to CO<sub>2</sub> sequestration in a warmer world.

It is reported that warming decreased the thickness of plant leaves (Luomala et al. 2005; Jin et al. 2011; Yang et al. 2011). For example, Jin et al. (2011) found that elevated temperature decreased the leaf thickness by 8 % in *A. thaliana*. Similarly, Yang et al. (2011) observed that warming decreased leaf thickness by 10 % in one of the dominant alpine sedge (*Kobresia pygmaea*) in the Qinghai-Tibetan Plateau, China. In this study, our results also showed that artificial warming decreased the leaf thickness by c. 10 %, which was mainly due to the thinner mesophyll tissue in maize leaves under elevated temperature. This thinner mesophyll tissue was mainly attributable to the decreased mesophyll cell size since the number of cell layer of mesophyll tissue was not changed. Moreover, warming significantly decreased palisade mesophyll cell area by 52 %, but had little effect on the spongy mesophyll cell area. These results showed that warming may decrease leaf thickness mainly via reducing the palisade mesophyll cell area of maize leaves. In addition, our results showed that warming decreased interveinal distance and vascular bundle area by c. 10 and 30 %, respectively, (Table 2). We also observed that maize leaves in elevated temperature have more and smaller vascular bundles than those in ambient temperature (Fig. 2). Similar results have been reported recently by Yang et al. (2011), who found that warming significantly decreased the width and length of the vascular bundle by 18 and 22 %, respectively. Our results suggested that more and smaller vascular bundles per unit leaf area may be more efficient for CO<sub>2</sub> assimilation under warmer conditions. Therefore, the stimulated photosynthetic rate may be also attributed to the decreases in the number and size of the vascular bundles in maize leaves.

The concentration of non-structural carbohydrates generally decreases in plants grown under warm conditions due to the accelerated leaf respiration (Zha et al. 2001; Tingey et al. 2003; Jin et al. 2011; Wang et al. 2012). Interestingly, in this study, we found that artificial warming significantly increased the TNC concentration, especially the concentration of soluble sugars including glucose, fructose, and sucrose. Similarly, a recent study also showed that artificial warming increased the concentration of

soluble sugar in soybean leaves (Djanaguiraman et al. 2011). In the current study, we found that artificial warming increased photosynthesis but decreased  $R_d$  (Table 1), thus leading to the carbohydrate accumulation in maize leaves (Table 4). Our results also suggest that warming may alter the sink-source balance, thus changing the carbon allocation in maize plants (Eamus and Jarvis 1989; Körner et al. 1995).

Our result that warming significantly increased the leaf C:N ratio suggests that the maize plants may become N limited in the NCP under future global warming. Other mineral nutrients, such as K, Ca and Mg, also have similar trends though their leaf concentrations were not statistically different between the warming and the control plots. Our results were supported by previous studies that warming will increase foliar C:N ratio in trees (Tjoelker et al. 1999; Olszyk et al. 2003; Wang et al. 2012), shrubs (Tolvanen and Henry 2001; Biasi et al. 2008a, b; Sardans et al. 2008b), and grasses (Yang et al. 2012). In addition, we found that warming had little impact on foliar P concentration of maize plants (Table 4). This result was in accordance with Sardans et al. (2008b), who found that warming had little effect on foliar P concentration in a shrub species *Globularia alypum*. However, Peñuelas et al. (2004) found that warming decreased leaf P concentration by 11 % in *G. alypum*. By contrast, Sardans et al. (2008a, b) reported that warming increased P concentration by 42 % in *Erica multiflora* leaves. These results suggest that the warming effects on foliar P concentration are more complicated and further studies are needed to clarify the differences.

There are many warming methods for simulating climatic warming in field conditions including greenhouse, open-top chamber, soil heating pipes and cables, infrared reflector, and infrared radiator (Niu et al. 2007). In this study, we examined the warming effects on physiological, biochemical, and structural characteristics of maize leaves with infrared radiators in northern China. It is important to note that the warming effects from the suspended infrared radiators on soil temperature and canopy temperature may be changed during the experiment, because the warming effects may also depend on the increase of maize growth and plant size within the artificial period. The growing maize plants are closer and closer to the radiators and simultaneously the soil surface was more and more shaded during the artificial period, which may result in artifacts of warming effects on the characteristics of maize leaves. Therefore, in terms of the increase of soil temperature, the soil was heated with the method of heating pipes and cables may be better for controlling the soil temperature.

**Author contribution** Y. Zheng and M. Xu designed the experiment and wrote the manuscript. Y. Zheng and R.

Shen conducted the experiment. Y. Zheng and S. Qiu analyzed data.

**Acknowledgments** We thank the associate editor Dr. Urs Feller and the two anonymous reviewers for their constructive comments and suggestions on revising the manuscript. This research was partially supported by the National Key Research and Development Program of China (2012CB417103 and 2010CB833500). This research was also partially supported by Forestry Department of Sichuan Province (Metering System Development and Application of the Regional Forestry Carbon Sink and Source, 2009-204). The authors gratefully acknowledge Dr. Joshua Caplan and Dr. Patrick Burgess (Rutgers University, USA) for their comments on the draft of the manuscript.

## References

- Albert KR, Ro-Poulsen H, Mikkelsen TN, Michelsen A, Linden L, Beier C (2011) Interactive effects of elevated CO<sub>2</sub>, warming, and drought on photosynthesis of *Deschampsia flexuosa* in a temperate heath ecosystem. *J Exp Bot* 62:4253–4266
- Alward RD, Detling JK, Milchunas DG (1999) Grassland vegetation changes and nocturnal global warming. *Science* 283:229–231
- Apple ME, Olszyk DM, Ormrod DP, Lewis J, Southworth D, Tingey D (2000) Morphology and stomatal function of Douglas fir needles exposed to climate change: elevated CO<sub>2</sub> and temperature. *Int J Plant Sci* 161:127–132
- Beerling DJ, Chaloner WG (1993) The impact of atmospheric CO<sub>2</sub> and temperature change on stomatal density: observations from *Quercus robur* Lammas leaves. *Ann Bot* 71:231–235
- Biasi C, Meyer H, Ruslimova O, Hämmerle R, Kaiser C, Baranyi C, Daims H, Lashchinsky N, Barsukov P, Richter A (2008a) Initial effects of artificial warming on carbon exchange rates, plant growth and microbial dynamics of a lichen-rich dwarf shrub tundra in Siberia. *Plant Soil* 307:191–205
- Biasi C, Meyer H, Ruslimova O, Hämmerle R, Kaiser C, Baranyi C, Daims H, Lashchinsky N, Barsukov P, Richter A (2008b) Initial effects of artificial warming on carbon exchange rates, plant growth and microbial dynamics of a lichen-rich dwarf shrub tundra in Siberia. *Plant Soil* 307:171–205
- Buckley TN, Farquhar GD, Mott KA (1997) Qualitative effects of patchy stomatal conductance distribution features on gas-exchange calculations. *Plant Cell Environ* 20:867–880
- Callaway RM, DeLucia EH, Thomas EM, Schlesinger WH (1994) Compensatory responses of CO<sub>2</sub> exchange and biomass allocation and their effects on the relative growth rate of ponderosa pine in different CO<sub>2</sub> and temperature regimes. *Oecologia* 98:159–166
- Chapin FS III, Shaver GR (1996) Physiological and growth responses of arctic plants to a field experiment simulating climatic change. *Ecology* 77:822–840
- Chaves MM (1991) Effects of water deficits on carbon assimilation. *J Exp Bot* 42:1–16
- Chun JA, Wang QG, Timlin D, Fleisher D, Reddy VR (2011) Effect of elevated carbon dioxide and water stress on gas exchange and water use efficiency in corn. *Agric For Meteorol* 151:37–384
- Cleland EE, Chiariello NR, Loarie SR, Mooney HA, Field CB (2006) Diverse responses of phenology to global changes in a grassland ecosystem. *PNAS* 103:13740–13744
- Colwell RK, Brehm G, Cardelús CL, Gilman AC, Longino JT (2008) Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *Science* 322:258–261
- Djanaguiraman M, Prasad PVV, Boyle DL, Schapaugh WT (2011) High-temperature stress and soybean leaves: leaf anatomy and photosynthesis. *Crop Sci* 51:2125–2131



- Eamus D, Jarvis PG (1989) The direct effect of increase in the global atmospheric CO<sub>2</sub> concentration on natural and commercial temperature trees and forests. In: Begon M, Fitter AH, Ford ED, Macfayden A (eds) Advantages in ecological research. Academic Press, New York, pp 1–41
- Epstein HE, Calef MP, Walker MD, Chapin FS III, Starfield AM (2004) Detecting changes in arctic tundra plant communities in responses to warming over decadal time scales. *Glob Chang Biol* 10:1325–1334
- Ferris R, Nijs I, Behaeghe T, Impens I (1996) Elevated CO<sub>2</sub> and temperature have different effects on leaf anatomy of perennial ryegrass in spring and summer. *Ann Bot* 78:489–497
- Fischer E, Schär C (2010) Consistent geographical patterns of changes in high-impact European heat waves. *Nat Geosci* 3:398–403
- Franco AMA, Hill JK, Kitschke C, Collingham YC, Roy DB, Fox R, Huntley B, Thomas CD (2006) Impacts of climate warming and habitat loss on extinctions at species' low-latitude range boundaries. *Glob Chang Biol* 12:1545–1553
- Franks PJ, Beerling DJ (2009) Maximum leaf conductance driven by CO<sub>2</sub> effects on stomatal size and density over geologic time. *PNAS* 106:10343–10347
- Franks PJ, Drake PL, Beerling DJ (2009) Plasticity in maximum stomatal conductance constrained by negative correlation between stomatal size and density: an analysis using *Eucalyptus globulus*. *Plant Cell Environ* 32:1737–1748
- Fraser LH, Greenall A, Carlyle C, Turkington R, Friedman CR (2009) Adaptive phenotypic plasticity of *Pseudoroegneria spicata*: response of stomatal density, leaf area and biomass to changes in water supply and increased temperature. *Ann Bot* 103:769–775
- Gorsuch PA, Pandey S, Atkin OK (2010) Temporal heterogeneity of cold acclimation phenotypes in *Arabidopsis* leaves. *Plant Cell Environ* 33:244–258
- Gunderson CA, Norby RJ, Wullschlegel SD (2000) Acclimation of photosynthesis and respiration to stimulated climatic warming in northern and southern populations of *Acer saccharum*: laboratory and field evidence. *Tree Physiol* 20:87–96
- Guo RP, Lin ZH, Mo XG, Yang CL (2010) Responses of crop yield and water use efficiency to climate change in the North China Plain. *Agric Water Manag* 97:1185–1194
- Han C, Liu Q, Yang Y (2009) Short-term effects of artificial warming and enhanced ultraviolet-B radiation on photosynthesis and antioxidant defense of *Picea asperata* seedlings. *Plant Growth Regul* 58:153–162
- Hartikainen K, Nerg AM, Kivimaenpää M, Kontunen-Soppela S, Maenpää M, Oksanen E, Rousi M, Holopainen T (2009) Emissions of volatile organic compounds and leaf structural characteristics of European aspen (*Populus tremula*) grown under elevated ozone and temperature. *Tree Physiol* 29:1163–1173
- Hendrix DL (1993) Rapid extraction and analysis of nonstructural carbohydrates in plant tissues. *Crop Sci* 33:1306–1311
- Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving environmental change. *Nature* 424:901–908
- Higuchi H, Sakuratani T, Utsunomiya N (1999) Photosynthesis, leaf morphology, and shoot growth as affected by temperatures in cherimoya (*Annona cherimola* Mill.) trees. *Sci Hortic* 80:91–104
- Holsten A, Vetter T, Vohland K, Krysanova V (2009) Impact of climate change on soil moisture dynamics in Brandenburg with a focus on nature conservation areas. *Ecol Model* 220:2076–2087
- Hou RX, Ouyang Z, Li Y, Wilson G, Li H (2012) Is the change of winter wheat yield under warming caused by shortened reproductive period? *Ecol Evol* 2:2999–3008
- Hovenden MJ (2001) The influence of temperature and genotype on the growth and stomatal morphology of southern beech, *Nothofagus cunninghamii* (Nothofagaceae). *Aust J Bot* 49:427–434
- Huxman TE, Hamerlynck EP, Loik ME, Smith SD (1998) Gas exchange and chlorophyll fluorescence responses of three southwestern *Yucca* species to elevated CO<sub>2</sub> and high temperature. *Plant Cell Environ* 21:1275–1283
- IPCC (2007) Climate change 2007: the physical science basis. Intergovernmental Panel on Climate Change, Geneva
- Jägerbrand AK, Alatalo JM, Chrimes D, Molau U (2009) Plant community responses to 5 years of simulated climate change in meadow and heath ecosystems at a subarctic-alpine site. *Oecologia* 161:601–610
- Jiang Y, Huang B (2000) Effects of drought or heat stress alone and in combination on Kentucky bluegrass. *Crop Sci* 40:1358–1362
- Jiang Y, Huang B (2001) Physiological responses to heat stress alone or in combination with drought: a comparison between tall fescue and perennial ryegrass. *Hortic Sci* 36:682–686
- Jin B, Wang L, Wang J, Jiang K, Wang Y, Jiang X, Ni C, Wang Y, Teng N (2011) The effect of artificial warming on leaf functional traits, leaf structure and leaf biochemistry in *Arabidopsis thaliana*. *BMC Plant Biol* 11:35
- Klein JA, Harte J, Zhao X (2004) Artificial warming causes large and rapid species loss, dampened by simulated grazing, on the Tibetan Plateau. *Ecol Lett* 7:1170–1179
- Körner C, Pellaen-Riedl S, Van Bel AJE (1995) CO<sub>2</sub> responsiveness of plants: a positive link to phloem loading. *Plant Cell Environ* 18:595–600
- Kouwenberg LLR, Kürschner WM, McElwain JC (2007) Stomatal frequency change over altitudinal gradients: prospects for paleoaltimetry. *Rev Miner Geochem* 66:215–241
- Kudo G, Suzuki S (2003) Warming effects on growth, production, and vegetation structure of alpine shrubs: a five-year experiment in northern Japan. *Oecologia* 135:280–287
- Lin D, Xia J, Wan S (2010) Climate warming and biomass accumulation of terrestrial plants: a meta-analysis. *New Phytol* 188:187–198
- Lin YS, Medlyn BE, Ellsworth DS (2012) Temperature responses of leaf net photosynthesis: the role of component processes. *Tree Physiol* 32:219–231
- Liu SX, Mo XG, Lin ZH, Xu YQ, Ji JJ, Wen G, Richey J (2010) Crop yield responses to climate change in the Huang-Huai-Hai Plain of China. *Agric Water Manag* 97:1195–1209
- Llorens L, Peñuelas J, Estiarte M, Bruna P (2004a) Contrasting growth changes in two dominant species of a Mediterranean shrubland submitted to artificial drought and warming. *Ann Bot* 94:843–853
- Llorens L, Peñuelas J, Beier C, Emmett B, Estiarte M, Tietema A (2004b) Effects of an artificial increase of temperature and drought on the photosynthetic performance of two ericaceous shrub species along a North–South European gradient. *Ecosystems* 7:613–624
- Lobell DB, Asner GP (2003) Climate and management contributions to recent trends in U.S. agricultural yields. *Science* 299:1032
- Loik ME, Redar SP, Harte J (2000) Photosynthetic responses to a climate warming manipulation for contrasting meadow species in the Rocky Mountains, Colorado, USA. *Funct Ecol* 14:166–175
- Luo Y, Sherry R, Zhou X, Wan S (2009) Terrestrial carbon-cycle feedback to climate warming: artificial evidence on plant regulation and impacts of biofuel feedstock harvest. *GCB Bioenergy* 1:62–74
- Luomala EM, Laitinen K, Sutinen S, Kellomäki S, Vapaavuori E (2005) Stomatal density, anatomy and nutrient concentrations of Scots pine needles are affected by elevated CO<sub>2</sub> and temperature. *Plant Cell Environ* 28:733–749
- Malcolm JR, Liu CR, Neilson RP, Hansen L, Hannah L (2006) Global warming and extinctions of endemic species from biodiversity hotspots. *Conserv Biol* 20:538–548



- McKown AD, Dengler NG (2007) Key innovations in the evolution of Kranz anatomy and C<sub>4</sub> vein pattern in *Flaveria* (Asteraceae). *Am J Bot* 94:382–399
- Meehl GA, Tebaldi C (2004) More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* 305:994–997
- Menzel A, Fabian P (1999) Growing season extended in Europe. *Nature* 397:659
- Mo XG, Liu SX, Lin ZH, Guo RP (2009) Regional crop yield, water consumption and water use efficiency and their responses to climate change in the North China Plain. *Agric Ecosyst Environ* 134:67–78
- Niinemets U, Díaz-Espejo A, Flexas J, Galmés J, Warren CR (2007) Role of mesophyll diffusion conductance in constraining potential photosynthetic productivity in the field. *J Exp Bot* 60:2249–2270
- Nijs I, Teughels H, Blum H, Hendrey G, Impens I (1996) Simulation of climate change with infrared heaters reduces the productivity of *Lolium perenne* L. in summer. *Environ Exp Bot* 36:271–280
- Niu SL, Han XG, Ma KP, Wan SQ (2007) Field facilities in global warming and terrestrial ecosystem research. *Chin J Plant Ecol* 31:262–271
- Niu S, Li Z, Xia J, Han Y, Wu M, Wan S (2008) Climate warming changes plant photosynthesis and its temperature dependence in a temperate steppe of northern China. *Environ Exp Bot* 63:91–101
- Olszyk DM, Johnson MG, Tingey DT, Rygielwicz PT, Wise C, VanEss E, Benson A, Storm MJ, King R (2003) Whole-seedling biomass allocation, leaf area, and tissue chemistry for Douglas-fir exposed to elevated CO<sub>2</sub> and temperature for 4 years. *Can J For Res* 33:269–278
- Osborne CP, Freckleton RP (2009) Ecological selection pressures for C<sub>4</sub> photosynthesis in the grasses. *Proc R Soc Sec B* 276:1753–1760
- Parkhurst DF (1982) Stereological methods for measuring internal leaf structure variables. *Am J Bot* 69:31–39
- Parnesan C, Yohe G (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421:37–42
- Parolo G, Rossi G (2008) Upward migration of vascular plants following a climate warming trend in the Alps. *Basic Appl Ecol* 9:100–107
- Pengelly JLL, Sirault XRR, Tazoe Y, Evans JR, Furbank RT, Caemmerer S (2010) Growth of the C<sub>4</sub> dicot *Flaveria bidentis*: photosynthetic acclimation to low light through shifts in leaf anatomy and biochemistry. *J Exp Bot* 61:4109–4122
- Peñuelas J, Gordon C, Llorens L, Nielsen T, Tietema A, Beier C, Bruna P, Emmett B, Estiarte M, Gorissen A (2004) Noninvasive field experiments show different plant responses to warming and drought among sites, seasons, and species in a north–south European gradient. *Ecosystems* 7:598–612
- Peñuelas J, Prieto P, Beier C (2007) Response of plant species richness and primary productivity in shrublands along a north–south gradient in Europe to seven years of artificial warming and drought. Reductions in primary productivity in the heat and drought year of 2003. *Global Change Biol* 13:2563–2581
- Prieto P, Peñuelas J, Llusà J, Asensio D, Estiarte M (2009a) Effects of artificial warming and drought on biomass accumulation in a Mediterranean shrubland. *Plant Ecol* 205:179–191
- Prieto P, Peñuelas J, Llusà J, Asensio D, Estiarte M (2009b) Effects of long-term artificial night-time warming and drought on photosynthesis, *Fv/Fm* and stomatal conductance in the dominant species of a Mediterranean shrubland. *Acta Physiol Plant* 31:729–739
- Rachmilevitch S, DaCosta M, Huang B (2006) Physiological and biochemical indicators for abiotic stress tolerance. In: Huang B (ed) *Plant–environment interaction*. CRC Press, Boca Raton, pp 321–356
- Reddy KR, Robana RR, Hodges HF, Liu XJ, McKinion JM (1998) Interactions of CO<sub>2</sub> enrichment and temperature on cotton growth and leaf characteristics. *Environ Exp Bot* 39:117–129
- Riyadh M, Sage RF, Dengler NG (2007) Diversity of kranz anatomy and biochemistry in C<sub>4</sub> eudicots. *Am J Bot* 94:362–381
- Roden JS, Ball MC (1996) The effect of elevated CO<sub>2</sub> on growth and photosynthesis of two eucalyptus species exposed to high temperatures and water deficits. *Plant Physiol* 111:909–919
- Rustad L, Campbell J, Marion G, Norby R, Mitchell M, Hartley A, Cornelissen J, Gurevitch J (2001) A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to artificial ecosystem warming. *Oecologia* 126:543–562
- Sage TL, Williams EG (1995) Structure, ultrastructure, and histochemistry of the pollen tube pathway in the milkweed *Asclepias exaltata* L. *Sex Plant Reprod* 8:257–265
- Sardans J, Peñuelas J, Prieto P, Estiarte M (2008a) Drought and warming induced changes in P and K concentration and accumulation in plant biomass and soil in a Mediterranean shrubland. *Plant Soil* 306:261–271
- Sardans J, Peñuelas J, Estiarte M, Prieto P (2008b) Warming and drought alter C and N concentration, allocation and accumulation in a Mediterranean shrubland. *Glob Change Biol* 14:2304–2316
- Schär C, Vidale PL, Lüthi D, Frei C, Häberli C, Liniger MA, Appenzeller C (2004) The role of increasing temperature variability in European summer heatwaves. *Nature* 427:332–336
- Shaw MR, Loik ME, Harte J (2000) Gas exchange and water relations of two Rocky Mountain shrub species exposed to a climate change manipulation. *Plant Ecol* 146:197–206
- Smith RA, Lewis JD, Ghannoum O, Tissue DT (2012) Leaf structural responses to pre-industrial, current and elevated atmospheric [CO<sub>2</sub>] and temperature affect leaf function in *Eucalyptus sideroxylon*. *Funct Plant Biol* 39:285–296
- Starr G, Oberbauer SF, Pop EW (2000) Effects of lengthened growing season and soil warming on the phenology and physiology of *Polygonum bistorta*. *Glob Change Biol* 6:357–369
- Taylor SH, Franks PJ, Hulme SP, Spriggs E, Christin PA, Edwards EJ, Woodward FI, Osborne CP (2012) Photosynthetic pathway and ecological adaptation explain stomatal trait diversity amongst grasses. *New Phytol* 193:387–396
- Tebaldi C, Hayhoe K, Arblaster JM, Meehl GA (2006) Going to the extremes. An intercomparison of model-simulated historical and future changes in extreme events. *Clim Change* 79:185–211
- Thomas A (2008) Agriculture irrigation demand under present and future climate scenarios in China. *Glob Planet Change* 60:306–326
- Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC, Erasmus BF, De Siqueira MF, Grainger A, Hannah L, Hughes L, Huntley B, Van Jaarsveld AS, Midgley GF, Miles L, Ortega-Huerta MA, Peterson AT, Phillips OL, Williams SE (2004) Extinction risk from climate change. *Nature* 427:145–148
- Tingey DT, Mckane RB, Olszyk DM, Johnson MG, Rygielwicz P, Henrylee E (2003) Elevated CO<sub>2</sub> and temperature alter nitrogen allocation in Douglass-fir. *Glob Change Biol* 9:1038–1050
- Tjoelker MG, Reich PB, Oleksyn J (1999) Changes in leaf nitrogen and carbohydrates underlie temperature and CO<sub>2</sub> acclimation of dark respiration in five boreal tree species. *Plant Cell Environ* 22:767–778
- Tolvanen A, Henry GHR (2001) Responses of carbon and nitrogen concentrations in high arctic plants to artificial warming. *Can J Bot* 79:711–718

- van Mantgem PJ, Stephenson NL, Byrne JC, Daniels LD, Franklin JF, Fulé PZ, Harmon ME, Larson AJ, Smith JM, Taylor AH, Veblen TT (2009) Widespread increase of tree mortality rates in the western United States. *Science* 323:521–524
- Walker MD, Wahren CH, Hollister RD (2006) Plant community responses to artificial warming across the tundra biome. *PNAS* 103:1342–1346
- Walther GR, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin JM, Hoegh-Guldberg O, Bairlein F (2002) Ecological responses to recent climate change. *Nature* 416:389–395
- Wan S, Hui D, Wallace L, Luo Y (2005) Direct and indirect effects of artificial warming on ecosystem carbon processes in a tallgrass prairie. *Glob Biogeochem Cycles* 19:GB2014
- Wang X, Gowik U, Tang H, Bowers JE, Westhoff P, Paterson AH (2009) Comparative genomic analysis of C<sub>4</sub> photosynthetic pathway evolution in grasses. *Genome Biol* 10:R68
- Wang J, Duan B, Zhang Y (2012) Effects of artificial warming on growth, biomass allocation, and needle chemistry of *Abies faxoniana* in even-aged monospecific stands. *Plant Ecol* 213:47–55
- Way DA, Sage RF (2008) Elevated growth temperatures reduce the carbon gain of black spruce [*Picea mariana* (Mill.) B.S.P.]. *Glob Change Biol* 14:624–636
- Wigley TML, Raper SCB (2001) Interpretation of high projections for global-mean warming. *Science* 293:451–454
- Wookey PA, Welker JM, Parsons AN, Press MC, Callaghan TV, Lee JA (1994) Differential growth, allocation and photosynthetic responses of *Polygonum viviparum* to simulated environmental change at a high arctic polar semi-desert. *Oikos* 70:131–139
- Xu CY, Salih A, Ghannoum O, Tissue DT (2012) Leaf structural characteristics are less important than leaf chemical properties in determining the response of leaf mass per area and photosynthesis of *Eucalyptus saligna* to industrial-age changes in [CO<sub>2</sub>] and temperature. *J Exp Bot* 63:5829–5841
- Xu ZZ, Zhou GS (2005) Effects of water stress and high nocturnal temperature on photosynthesis and nitrogen level of a perennial grass *Leymus chinensis*. *Plant Soil* 269:131–139
- Xu ZZ, Zhou GS, Shimizu H (2009a) Effects of soil drought with nocturnal warming on leaf stomatal traits and mesophyll cell ultrastructure of a perennial grass. *Crop Sci* 49:1843–1851
- Xu Z, Hu T, Wang K, Zhang Y, Xian J (2009b) Short-term responses of phenology, shoot growth and leaf traits of four alpine shrubs in a timberline ecotone to simulated global warming, Eastern Tibetan Plateau, China. *Plant Spec Biol* 24:27–34
- Yang Y, Wang G, Klanderud K, Yang L (2011) Responses in leaf functional traits and resource allocation of a dominant alpine sedge (*Kobresia pygmaea*) to climate warming in the Qinghai-Tibetan Plateau permafrost region. *Plant Soil* 349:377–387
- Yin H, Liu Q, Lai T (2008) Warming effects on growth and physiology in the seedlings of the two conifers *Picea asperata* and *Abies faxoniana* under two contrasting light conditions. *Ecol Res* 23:459–469
- Zha T, Ryppö A, Wang K, Kellomäki S (2001) Effects of elevated carbon dioxide concentration and temperature on needle growth, respiration and carbohydrate status in field-grown Scots pine during the needle expansion period. *Tree Physiol* 21:1279–1287
- Zhao C, Liu Q (2009) Growth and physiological responses of *Picea asperata* seedlings to elevated temperature and to nitrogen fertilization. *Acta Physiol Plant* 31:163–173