

---

# CONGRUENCE

---

REVISION: 2983

20 OCTOBER 2025

AZIZ MANVA

[AZIZMANVA@GMAIL.COM](mailto:AZIZMANVA@GMAIL.COM)

ALL RIGHTS RESERVED

# TABLE OF CONTENTS

TABLE OF CONTENTS .....	2
-------------------------	---

1. CONGRUENCE.....	3
--------------------	---

1.1 SSS Theorem	3
-----------------	---

1.2 RHS Theorem	7
1.3 SAS Theorem	10
1.4 ASA/SAA Theorem	11
1.5 Applications	16
1.6 Further Topics	19

# 1. CONGRUENCE

## 1.1 SSS Theorem

### A. Basics

#### 1.1: Summary of Congruence Theorems

Theorem	Meaning
SSS	Side-Side-Side
SAS	Side-Angle-Side
SAA / ASA	Side-Angle-Angle Angle Side Angle
RHS/Hy Leg	Right Angle, Hypotenuse, Side OR Hypotenuse Leg

#### 1.2: Congruence Definition

Two figures are congruent if they are the same size and shape.

- This definition is very simple, but very powerful. Many problems can be solved using congruence.
- We will focus on congruence of triangles, since it is a fundamental skill. It is very useful in the geometry of other shapes, such as quadrilaterals, circles and many other shapes.

#### 1.3: Corresponding Parts of Congruent Triangles are Congruent (CPCTC)


Corresponding parts of congruent triangles are congruent.

- If two triangles are congruent, then their corresponding parts are also congruent.
- The order in which you mention the vertices of two congruent triangles is important.

#### Example 1.4

$$\triangle ABC \cong \triangle XYZ$$

Identify the corresponding sides and corresponding angles which are congruent.



For angles, we need to pick letters in corresponding positions:

$$\text{First Position: } \triangle ABC \cong \triangle XYZ \Rightarrow \angle A \cong \angle X$$

$$\text{2nd Position: } \triangle ABC \cong \triangle XYZ \Rightarrow \angle B \cong \angle Y$$

$$\text{3rd Position: } \triangle ABC \cong \triangle XYZ \Rightarrow \angle C \cong \angle Z$$

We need to do the same for sides:


$$\text{1st and 2nd Position: } \triangle ABC \cong \triangle XYZ \Rightarrow AB \cong XY$$

$$\text{2nd and 3rd Position: } \triangle ABC \cong \triangle XYZ \Rightarrow BC \cong YZ$$

$$\text{1st and 3rd Position: } \triangle ABC \cong \triangle XYZ \Rightarrow AC \cong XZ$$

### Example 1.5

Consider the diagram drawn alongside (which is not drawn to scale). Suppose that  $\triangle ABC \cong \triangle YZX$ . Identify the corresponding sides and corresponding angles which are congruent.



Sides:

$$AB \cong YZ, BC \cong ZX, AC \cong YX$$

Angles:

$$\angle A \cong \angle Y, \angle B \cong \angle Z, \angle C \cong \angle X$$

### 1.6: Showing Congruent Parts

Congruent parts are shown using the same symbols.

*Lines are indicated using cuts*

*Angles are indicated using curves*

### Example 1.7


Which parts are congruent in the figure alongside?

*AD has a single cut, BC has a single cut  $\Rightarrow AD = BC$*

*DC has a double cut, AB has a double cut  $\Rightarrow DC = AB$*

*$\angle DAC$  is drawn with a single curve,  $\angle ACB$  is drawn with a single curve  $\Rightarrow \angle DAC = \angle ACB$*


*$\angle CAB$  is drawn with a double curve,  $\angle ACD$  is drawn with a double curve  $\Rightarrow \angle CAB = \angle ACD$*



### 1.8: Reflexive Property

Reflexive property:

*Anything is equal to itself*



### Example 1.9

Use the information in the figure to show that the two triangles are congruent.

Statement	Reason
$AD \cong CB$ $CD \cong AB$ $\angle DAC \cong ACB$ $\angle DCA \cong \angle CAB$	Given
$AC \cong CA$	Reflexive Property
$\angle D \cong \angle B$	Third Angles Theorem

## B. SSS Theorem

There are five theorems for proving the congruence of triangles.

### 1.10: SSS (Side-Side-Side) Congruence Theorem

Two triangles are congruent if:

*three sides of the first triangle are congruent to three sides of the second triangle*

- Order is not mentioned in the property
- Order must be taken into when establishing the congruence

### Example 1.11

Show that a triangle is congruent to itself.

Consider  $\triangle ABC$ :

$$AB \cong AB \text{ (Reflexive Property)}$$

$$BC \cong BC \text{ (Reflexive Property)}$$

$$CA \cong CA \text{ (Reflexive Property)}$$

Hence, by SSS,

$$\triangle ABC \cong \triangle ABC$$

This is an example of the “reflexive” property: Anything is congruent to itself.

### Example 1.12

A right triangle has legs 3 and 4. Another right triangle has hypotenuse 5 and a leg 4. Are the two triangles congruent?

Note: Use SSS Theorem for congruence.

In the first triangle, the sides are:

3,4,5

In the second triangle, the sides are:

3,4,5

By SSS:

*Triangles are congruent*


### 1.13: Order in a Diagram

In a diagram, two congruent triangles need to be oriented the same way. However, if you are able to establish congruence using any order, then the two triangles are congruent.

### Example 1.14

The adjoining diagram (which is not drawn to scale) has two triangles.

Prove or disprove the two triangles to be congruent.



$\triangle ABC \cong \triangle EFD$  by SSS Theorem:

$$AB = EF \text{ (Given)}$$

$$BC = DF \text{ (Given)}$$

$$AC = DE \text{ (Given)}$$

## C. Insufficient Information

### 1.15: Insufficient Information

In certain cases, we may not have enough information to decide whether two triangles are congruent.

#### Example 1.16

A right triangle has sides 3 and 4. Another right triangle has sides 4 and 5. Are the two triangles congruent?

##### Case I

Suppose the first triangle has

$$\text{Legs } 3 \text{ and } 4 \Rightarrow \text{Hypotenuse} = 5 \Rightarrow \text{Sides} = \{3, 4, 5\}$$

Suppose the second triangle has

$$\text{Leg } 4 \text{ and Hypotenuse } 5 \Rightarrow \text{Second Leg} = 4 \Rightarrow \text{Sides} = \{3, 4, 5\}$$

By SSS:

*Two Triangles are congruent*

##### Case II

Suppose the first triangle has

$$\text{Legs } 3 \text{ and } 4 \Rightarrow \text{Hypotenuse} = 5 \Rightarrow \text{Sides} = \{3, 4, 5\}$$

Suppose the second triangle has

$$\text{Legs } 4 \text{ and } 5 \Rightarrow \text{Hypotenuse} = \sqrt{4^2 + 5^2} = \sqrt{16 + 25} = \sqrt{41} \Rightarrow \text{Sides} = \{4, 5, \sqrt{41}\}$$

Then:

*The Two Triangles may or may not be congruent*

## D. Disproving Congruence

### 1.17: Proving Not Congruent using SSS

SSS requires three sides to be congruent. If even one side does not match, then the two triangles will not be congruent.

### 1.18: Triangle Inequality

In a triangle:

- Sum of any two sides is greater than the third side
- Third side is less than sum of the other two sides, and greater than difference of other two sides

#### Example 1.19

Consider non-congruent triangles  $ABC$  and  $XYZ$  with integer side lengths.  $\triangle ABC$  has sides 3, 4 and 5.  $\triangle XYZ$  has sides 3 and 4. The two triangles are not congruent. Determine the sum of the possible values of the third side of  $\triangle XYZ$ .

Using triangle inequality for  $\triangle XYZ$ :

$$\text{Third Side} < 3 + 4 = 7$$

$$\text{Third Side} > 4 - 3 = 1$$

Possible values using the triangle inequality are:

$\{2, 3, 4, 5, 6\}$

But  $\triangle ABC$  is not congruent with  $\triangle XYZ$ . Hence,

$\text{Third Side} \neq 5$

Final answer is:

$$\text{Sum of possible values} = 2 + 3 + 4 + 6 = 15$$

## 1.2 RHS Theorem

### A. SSA: Not a Congruence Theorem

#### 1.20: SSA: Not a Congruence Theorem

If two sides and the *angle which is not included* of one triangle are congruent to two sides and the angle which is not included of a second triangle:

*we cannot conclude anything without further information*

In general:


- Information with SSA will give two triangles that meet the required conditions.
- Hence, this case is called the ambiguous case.

#### Example 1.21

$\triangle ABC$  has  $AB = 1.5$ ,  $AC = 2$  and  $\angle BCA = 40^\circ$ .  $\triangle AGC$  has  $AC = 2$ ,  $AG = 1.5$ , and  $\angle GCA = 40^\circ$ . Are the two triangles congruent?

Without further information, we cannot tell.

For example, the diagram on the right shows the case where the two triangles are not congruent.




### B. RHS

If the angle which is not included is a right angle, then the two triangles that are created are both the same. In other words, you get only one triangle.

Hence, you can create a congruence condition based on SSA with a right angle.

### 1.22: RHS (Right Angle-Hyp-Side) / Hyp-Leg

If the hypotenuse and one leg of a right-angled triangle are congruent to the hypotenuse and a leg of another right-angled triangle, then the two triangles are congruent.



### Example 1.23

On checking congruence for  $\triangle ABC$ :  $\angle ABC = 90^\circ$ ,  $AB = 3$ ,  $BC = 4$  and  $\triangle PQR$ :  $\angle PQR = 90^\circ$ ,  $PQ = 3$ ,  $QR = 4$ , a student says

$$\begin{aligned}\angle ABC &= 90^\circ, AB = 3, BC = 4 \\ \angle PQR &= 90^\circ, PQ = 3, QR = 4 \\ \therefore \triangle ABC &\cong \triangle PQR \text{ (RHS congruence)}\end{aligned}$$

Is the above solution correct or incorrect? Justify.

In the given solution, the hypotenuse has not been shown to be congruent, and hence *RHS* congruence is incorrectly used.

To use *RHS* congruence, we can first calculate, using the Pythagorean Theorem:

$$\begin{aligned}AC &= \sqrt{3^2 + 4^2} = \sqrt{25} = 5 \\ PR &= \sqrt{3^2 + 4^2} = \sqrt{25} = 5\end{aligned}$$

And then establish:


$$\begin{aligned}\angle ABC &= 90^\circ, AB = 3, AC = 5 \\ \angle PQR &= 90^\circ, PQ = 3, PR = 5 \\ \therefore \triangle ABC &\cong \triangle PQR \text{ (RHS congruence)}\end{aligned}$$

### Example 1.24

In the diagram alongside, triangles  $CDB$  and  $GFE$  are right angled at  $D$  and  $F$  respectively, and

$$\begin{aligned}FB &= GD = 1 \\ CB &= GE = \sqrt{2}\end{aligned}$$

Show that  $\triangle CDB$  is congruent to  $\triangle GFE$ .



$$\begin{aligned}GF &= GD + DF = 1 + DF \\ BD &= BF + DF = 1 + DF = GF\end{aligned}$$

$\triangle APC \cong \triangle BPC$  by *RHS*:


*Right Angle*:  $\angle CDB = \angle EFG = 90^\circ$  (Given)

*Hyp*:  $GE = CB$  (Given)

*Leg*:  $GF = BD$  (Shown above)

### Example 1.25

Show that  $CD$  is equal to  $DB$  in the diagram alongside.





$\triangle ADC \cong \triangle ADB$  by RHS because

$$\angle ADC = \angle ADB = 90^\circ \text{ (Right Angle) (Given)}$$

$$AC = AB \text{ (Hyp)}$$


$$AD = AD \text{ (Leg) (Reflexive Property)}$$

Hence,

$$CD = DB \text{ (CPCT)}$$

### Example 1.26

In the diagram, the lengths of AD and DB are equal. Prove that CA is congruent to CB.



$\triangle CDA \cong \triangle CDB$  by SAS

$$\angle CDA = \angle CDB = 90^\circ \text{ (Right Angle) (Given)}$$

$$AD = DB \text{ (Given)}$$

$$CD = CD \text{ (Reflexive Property)}$$

Hence:

$$CA = CB \text{ (CPCT)}$$

## C. Altitudes

### Example 1.27

In  $\triangle ABC$ , the altitude from C intersects AB at P, and the altitude from B intersects AC at Q. If PB and QC are congruent, then prove that  $\triangle ABC$  is isosceles.

$\triangle BPC \cong \triangle CQB$  by RHS because:

$$\angle BPC = \angle CQB = 90^\circ \text{ (Right Angle) (Altitudes are } \perp \text{ to sides)}$$

$$BC = BC \text{ (Hyp) (Reflexive Property)}$$


$$PB = QC \text{ (Leg) (Given)}$$

Hence:

$$\angle PBC = \angle QCB \text{ (CPCT)}$$

Hence,  $\triangle ABC$  is isosceles.

SAA, SAS, RHS, SSS



### Example 1.28

If two altitudes in a triangle are equal, they are drawn from equal angles.

$\triangle ABX \cong \triangle CYB$  by RHS:


$\angle BYC = \angle BXC = 90^\circ$  (Definition of Altitude)

$BX \cong CY$  (Leg) (Given)

$BC = BC$  (Hyp) (Reflexive Property)

By CPCTC:

$$\angle YBC = \angle XCB$$



## D. Perpendicular Bisector

### 1.29: Perpendicular Bisector of a Line Segment

Line segment  $CP$  is the perpendicular bisector of line segment  $AB$  if and only if:

- $P$  bisects  $AB$
- $CP \perp AB$

### Example 1.30

$C$  is a point equidistant from the endpoints of line segment  $AB$ . Show that it lies on the perpendicular bisector of  $AB$ .

Construct line segment  $CP$  perpendicular to line segment  $AB$ .

$\triangle APC \cong \triangle BPC$  by RHS:

Right Angle:  $\angle APC = \angle BPC = 90^\circ$  (Definition of Perpendicular)

Hyp:  $CA = CB$  (Given)


Leg:  $PC = PC$  (Reflexive Property)

By CPCTC:

$$PA = PB \Rightarrow P \text{ bisects } AB$$

Hence,  $C$  lies on the perpendicular bisector of  $AB$  because

$$CP \perp AB, P \text{ bisects } AB$$



## 1.3 SAS Theorem

### A. Basics

#### 1.31: SAS (Side-Angle-Side)

If two sides and the *included angle* of one triangle are congruent to two sides and the included angle of another triangle,  
then the two triangles are congruent.

- The angle must be included. Included means it is between the two sides.
- If the angle is not included then, we get ASS, which is not a congruence theorem.

### Example 1.32

In  $\triangle ABC$ ,  $AB = AC$ . Line segment  $AD$  is the angle bisector of  $\angle BAC$ . Prove that  $AD$  is the median to  $BC$ , and the altitude to  $BC$ .


(Do not assume any properties of isosceles triangles other than: an Isosceles triangle is a triangle with at least two congruent sides).

$\triangle ADB \cong \triangle ADC$  by SAS:

$$\begin{aligned} AD &= AD \text{ (Side) (Reflexive Property)} \\ \angle BAD &= \angle CAD \text{ (Angle) (Definition of Angle Bisector)} \\ AB &= AC \text{ (Side) (Given)} \end{aligned}$$

Show that  $AD$  is the median:

$$\begin{aligned} BD &= DC \text{ (CPCT)} \\ AD &\text{ is Median } (BD = DC) \end{aligned}$$



Show that that  $AD$  is the altitude:

$$\angle ADB = \angle ADC \text{ (CPCT)}$$

By angles in a linear pair:

$$\begin{aligned} \angle ADB + \angle ADC &= 180 \\ \angle ADB + \angle ADB &= 180 \\ \angle ADB &= \angle ADC = 90 \Rightarrow AD \text{ is altitude} \end{aligned}$$

## 1.4 ASA/SAA Theorem

### A. Basics

#### 1.33: ASA / SAA (Angle-Side-Angle / Side-Angle-Angle)

A side and two angles of a triangle are congruent

Note: Both conditions on the left are the same, since if two angles of a triangle are congruent, the third angle is also congruent.

#### Example 1.34


If a triangle is isosceles, then the altitudes drawn to the congruent sides are also congruent.

$\triangle BYC \cong \triangle BXC$  by ASA:

$$\begin{aligned} \angle BYC &= \angle BXC = 90^\circ \text{ (Angle) (Given)} \\ BC &= BC \text{ (Side) (Reflexive Property)} \\ \angle XBC &= \angle YCB \text{ (Angle) (Given)} \end{aligned}$$

Hence,

$$CX \cong BY \text{ (CPCT)}$$




### 1.35: ASS

- Not a congruence theorem
- **Ambiguous Case – Be careful**

Construct  $\triangle ABC$  with

$$\angle A = 26^\circ, AB = 3 \text{ units}, BC = 2 \text{ units}$$

$$\angle P = 26^\circ, PQ = 3 \text{ units}, QR = 2 \text{ units}$$



## B. Isosceles Triangles

### 1.36: Special Lines are Identical

In an Isosceles Triangle, the median, the angle bisector, and the altitude drawn from a vertex are all the same.

- We prove this in three different ways below.
- This property does not hold true in general for all triangles. In fact, it is not true for scalene triangles.

#### Example 1.37

Prove, using *RHS Congruence*, that, in an Isosceles Triangle, the median to the base, the angle bisector to the base, and the altitude to the base are the same.

In Isosceles  $\triangle ABC$ , with  $AB = AC$ , draw the altitude from point  $A$ , intersecting base  $BC$  at  $D$ . (Construction)

$\triangle ADB \cong \triangle ADC$  by *RHS*

$$AD = AD \text{ (Reflexive Property)}$$


$$AB = AC \text{ (Definition of Isosceles Triangle)}$$

$$\angle ADB = \angle ADC = 90^\circ \text{ (Definition of Altitude)}$$

Hence, by CPCT:

$$BD = DC \Rightarrow AD \text{ is Median}$$

$$\angle BAD = \angle CAD \Rightarrow AD \text{ is angle bisector}$$



#### Example 1.38

Prove, using *SSS Congruence*, that, in an Isosceles Triangle, the median to the base, the angle bisector to the base, and the altitude to the base are the same.

(Don't assume any properties of an isosceles triangle other than the definition, which for this question is: An Isosceles triangle is a triangle with at least two congruent sides).

Construction:


In Isosceles  $\triangle ABC$ , with  $AB = AC$ , draw the median from point  $A$ , intersecting base  $BC$  at  $D$ .

$\triangle ADB \cong \triangle ADC$  by *SSS*:

$$AD = AD \text{ (Reflexive Property)}$$

$$AB = AC \text{ (Definition of Isosceles Triangle)}$$

$$BD = DC \text{ (Definition of Median)}$$




In  $\triangle ADB$  and  $\triangle ADC$ , by CPCT

$$\begin{aligned}\angle BAD &= \angle CAD \Rightarrow AD \text{ is angle bisector} \\ \angle ADB &= \angle ADC \Rightarrow \angle ADB + \angle ADC = 180 \Rightarrow \angle ADB = \angle ADC = 90 \Rightarrow AD \text{ is altitude}\end{aligned}$$

### Example 1.39

Prove that the base angles of an isosceles triangle are congruent.  
(Don't assume any properties of an isosceles triangle other than the definition, which for this question is:

An Isosceles triangle is a triangle with at least two congruent sides).



Take any of the three proofs above, and show that

$$\triangle ADB \cong \triangle ADC$$

Then, since corresponding parts of congruent triangles are equal:

$$\angle B = \angle C \Rightarrow \text{Base angles are equal}$$

## 1.40: Converse: Special Lines are Identical

If, in a triangle:

- The altitude is the same as the angle bisector, then the triangle is Isosceles
- The altitude is the same as the median, then the triangle is Isosceles
- The median is the same as the angle bisector, then the triangle may or may not be Isosceles

### Example 1.41

Prove that a triangle where the altitude is the same as the angle bisector is an Isosceles Triangle.


In  $\triangle ABC$  let  $AD$  be the common altitude and angle bisector.

$\triangle ABD \cong \triangle ACD$  by ASA:

$$\begin{aligned}\angle BDA &= \angle CDA = 90^\circ \text{ (Angle) (Definition of Altitude)} \\ AD &\cong AD \text{ (Side) (Reflexive Property)} \\ \angle BAD &\cong \angle CAD \text{ (Angle) (Definition of Angle Bisector)}\end{aligned}$$

By CPCT:

$$AB \cong AC$$



### Example 1.42

Prove that a triangle where the altitude is the same as the median is an Isosceles Triangle.


In  $\triangle ABC$  let  $AD$  be the common altitude and median.

$\triangle ABD \cong \triangle ACD$  by SAS:

$$\begin{aligned}AD &\cong AD \text{ (Side) (Reflexive Property)} \\ \angle BDA &= \angle CDA = 90^\circ \text{ (Angle) (Definition of Altitude)} \\ BD &= DC \text{ (Side) (Definition of Median)}\end{aligned}$$

By CPCT:

$$AB \cong AC$$



### Example 1.43

- Draw a counterexample where the median is also the angle bisector in a triangle, but the triangle is not isosceles.
- If you attempt to set up a congruence, which “congruence theorem” do you get? What is the pitfall here?

In  $\triangle ABC$  let  $AD$  be the common median and angle bisector:

$$\begin{aligned}BD &= DC \text{ (Side) (Definition of Median)} \\AD &\cong AD \text{ (Side) (Reflexive Property)} \\ \angle BAD &= \angle CAD = 90^\circ \text{ (Angle) (Definition of Angle Bisectore)}\end{aligned}$$

What we get is

$$SSA \Rightarrow \text{Not a congruence theorem}$$

### 1.44: Altitudes in an isosceles triangle

In an isosceles triangle, the altitudes drawn to equal sides are equal.

The converse is also true: if the altitudes drawn to two sides are equal, then the triangle is isosceles.

### Example 1.45

Prove that if the perpendiculars to two sides of a triangle from the midpoint of the third side are congruent, then the triangle is isosceles.

Draw  $\triangle ABC$ . Let  $D$  be the midpoint of side  $BC$ . Then:

$$DB = DC \text{ (Hyp)}$$

Draw  $DE \perp AB$  and  $DF \perp AC$ .

$$\angle DEB = \angle DFC = 90^\circ \text{ (Right Angle)}$$

From the question, we are given that:


$$DE = DF \text{ (Side)}$$

From the above three statements:

$$\triangle DEB \cong \triangle DFC \text{ (RHS Congruence)}$$

$$\angle EBD = \angle FCD \text{ (CPCT)}$$

Hence, since its base angles are congruent,  $\triangle ABC$  is isosceles.




### 1.46: Medians in an isosceles triangle

In an isosceles triangle, the medians drawn to equal sides are equal.

The converse is also true: if the medians drawn to two sides are equal, then the triangle is isosceles.

### Example 1.47

Show that the medians drawn to equal sides in a triangle are equal.



Prove the converse of the above statement: If two medians in a triangle are equal, they are drawn from equal angles

### 1.48: Angle Bisectors in an isosceles triangle

In an isosceles triangle, the angle bisectors drawn to equal sides are equal.

The converse is also true: if the angle bisectors drawn to two sides are equal, then the triangle is isosceles.

### Example 1.49

Show that the angle bisectors drawn to equal sides in a triangle are equal.

### Example 1.50

Show that if the angle bisectors drawn to two sides in a triangle are equal, then the triangle is isosceles.

Note: the proof here is based on inequalities, not on congruence.

## C. Perpendicular Bisectors

### Example 1.51

Prove

- If a point is on the perpendicular bisector of a line segment, then it is equidistant from the ends of the line segment.
- If a point is equidistant from the ends of a line segment, then it is on the perpendicular bisector of the line segment.

#### Part A

On line segment  $AB$  let  $M$  be the midpoint.


Construct the perpendicular bisector to line segment  $AB$ . Note that the bisector passes through  $M$ .

Let  $P$  be a point on the perpendicular bisector.

Then, in  $\triangle PAM$  and  $\triangle PBM$

$$\begin{aligned} AM &= BM \text{ (} M \text{ is midpoint)} \\ \angle PMA &= \angle PMB = 90^\circ \text{ (Def of } \perp) \\ PM &= PM \text{ (Reflexive property)} \\ \triangle PAM &\cong \triangle PBM \text{ (SAS Theorem)} \\ PA &= PB \text{ (CPCTC)} \end{aligned}$$

$P$  is equidistant from  $A$  and  $B$ . (Def of distance and  $PA = PB$ )




### Part B

Let  $P$  be equidistant from  $A$  and  $B$ .

Draw the perpendicular from  $P$  to  $AB$ , intersecting  $AB$  at  $M$ .

Then, in  $\triangle PAM$  and  $\triangle PBM$

$PA = PB$  (Def of Distance)  
 $PM = PM$  (Reflexive Property)  
 $\angle PMA = \angle PMB = 90^\circ$  (Def of  $\perp$ )  
 $\triangle PMA \cong \triangle PMB$  (RHS Theorem)  
 $AM = BM$  (CPCTC)  
 $P$  lies on perpendicular bisector



## 1.5 Applications

### A. Parts of Congruent Triangles

#### 1.52: More corresponding parts

The heights, angle bisectors, medians, and other parts of two congruent triangles are also congruent.

#### Example 1.53

Prove that if two triangles are congruent, their medians are also congruent.

Let  $\triangle ABC \cong \triangle DEF$ .

Draw medians  $AG$  and  $DH$


$\triangle ABG \cong \triangle DEH$  by SAS:

$$AB = DE \text{ (CPCT)}$$

$$\angle ABG = \angle DEH \text{ (CPCT)}$$

$$BC = EF \text{ (CPCT)}$$

$$\frac{BC}{2} = BG = EH = \frac{EF}{2} \text{ (Def. of Median)}$$



Hence:

*Angle Bisectors  $AG$  and  $DH$  are congruent (CPCT)*

Similarly, the other two medians can also be proved congruent.

#### Example 1.54

Prove that if two triangles are congruent, their angle bisectors are also congruent.



Let  $\triangle ABC \cong \triangle DEF$ .

Draw angle bisectors  $AG$  and  $DH$


$\triangle ABG \cong \triangle DEH$  by SAA:

$$\angle ABG = \angle DEH \text{ (CPCT)}$$

$$AB = DE \text{ (CPCT)}$$

$$\angle BAC = \angle EDF \text{ (CPCT)}$$

$$\frac{\angle BAC}{2} = \angle BAG = \angle EDH = \frac{\angle EDF}{2}$$



Hence:

*Angle Bisectors  $AG$  and  $DH$  are congruent (CPCT)*

Similarly, the other two angle bisectors can also be proved congruent.

### Example 1.55

Prove that if two triangles are congruent, their heights are also the same.

Let  $\triangle ABC \cong \triangle XYZ$ .

Draw altitudes  $AD$  and  $XP$ .


#### Case I: The altitude lies inside the triangle

$\triangle ABD \cong \triangle XYP$  By SAA

$$AB \cong XY \text{ (CPCT)}$$

$$\angle ABD \cong \angle XYP \text{ (CPCT)}$$

$$\angle ADB = \angle XPY = 90^\circ \text{ (AD} \perp \text{BC, XP} \perp \text{YZ)}$$




#### Case II: The altitude lies outside the triangle

$\triangle ACD \cong \triangle XZP$  By SAA

$$AC \cong XZ \text{ (CPCT)}$$

$$\angle ACD \cong \angle XZP \text{ (CPCT)}$$

$$\angle ADC = \angle XPZ = 90^\circ \text{ (AD} \perp \text{BC, XP} \perp \text{YZ)}$$



In both the cases above, by CPCT:

$$AD \cong XP \Rightarrow \text{Altitudes are congruent}$$

Since there is nothing special about the altitude from  $A$ , we can prove similarly that the altitudes from  $B$  and  $C$  are also congruent.

## B. Midpoint Theorem

### 1.56: Midpoint Theorem

- In a triangle, the line joining the midpoints of two sides is parallel to the third side, and half of the third side.

In  $\triangle ABC$ , let  $X$  and  $Y$  be the midpoints of  $AB$  and  $AC$ , respectively. Draw  $XY$  and produce  $XY$  to  $Z$  such that  $XY = YZ$ .

In  $\triangle AYX$  and  $\triangle CYZ$ :

$$\angle AYX = \angle CYZ \text{ (Vertically opposite angles)}$$

$$XY = YZ \text{ (Construction)}$$

$$AY = YC \text{ (Y is midpoint of AC)}$$

$$\therefore \triangle AYX \cong \triangle CYZ \text{ (SAS Theorem)}$$

Consider  $AC$  as a transversal of  $BA$  and  $CZ$ :

$$\angle XAY \text{ and } \angle ZCY \text{ are alternate interior angles}$$

$$\angle XAY \cong \angle ZCY \text{ (CPCTC in } \triangle AYX \text{ and } \triangle CYZ)$$

$$\therefore AB \parallel CZ$$

$$\therefore BX \parallel CZ$$

Also:

$$BX \cong AX \text{ (X is midpoint of AB)}$$

$$AX \cong CZ \text{ (CPCTC in } \triangle AYX \text{ and } \triangle CYZ)$$

$$\therefore BX \cong CZ$$


Hence,

$$BX \parallel CZ, BX \cong CZ \Rightarrow XZCB \text{ is a parallelogram}$$

Since  $XZCB$  is a parallelogram:

$$XZ \parallel BC \Rightarrow \text{Parallel condition}$$

$$XZ = BC \Rightarrow XY = \frac{1}{2}XZ = \frac{1}{2}BC \Rightarrow \text{Half Condition}$$



## C. Right Triangles

### 1.57: Median to Hypotenuse

The median to the hypotenuse of a right-angled triangle is half the length of the hypotenuse.

#### Using Congruence

In right  $\triangle ABC$ , draw median  $BD$ , and hence

$$AD = DC$$

Draw  $DE$  such that  $E$  is the midpoint of  $BC$ .

Then, in  $\triangle BDE$  and  $\triangle CDE$ :


$$DE = DE \text{ (Common Side)}$$

$$BE = EC \text{ (Construction)}$$

$$\angle DEC = \angle DEB = 90^\circ \text{ (Midpoint Theorem)}$$

$$\triangle BDE \cong \triangle CDE \text{ (SAS)}$$

$$BD \cong CD \text{ (CPCTC)}$$



### Using Isosceles Triangles

In right  $\triangle ABC$ , draw  $BD$  such that

$$\angle DBC = \angle DCB = \theta$$

This means that  $\triangle DBC$  is isosceles, and hence

$$\underline{BD = DC}$$

*Equation I*

By sum of angles of a triangle,

$$\angle DAB = 180 - 90 - \theta = 90 - \theta$$

Also, since  $\angle ABC = 90^\circ$ :

$$\angle DAB = 90 - \theta$$

Hence,  $\triangle ADB$  is isosceles and

$$\underline{AD = BD}$$


*Equation II*

Combine Equations I and II to get:

$$AD = BD = DC \Rightarrow AD = DC$$

Hence,  $BD$  is the median, and since the median is unique, we have also shown that

$$BD = DC = \frac{1}{2}AC$$



## 1.6 Further Topics

### 58 Examples