# +44 7719 773067 adam.morris@cern.ch

## **Adam Morris**

### Curriculum Vitae

### Personal statement

I am a PhD student with the LHCb group at the University of Edinburgh. My research experience is in studies of four-body charmless hadronic decays of B mesons.

## Education

#### PhD in experimental particle physics

2013 —

University of Edinburgh

Thesis title Measurements of charmless  $B_s$  meson decays at LHCb

Supervisors Matthew Needham and Franz Muheim

**MSci in physics** 2009 — 2013

University of Birmingham

Thesis title A possible lepton flavour violation search at NA62 Supervisors Cristina Lazzeroni and Evqueni Goudzovski

Classification 2:1

## Physics analysis

### Amplitude analysis of $B^0_s \to \phi K^+ K^-$ decays

2015 —

An amplitude fit to  $B_s^0 \to \phi K^+ K^-$  events with  $m(K^+ K^-) < 1.8$  GeV. The decay  $B_s^0 \to \phi f_2'(1525)$  is observed for the first time, and measurements are made of its branching fraction and longitudinal polarisation fraction.

This is my primary thesis work, and I was responsible for performing the analysis, from selection, modelling the acceptance, signal and background, fitting the data and calculating results and systematic uncertainties. The amplitude fit was performed using Edinburgh's RapidFit fitting framework, which I modified to make more suited to the task. The PDF was adapted from the  $B^0 \to \psi(2S)K^+\pi^-$  analysis, and extended to become extensively runtime-configurable, allowing many versions of the fit to run simultaneously on a HPC cluster without the need to re-compile.

## Measurement of the $B^0_s o \phi \phi$ branching fraction and search for $B^0 o \phi \phi$ [1]

2014 — 2015

The  $B^0_s \to \phi \phi$  branching fraction was measured using  $B^0 \to \phi K^{*0}$  for normalisation, resulting in a factor of five reduction in the statistical uncertainty compared to the previous best result. In addition, a search was performed for the suppressed  $B^0 \to \phi \phi$  decay, and an upper limit placed on its branching fraction, which is a factor of seven improvement compared to the previous best result.

This analysis forms part of my thesis work, and I was responsible for the mass fits and most of the calculations of efficiencies and systematics, as well as setting the limit on the  $B^0$  mode using the  $CL_s$  method.

## Measurement of CP violation in $B_s^0 o \phi \phi$ decays [2]

2013 - 2014

A time-dependent angular fit performed to  $B_s^0 \to \phi \phi$  events to extract the weak phase  $\phi_s$ , which is expected to be close to zero due to cancellation between the leading-order mixing and decay diagrams.

I was responsible for parametrising the peaking backgrounds and performing fits to the  $\phi\phi$  invariant mass to obtain event yields.

## Hardware and operations

LHCb control room shifts (4× shift leader & 11× data manager)	2015 — 2016
RICH piquet shifts (3 weeks)	2015 - 2016
Ion feedback monitoring of the RICH HPDs	2015 - 2016
RICH upgrade testbeam [3]	2014 - 2016
Performance of MaPMTs in magnetic fields for the RICH upgrade [4]	2013 - 2014

## Schools, workshops and conferences

**Exotic hadron spectroscopy workshop** 

University of Edinburgh, September 2016

Heavy Quarks and Leptons (HQL) conference

Virginia Tech, May 2016

Talk: Charmless b-meson and b-baryon decays at LHCb [5]

**LHCC Poster Session** 

CERN, March 2016

Poster: Measurement of the  $B_s \to \phi \phi$  branching fraction and angular analysis of  $B_s \to \phi \pi^+ \pi^-$  at LHCb

UK HEP forum: "Anomalies and Deviations" workshop

STFC, Abingdon, November 2015

Poster: Measurement of the  $B_s \to \phi \phi$  branching fraction and search for  $B^0 \to \phi \phi$  at LHCb

**European School of High-Energy Physics** 

Bansko, Bulgaria, September 2015

Young Experimentalists and Theorists Institute (YETI) workshop

Durham University, January 2015

STFC HEP summer school

University of Warwick, September 2014

B-physics at frontier machines (Beauty) conference

University of Edinburgh, July 2014

Poster: Measurement of CP violation in  $B_s^0 o \phi \phi$  decays [6]

#### **Public outreach**

Participant in an 'Ask Me Anything' session organised by CERN on *reddit.com*May 2015

Participant in a *Speed Science* event at the *Glasgow Science Festival*Demonstrator at the *Discovering the Higgs Boson* exibit at the *Big Bang Fair*, Birmingham NEC

March 2014

## **Skills**

- HEP data analysis techniques: multivariate selection, maximum-likelihood fits, amplitude analysis and limit-setting
- Proficiency with C++, Python and shell scripting
- Familiarity with ROOT (incl. TMVA, RooFit and RooStats), GSL, Boost and the LHCb software
- Experience using parallel computing resources, such as the Edinburgh and CERN batch services and the LHC Grid
- Frequent user of version control systems (Git and SVN) and experience with collaborative software development

#### **Publications**

#### **Papers**

- [1] Measurement of the  $B_s^0 \to \phi \phi$  branching fraction and search for the decay  $B^0 \to \phi \phi$ , LHCb collaboration, JHEP **10** (2015) 053
- [2] Measurement of CP violation in  $B_s^0 \to \phi \phi$  decays, LHCb collaboration, Phys. Rev. **D90**, (2014) 052011
- [3] Test of the photon detection system for the LHCb RICH Upgrade in a charged particle beam, M. K. Baszscyk et al, LHCb-PUB-2016-019, to be submitted to JINST

### **Proceedings**

- [4] Characterisation and magnetic field properties of multianode photomultiplier tubes,
  - S. Eisenhart et al, on behalf of the LHCb RICH collaboration, Nucl. Instrum. Meth. A766 (2014) 167-170
- [5] Charmless b-meson and b-baryon decays at LHCb,
  - A. Morris, on behalf of the LHCb collaboration, PoS (HQL 2016) 059
- [6] Measurement of CP violation in  $B^0_s \to \phi \phi$  decays,
  - A. Morris, on behalf of the LHCb collaboration, PoS (Beauty 2014) 064