

Lecture 1

1 Introduction

Definition 1 (Partial Differential Equations (PDEs)) ... □

However we only concentrate on those questions that naturally occur in various applications, such as physics and other sciences, engineering and economics.

Now let's give some example of PDEs.

Example 1

$$\Delta u := \sum_{i=1}^d u_{x^i x^i} = 0 \quad (\text{Laplace Equation})$$

$$\Delta u = f \quad (\text{Poisson Equation})$$

$$u_t = \Delta u \quad (\text{Heat Equation})$$

$$u_{tt} = \Delta u \quad (\text{Wave Equation})$$

$$(1 + u_y^2)u_{xx} - 2u_x u_y u_{xy} + (1 + u_x^2)u_{yy} = 0 \quad (\text{Minimal Surface Equation})$$

$$\begin{cases} \operatorname{div} B = 0 \\ B_t - \operatorname{curl} B = 0 \\ \operatorname{div} E = 4\pi \rho \\ E_t - \operatorname{curl} E = -4\pi j \end{cases} \quad (\text{Maxwell Equations})$$

$$\det(u_{x^i x^j}) = f \quad (\text{Monge-Ampere Equation})$$

$$u_t - 6uu_x + u_{xxx} = 0 \quad (\text{Korteweg-de Vries Equation})$$

1.1 Classification

By of linear of nonlinear

- Linear equations
- Non-linear equations
 - Quasilinear equations
 - Semilinear equations

By order

- First order
- Second order
 - elliptic
 - parabolic
 - hyperbolic
 - etc.
- Higher order

Exercise 1 *Classify all the equations above.*

Exercise 2 *Give more equations and try to classify them according to the standard given above.*

2 What will we learn from this textbook

Question 1 How to find a solution of elliptic PDES.

In general we have the following five methods:

- 0 Write down an *explicit* formula for the solution in terms of the given data.
具有极大的局限性。仅仅对少数方程有效。
- I Solve a sequence of auxiliary problems that *approximate* the given one and show ...
偏微分方程可以看作一个无穷维函数空间上的函数方程，从而使用有限维空间逼近。
- II Start anywhere, with the required constraints satisfied, and let things *flow* toward a solution.
解是一个扩散过程的渐进平衡态。
- III Solve an *optimization* problem, and ...
变分问题的极值问题，既是偏微分方程的一个来源，也是种解方程的方法。
- IV *Connect* what you want to know to what you already know, that is the *continuity method*.
由已知问题连续的过渡到未知问题。

Question 2 Regularity of Solutions and Uniqueness of Solutions.

generalized solutions vs. smooth solutions

The understanding of the above summary will be deepened after our study.

In any case, it's usually not very efficient to read a mathematical textbook linearly, and the reader should rather try first to grasp the central statements.

可以参考 MIT 公开课程的教学计划。地址见：haoyun.github.com/academic/seminars/2012fall-pde/

3 Harmonic Functions

Definition 2 (Harmonic functions) A function u on a domain Ω that satisfies the Laplace Equation:

$$\Delta u = 0$$

is called a harmonic function on Ω . □

Remark 1 All harmonic functions constitute a linear space. Simple verification will prove this. A higher view of point is that all harmonic functions is the kernel of Laplace operator, which is a linear operator. □

Example 2 Real and imaginary part of a complex analytic function. (As an exercise, give some examples, and note that the derivative of an analytic function is also analytic, thus we can find even more desired functions.) This is a consequence of Cauchy-Riemann Equation. □

Remark 2 This view of point enable us to solve some problems on harmonic functions using the method and theories developped in Complex Analysis. We will give some example later. \square

Lemma 1 *Laplace operator is rotation-invariant.*

PROOF Note that $\Delta u(x) = \text{tr}(H[u(x)])$, where $H[u(x)]$ is the Hessein matrix of $u(x)$.

$$\Delta_x u(A(x - y) + y) = \text{tr}(A^T H[u(A(x - y) + y)]A) \quad \blacksquare$$

Another proof? \square

Taking into consideration that Δ is rotation-invariant, we may suppose that the solution of Δu is also rotation-invariant, or radial, i.e.,

$$u(x) = \varphi(r), \text{ where } r = |x - y| \text{ with fixed } y$$

Then $\Delta u = 0$ can be written as

$$\varphi''(r) + \frac{d-1}{r}\varphi'(r) = 0$$

Solve the second order ordinary differential equation we obtain that

$$\varphi'(r) = Cr^{1-d}$$

and then

$$\varphi(r) = \begin{cases} C \log r & d = 2 \\ \frac{C}{2-d} r^{2-d} & d > 2 \end{cases}$$

Where C is a constant.

Definition 3 (Fundamental Solution) ... \square

Lemma 2 (Divergence Theorem) ... \square

Lemma 3 (Green's Formula) ... \square

Lemma 4 (Lebesgue Dominated Convergence Theorem) ... \square

Lemma 5 (Lebesgue Theorem) ... \square

Theorem 1 (Green Representation Formula)

$$u(y) = \int_{\partial\Omega} \left(u \frac{\partial \Gamma}{\partial \nu} - \Gamma \frac{\partial u}{\partial \nu} \right) + \int_{\Omega} \Gamma \Delta u \quad \square$$

Remark 3 From Green Representation Formula we know that u is determined by its normal derivatives on $\partial\Omega$, provided Δu is given, in particular u is harmonic. \square

Remark 4 Conversely, we cannot, however, obtain a harmonic function using this formula by giving an arbitrary $\frac{\partial u}{\partial \nu}|_{\partial\Omega}$, because it must satisfy some constraints, say, at least,

$$\int_{\partial\Omega} \frac{\partial u}{\partial \nu} = \int_{\Omega} \Delta u = 0 \quad \square$$

We now turn to another method.

Definition 4 (Green Function) ... □

Remark 5 The idea of why we introduce Green function is unknown from this textbook. □

We suppose there do exist a Green function for Ω , then by Green Representation Formula, we have

$$u = \int_{\partial\Omega} u \frac{\partial G}{\partial \nu} + \int_{\Omega} G \Delta u \quad (1)$$

Remark 6 From (1) we know that u is determined by its value on the boundary, provided Δu is given, in particular u is harmonic. □

Remark 7 Under what condition can we reconstruct u by given $t|_{\partial\Omega}$ and Δu is the next question.

...

□

Next, we turn to the Dirichlet Problem on a ball.

$$G(x, y) := \begin{cases} \Gamma(|x - y|) - \Gamma(\frac{|y|}{R}|x - \bar{y}|) & y \neq 0 \\ \Gamma(|x|) - \Gamma(R) & y = 0 \end{cases} \quad (2)$$

First of all, $G - \Gamma$ is harmonic

Secondly, ... $G(R) = 0$

Therefore $G(x, y)$ is a Green function of $B(0, R)$.

Next we calculate $\frac{\partial G}{\partial \nu}$, ..., we have

$$\frac{\partial G}{\partial \nu_x} = \dots = \frac{R^2 - |y|^2}{d\omega_d R} \frac{1}{|x - y|^d} \quad (3)$$

Then we can obtain the following theorem.

Theorem 2 (Poisson Representation Formula) ...

PROOF the fact that Δ commutes with \int leads to u is harmonic about y .

Next we show the continuity ...

■

Corollary 1 (smoothness) u is real analytic on Ω .

PROOF ...

■

Corollary 2 (uniqueness) ...

PROOF ...

■