

Forecasting ionospheric Total Electron Content maps with deep neural networks

Noëlie Cherrier, Thibaut Castaings, **Alexandre Boulch**

DeLTA : Deep Learning for Aerospace Applications

Electromagnetism

Remote sensing

Robotics

Optics



Materials and structures

Fluid mechanics

Table of contents

TEC

Approach for TEC prediction


Experiments

Conclusion and future work

Ionosphere

Ionosphere

Highly ionized region in upper atmosphere.



Ionosphere state

Influence




Affects trans-ionospheric
radiowaves



GPS / space
based telecoms

TEC Map



Total Electron Content (TEC) measures ionospheric activity.

TEC = Integration of electron density along a $1m^2$ sect. tube between GNSS station and GNSS satellite


Table of contents

TEC

Approach for TEC prediction

Experiments

Conclusion and future work



Code TEC data


aiuws.unibe.ch/ionosphere/
based 200 stations
1 TEC map every 2 hours since
2003

- ▶ 72×80
- ▶ Resolution : $5^\circ \times 2.5^\circ$

Approach

Preprocessing


Heliocentric coordinates : remove rotation effect.



Approach

Preprocessing


Heliocentric coordinates : remove rotation effect.




Approach

Preprocessing


Heliocentric coordinates : remove rotation effect.



Network architecture



Encoder - Decoder architecture¹



-
1. Work presented at ICONIP : *Deep sequence-to-sequence neural networks for ionospheric activity map prediction* [1]

Recurrent U-net




Table of contents


TEC

Approach for TEC prediction

Experiments

Conclusion and future work

Comparison with Encoder-Decoder method



Prediction difference between [1] and Rec-Unet.

Quantitative results

Whole test set

Method	RMS 48h	First 24h	Last 24h
Priodic	2.74	2.88	2.53
ICONIP	2.65	2.65	2.65
Ugru	2.66	2.46	2.85

First half of test set

Method	RMS 48h	First 24h	Last 24h
Priodic	2.88	2.87	2.89
ICONIP	2.75	2.74	2.76
Ugru	2.60	2.46	2.74

Note : mean over 6 runs, numbers updated compared to paper. Different test set.

Comparison with other approaches

	Reference	RMS (ref)	RMS (best run)
[2]	Chunli D., Jinsong P.	1.45	2.1
[3]	Huang, Z., Yuan, H.	≤ 2	1.53
[4]	Niu, R. <i>et al.</i>	3.1	0.73

2

2. Erratum : in paper numbers from [1]. Replaced at aboulch.github.

io

Table of contents

TEC

Approach for TEC prediction

Experiments

Conclusion and future work

Conclusion

Our method

- ▶ Global TEC prediction
- ▶ Recurrent Unet

Perspectives

- ▶ Improve prediction from 24h to 48h
- ▶ Improve convergence (may diverge)
- ▶ Reduce time dependency to training set (train on more data)
- ▶ Involve other sources (e.g. sun imagery)

Thanks for your attention

Slides and updated paper at : aboulch.github.io

Implementation

- ▶ PyTorch framework
- ▶ Code to be released

References

-  Noëlie Cherrier, Thibaut Castaings, and Alexandre Boulch.
Deep sequence-to-sequence neural networks for ionospheric activity map prediction.
In *International Conference on Neural Information Processing*, pages 545–555. Springer, 2017.
-  D. Chunli and P. Jinsong.
Modeling and prediction of TEC in China region for satellite navigation.
In *2009 15th Asia-Pacific Conference on Communications*, pages 310–313, Oct 2009.
-  Z. Huang and H. Yuan.
Ionospheric single-station TEC short-term forecast using RBF neural network.
Radio Science, 49(4) :283–292, 2014.
-  R. Niu, C. Guo, Y. Zhang, L. He, and Y. Mao.
Study of ionospheric TEC short-term forecast model based on combination method.
In *2014 12th International Conference on Signal Processing (ICSP)*, pages 2426–2430, Oct 2014.