

# Class Field Theory

## Lectures by Frank Calegari

Typed by Andrea Bourque

Fall 2025

### Preface

There will be some gaps in explanation, either due to the lecturer's admission or my own lack of understanding. In particular, many "proofs" are sketches of proofs. Gaps due to my own misunderstanding will be indicated by three red question marks: ???. More generally, my own questions about the material will also be in red. Things like "**Question**" will be questions posed by the lecturer. Feel free to reach out to me with explanations.

### Contents

<b>1</b>	<b>Sep 29: Intro</b>	<b>2</b>
1.1	Prereqs . . . . .	2
1.2	Galois Theory . . . . .	2
<b>2</b>	<b>Oct 3 25</b>	<b>5</b>

# 1 Sep 29: Intro

## 1.1 Prereqs

The two more important prereqs are working knowledge of Galois theory and representation theory of finite groups. The less important prereqs are knowledge of the statements of core theorems in a first course of algebraic number theory, particularly Dirichlet's unit theorem and finiteness of class groups, as well as some homological algebra.

## 1.2 Galois Theory

Let  $L/K$  be a Galois extension with Galois group  $G$ . The **Galois correspondence** states that there is an inclusion-reversing bijection between fields  $L'$  with  $K \subseteq L' \subseteq L$  (also called **subextensions** of  $L/K$ ) and subgroups of  $G$ ; explicitly, a subgroup  $H \leq G$  gives the subextension  $L^H$  of fixed points of the  $H$ -action, and a subextension  $L'$  gives the subgroup  $\text{Gal}(L/L')$ .

The action of  $G$  on  $L$  is  $K$ -linear, meaning  $L$  has the structure of a  $K[G]$ -module. Also, the unit group  $\mathcal{O}_L^\times$  is stable under the  $G$ -action, so  $\mathcal{O}_L^\times$  has a  $\mathbb{Z}[G]$ -module structure. We aim to answer the following questions:

### Questions:

1. What is the structure of  $L$  as a  $K[G]$ -module?
2. What is the structure of  $\mathcal{O}_L^\times$  as a  $\mathbb{Z}[G]$ -module?
3. What is the structure of  $\mathcal{O}_L^\times \otimes_{\mathbb{Z}} \mathbb{C}$  as a  $\mathbb{C}[G]$ -module?

The first question has a nice answer:

**Theorem 1.1** (Normal Basis Theorem). *If  $L/K$  is a Galois extension with Galois group  $G$ , then  $L \cong K[G]$  as a  $K[G]$ -module. Equivalently, there exists some  $\theta \in L$ , called a **normal basis element**, such that the set  $\{g\theta \mid g \in G\}$  is a  $K$ -basis for  $L$ .*

**Example 1.1.** Suppose  $[L : K] = 2$  and  $\text{char}(K) \neq 2$ . Then  $L = K(\sqrt{D})$  for some square free  $D \in K$ . Any  $y \in L$  can be written in the form  $y = a + b\sqrt{D}$  for  $a, b \in K$ . Then  $\theta$  is a normal basis element iff  $y$  and  $\bar{y} = a - b\sqrt{D}$  span  $L$ , which happens iff  $ab \neq 0$ .

We will prove 1.1 in a bit. First, we showcase a certain proof technique.

**Theorem 1.2** (Cayley-Hamilton). *Let  $M \in M_n(R)$  be an  $n$  by  $n$  matrix with entries in a commutative ring  $R$ . Then  $\text{ch}_M(M) = 0$ , where  $\text{ch}_M(x) = \det(xI - M) \in R[x]$  is the characteristic polynomial of  $M$ .*


*Proof.* We will make several reductions and then pass to a lemma.

1. We can reduce to the case where  $R$  is finitely generated by simply taking the subring  $S$  of  $R$  generated by the entries of  $M$ . Notably,  $ch_M(x) \in S[x]$ , so there will be no issue in replacing  $R$  by  $S$ .
2. Using a choice of generators for  $R$ , we get a surjection  $\mathbb{Z}[x_1, \dots, x_d] \rightarrow R$ . We can lift  $M$  to a matrix  $M'$  over  $\mathbb{Z}[x_1, \dots, x_d]$  by making arbitrary choices for the preimages of the entries of  $M$ . The matrix  $ch_{M'}(M')$  will map down to  $ch_M(M)$ , so it suffices to prove the theorem when  $R = \mathbb{Z}[x_1, \dots, x_d]$ .
3. We can embed into  $\mathbb{Q}(x_1, \dots, x_d)$ , and better yet, an algebraic closure thereof.

To conclude the proof from here, we use the following lemma. 

**Lemma 1.1.** *Let  $K$  be an infinite field, and let  $P \in K[x_1, \dots, x_d]$ . Then TFAE:*

1.  $P \neq 0$ .
2. If  $L/K$  is any field extension, then  $P \in L[x_i]$  is not 0.
3. There exist elements  $a_i \in K$  such that  $P(a_i) \neq 0$ .
4. If  $L/K$ , there exist elements  $a_i \in L$  such that  $P(a_i) \neq 0$ .

*Proof.* It is clear that 1 is equivalent to 2, and that 3 implies 4 implies 1. Then it suffices to show that 1 implies 3. Assume  $P \neq 0$ . We induct on  $d$ . For  $d = 1$ ,  $P \in K[x]$  has only finitely many roots, while  $K$  itself is infinite, so there must be some  $a \in K$  with  $P(a) \neq 0$ . For  $d > 1$ , we can write  $P$  as a polynomial in one variable over the field  $L = K(x_1, \dots, x_{d-1})$ . Certainly  $L$  is infinite, so by the base case there is some  $a_d \in L$  with  $P(x_1, \dots, x_{d-1}, a_d) \neq 0$  in  $L$ . By clearing denominators, we obtain a nonzero polynomial  $Q \in K[x_1, \dots, x_{d-1}]$ , equal to  $P(y_1, \dots, y_d)$  for some  $y_1, \dots, y_d \in K[x_1, \dots, x_{d-1}]$ . By induction, there are some  $a_1, \dots, a_{d-1}$  such that  $Q(a_i) \neq 0$ . Using  $x_i = a_i$  then gives values  $b_i$  for the  $y_i$ , so that  $0 \neq Q(a_i) = P(b_i)$ , as desired. 

To finish the proof of Theorem 1.2 using this lemma, we (???) (something about  $\mathbb{C}$  and Jordan normal forms?).

Before proving the normal basis theorem (1.1), we require a useful lemma.

**Lemma 1.2** (Independence of Characters). *Let  $\Delta$  be an abelian group with distinct characters  $\chi_1, \dots, \chi_d : \Delta \rightarrow F^\times$  for some field  $F$ . If there are  $c_i \in F$  such that  $\sum_i c_i \chi_i = 0$ , then all  $c_i = 0$ .*

*Proof.* Assume the  $c_i$  are not all 0. Relabeling if necessary, assume  $c_d \neq 0$ . Since the characters are distinct, there is some  $\delta_0 \in \Delta$  such that  $\chi_1(\delta_0) \neq \chi_d(\delta_0)$ . We

then have two equations that hold for all  $\delta \in \Delta$ :

$$\begin{aligned} 0 &= \sum_i c_i \chi_i(\delta_0 \delta) = \sum_i c_i \chi_i(\delta_0) \chi_i(\delta), \\ 0 &= \chi_1(\delta_0) \sum_i c_i \chi_i(\delta) = \sum_i c_i \chi_1(\delta_0) \chi_i(\delta). \end{aligned}$$

Subtracting these equations gives a new linear dependence equation:  $\sum c'_i \chi_i = 0$ , where  $c'_i = c_i(\chi_i(\delta_0) - \chi_1(\delta_0))$ . By the choice of  $\delta_0$ ,  $c'_d \neq 0$  and  $c'_1 = 0$ . In effect, we have removed  $\chi_1$  from our list of characters and obtained a new non-trivial dependence equation. We may repeat until we obtain  $c\chi_d = 0$  for  $c \neq 0$ , at which point we have a contradiction. Thus, the original  $c_i$  must have all been 0. 🇺🇸

We now prove the normal basis theorem (1.1).

*Proof.* For all  $x \in L$ , there are  $a_g \in K$  indexed over  $g \in G$  so that  $\sum_g a_g g x = 0$ ; for instance, we can choose  $a_g = 0$  for all  $g$ . We want to find  $x$  for which there is no choice of these  $a_g$  where not all  $a_g = 0$ . The previous equation implies that  $\sum_g a_g h^{-1} g x = 0$  for all  $h$ , so we obtain a matrix equation  $M\vec{a} = 0$ , where  $M = [h^{-1} g x]_{h,g}$  and  $\vec{a} = (a_g)_g$ . Note that  $M$  has nothing to do with  $a$ , so if we show that there is a choice of  $x$  for which  $M$  is invertible, we will be done.

As an aside, an explicit example with  $L = K(\sqrt{D})$  and  $x = a + b\sqrt{D}$  is  $M = \begin{pmatrix} x & \bar{x} \\ \bar{x} & x \end{pmatrix}$ , where  $\bar{x} = a - b\sqrt{D}$ .

Now, fix a basis  $e_1, \dots, e_d$  for  $L/K$ . Write  $x = \sum_i x_i e_i$ . Then  $M$  becomes  $[\sum_i x_i h^{-1} g e_i]_{h,g}$ .

We will first treat the case where  $K$  is infinite, so that we can use Lemma 1.1. To that end, we want to find a specialization of the  $x_i$  to  $\eta_i \in L$  such that  $\sum_i \eta_i e_i = 1$  and  $\sum_i \eta_i g e_i = 0$  for  $g \neq 1 \in G$ . We can do this by showing  $A = [g e_i]_{i,g}$  is invertible.  $A$  is invertible iff its transpose is invertible iff its transpose is injective. So, suppose we have  $c_g \in L$  with  $\sum_g c_g g e_i = 0$  for all  $i$ . Since the  $e_i$  are a basis for  $L$ , this means that  $\sum_g c_g g x = 0$  for all  $x \in L$ , and thus that  $\sum_g c_g g = 0$  as functions on  $L$ . But each  $g$  is a character of  $L^\times$ , so by linear independence of characters (Lemma 1.2), we have that all  $c_g = 0$ . Thus  $A$  is invertible and we can find the  $\eta_i$ . The ability to find  $\eta_i$  implies that the polynomial defining  $\det(M)$  can be specialized to be 1, which is in particular not 0. Therefore, by Lemma 1.1, we can find a specialization of the  $x_i$  in  $K$  that makes  $\det(M) \neq 0$ , so that  $x = \sum x_i e_i$  is a normal basis element.

Now we treat the case where  $K = \mathbb{F}_q$  is a finite field. Then  $L = \mathbb{F}_{q^d}$ , and  $G = \mathbb{Z}/d\mathbb{Z}$  is generated by the  $q$ -power Frobenius  $\Phi : x \mapsto x^q$ . In particular, the group ring  $K[G]$  is isomorphic to  $K[x]/(x^d - 1)$ , with the isomorphism sending  $\Phi$  to  $x$ . In particular, we have a map  $K[x] \twoheadrightarrow K[G]$ , so that any  $K[G]$ -module,

like  $L$ , can be considered a  $K[x]$ -module, where  $x$  acts by  $\Phi$ . This allows us to use the structure theorem for f.g. modules over a PID;  $L \cong \bigoplus_{i=1}^n K[x]/(a_i)$ , where we can arrange the  $a_i \in K[x]$  to satisfy  $a_{i+1}$  is a multiple of  $a_i$ . Since  $[L : K] = d < \infty$ , we must have  $a_n \neq 0$ , since otherwise  $L$  would have a subspace isomorphic to  $K[x]$ , which is infinite dimensional over  $K$ . Since  $a_n \neq 0$  and it is a multiple of all other  $a_i$ , the other  $a_i$  cannot be 0. We may assume the  $a_i$  are not constants, since  $K[x]$  modulo a non-zero constant is 0. Each subspace  $K[x]/(a_i)$  has  $K$ -dimension equal to  $\deg a_i$ , so  $d = [L : K] = \sum_i \deg a_i$ .

Now, suppose  $L$  is not cyclic as a  $K[x]$ -module, so that  $n > 1$  above. Then  $\deg a_n < d$ . We also know that  $a_n$  annihilates  $L$ , since it annihilates each  $K[x]/(a_i)$  by virtue of  $a_i | a_n$ . What this means is that if  $a_n = \sum_{j=0}^{d-1} c_j x^j$ , then  $\sum_{j=0}^{d-1} c_j \Phi^j(y) = 0$  for all  $y \in L$ . In other words,  $\sum_j c_j \Phi^j = 0$  as functions on  $L$ . We noted that  $a_n \neq 0$ , so the  $c_j$  are not all 0. But the  $\Phi^j$  are distinct characters of  $L^\times$ , so we obtain a contradiction by independence of characters (Lemma 1.2).


So now we are in the case that  $L \cong K[x]/(a)$  as  $K[x]$ -modules for some degree  $d$  polynomial  $a$ . Suppose  $a = \sum_{j=0}^d c_j x^j$ . Then by similar reasoning as above, we have  $\sum_j c_j \Phi^j = 0$  as functions on  $L$ . But notice that, since  $\Phi$  is an order  $d$  generator for  $G$ , we have  $\Phi^d = 1$ , so  $\sum_j c_j \Phi^j = c_0 + c_d + \sum_{j=1}^{d-1} c_j \Phi^j = 0$ . This is another linear dependence equation of the characters  $1, \Phi, \dots, \Phi^{d-1}$ , so we must have  $c_j = 0$  for  $0 < j < d$  and  $c_0 = -c_d$ . In other words, up to scalar,  $a = x^d - 1$ . Therefore,  $L \cong K[x]/(x^d - 1)$  as  $K[x]$ -modules, say with an isomorphism  $f : L \rightarrow K[x]/(x^d - 1)$ . Let  $y = f^{-1}(1)$ . Then the set  $\{gy \mid g \in G\}$  is the set  $\{\Phi^j y \mid 0 \leq j < d\}$ , which under the isomorphism  $f$  corresponds to the set  $\{x^j \cdot 1 \mid 0 \leq j < d\}$ . This set is certainly a basis for  $K[x]/(x^d - 1)$ , so  $\{gy \mid g \in G\}$  is a basis for  $L$ . Thus,  $y$  is a normal basis element as desired. 🇧🇷

## 2 Oct 3 25

Came in 9 mins late.

Let  $L/\mathbb{Q}$  be Galois with Galois group  $G$ . Let  $\Sigma_L = \text{Hom}(L, \mathbb{C})$ .  $G$  acts on  $\Sigma_L$  by  $g\sigma = \sigma \circ g^{-1}$ .

$\Sigma_L$  up to complex conj. in bijection with  $G/\langle c \rangle$ .



We want to understand the relationship between  $c \in G$  corresponding to  $\sigma \in \Sigma_L$  and  $\Sigma_K$ . There is a surjective map  $\Sigma_L \twoheadrightarrow \Sigma_K$  given by restriction to  $K$ . Elements  $\sigma, g\sigma$  map to the same thing iff  $g \in H$ . Similarly  $g_1\sigma$  and  $g_2\sigma$  map to the same thing iff  $g_1^{-1} = g_2^{-1}h$  for some  $h \in H$ , which means that the fibers of the map  $\Sigma_L \rightarrow \Sigma_K$  are right cosets of  $H$ .

How many embeddings in  $\Sigma_K$  are real? How many are complex? Can write this in terms of cosets...

Can map  $L^\times \rightarrow \prod_{\nu \text{ cplx}} \mathbb{C}^\times \xrightarrow{\log|\cdot|} \prod \mathbb{R}$ .  $G$  acts on the rightmost set by permutation, corresponding to the action of  $G$  on  $G/\langle c \rangle$ . We can express this as  $\mathbb{R}[G/\langle c \rangle]$  or  $\text{Ind}_{\langle c \rangle}^G \mathbb{R}$ . If we restrict to units, we get  $\mathcal{O}_L^\times \otimes \mathbb{R} \rightarrow \text{Ind}_{\langle c \rangle}^G \mathbb{R} = \mathbb{R}^{[G:\langle c \rangle]}$ . Since units have norm one, this map lands in the subspace whose coordinates sum to 0. In particular, letting  $d$  be the rank of  $\mathcal{O}_L^\times$ , we find  $d = [G:\langle c \rangle] - 1$ . Denoting by  $\mathbb{R}$  the trivial representation in  $\mathbb{R}^{[G:\langle c \rangle]}$ , sitting in the diagonal of vectors with equal coordinates, we obtain an isomorphism of  $G$ -modules  $\mathcal{O}_L^\times \otimes R \cong (\text{Ind}_{\langle c \rangle}^G \mathbb{R})/\mathbb{R}$ . For example, if  $L$  is totally real, we get  $\mathcal{O}_L^\times \otimes \mathbb{R} \cong \mathbb{R}[G]/\mathbb{R}$ . Otherwise (or in any case?), we also obtain  $\mathcal{O}_L^\times \otimes \mathbb{C} \cong (\text{Ind}_{\langle c \rangle}^G \mathbb{C})/\mathbb{C}$ .

What is the multiplicity of a  $G$ -irrep  $V$  in  $\mathcal{O}_L^\times \otimes \mathbb{C}$ ? It is  $\dim \text{Hom}_G(V, \mathcal{O}_L^\times \otimes \mathbb{C}) = \dim \text{Hom}_G(V, \text{Ind}_{\langle c \rangle}^G \mathbb{C}) - \dim \text{Hom}_G(V, \mathbb{C})$ . The subtracted term is 1 if  $V$  is trivial and 0 otherwise. The other term can be rewritten with Frobenius reciprocity as  $\dim \text{Hom}_{\langle c \rangle}(V, \mathbb{C}) = \dim(V|c = 1)$ .

**Example 2.1.** Let  $G \cong \mathbb{Z}/3$ . There are three irreducible characters over  $\mathbb{C}$ , called  $1, \chi, \chi^2$ . They are all defined over  $K = \mathbb{Q}(\zeta_3)$ . You can look at  $\chi$  as a 2-dimensional representation over  $\mathbb{Q}$ , by taking a basis  $1, \zeta_3$  for  $K/\mathbb{Q}$ .

In fact for representations of finite groups over  $\mathbb{C}$ , we can always define over a number field. There is a moduli space perspective on this; a purely transcendental representation must correspond to some deformation, and then the deformation corresponds to a non-split extension. By semisimplicity, this cannot happen.

If  $W$  is a representation of  $\mathbb{Q}[G]$ , then  $W \otimes_{\mathbb{Q}} K$  is a representation of  $\mathbb{Q}[G]$  isomorphic to  $W^{[K:\mathbb{Q}]}$ .

**Lemma 2.1.** *If  $W, V$  are reps of  $G$  over a number field  $K$  and if  $W \otimes_K \mathbb{C} \cong V \otimes_K \mathbb{C}$ , then  $V \cong W$ .*

How close is  $\mathcal{O}_L^\times$  to  $\text{Ind}_{\langle c \rangle}^G \mathbb{Z}$ ? Similarly, if  $L/K$  Galois, how close is  $\mathcal{O}_L$  to  $\mathcal{O}_K[G]$ ?

**Example 2.2.** Let  $K = \mathbb{Q}$ ,  $L = \mathbb{Q}(\sqrt{D})$ . Let  $\theta = a + b\sqrt{D}$  with  $ab \neq 0$ . We have  $L \cong \mathbb{Q}[\mathbb{Z}/2]$ . Integrally, the question is whether  $\mathcal{O}_L = \mathbb{Z}\theta \oplus \mathbb{Z}\bar{\theta}$ .

If  $L = \mathbb{Q}(\sqrt{-3})$ , we can pick  $\theta = \frac{1+\sqrt{-3}}{2}$ . If  $L = \mathbb{Q}(\sqrt{-1})$ , there is no choice of  $\theta$ .

In general, if  $\Delta_D \equiv 1 \pmod{4}$ , then we can find  $\theta$ . But if  $\Delta_D \equiv 0 \pmod{4}$ , then we cannot find  $\theta$ . However, we can get an index 2 subgroup.

In general we can do things after inverting finitely many primes.