

# Linear Algebra 2

## Homework 3 – Linear Transformations II

Abraham Murciano

May 10, 2021

1. For the following transformations, our task is to find their matrix representation using standard bases, if the transformation is linear.

(b)  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $T(x, y) = (2x + |y|, -2y)$

This transformation is not linear.

(d)  $T : \mathbb{R}^3 \rightarrow \mathbb{R}$  such that  $T(x, y, z) = y$

Since  $T$  maps from  $\mathbb{R}^3$  to  $\mathbb{R}$ , the matrix must have dimensions such that when multiplying it by a vector in  $\mathbb{R}^3$ , the result must be in  $\mathbb{R}$ . Therefore its dimensions must be three by one. And since we are only interested in the result containing  $y$ , we can use the following matrix which multiplies any input vector  $(x, y, z)$  as follows.

$$\begin{bmatrix} 0 & 1 & 0 \end{bmatrix} \times \begin{bmatrix} x \\ y \\ z \end{bmatrix} = [y]$$

(g)  $T : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  such that  $T(z_1, z_2) = (|z_1 + z_2|, z_1 - 2z_2)$  where  $\mathbb{C}^2$  is a vector space over  $\mathbb{C}$ .

This is not a linear transformation either.

2. Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  be the transformation defined by  $T(x, y, z) = (x - 2y, 2x + y - z)$ .

- (a) To show that this transformation is linear, we must show that it is commutative with addition and with scalar multiplication.

$$\begin{aligned}
T(\vec{u} + \vec{v}) &= ((x_u + x_v) - 2(y_u + y_v), \\
&\quad 2(x_u + x_v) + (y_u + y_v) - (z_u + z_v)) \\
&= (x_u + x_v - 2y_u - 2y_v, 2x_u + 2x_v + y_u + y_v - z_u - z_v) \\
&= (x_u - 2y_u, 2x_u + y_u - z_u) + (x_v - 2y_v, 2x_v + y_v - z_v) \\
&= T(\vec{u}) + T(\vec{v})
\end{aligned}$$

$$\begin{aligned}
T(c\vec{u}) &= (cx - 2cy, 2cx + cy - cz) \\
&= (c(x - 2y), c(2x + y - z)) \\
&= c(x - 2y, 2x + y - z) \\
&= cT(\vec{u})
\end{aligned}$$

- (b) The inverse image of  $(1, 3)$  under  $T$ , i.e.  $\{v \in \mathbb{R}^3 : T(v) = (1, 3)\}$ , can be described with the following system of equations.

$$\begin{aligned}
x - 2y &= 1 \\
2x + y - z &= 3
\end{aligned}$$

Solving this system gives us the line  $(x, y, z) = (\frac{7}{5}, \frac{1}{5}, 0) + t(-\frac{2}{5}, \frac{1}{5}, \frac{1}{5})$ .

- (c) To find a basis for the kernel of  $T$ , it suffices to find one vector  $\vec{v} \in \ker T$ , since we know from the previous question that the dimension of the kernel must be one, i.e. a line.

Letting  $x := 2$  and  $y := 1$ , we have  $x - 2y = 0$ . Now to obtain  $2x + y - z = 0$  as well, we can simply set  $z := 2x + y = 5$ . Therefore  $(2, 1, 5)$  is a basis of the kernel of  $T$ .

- (d) Since the dimension of the kernel is one, and the domain of the transformation has a dimension of three, we must conclude that the image of  $T$  has a dimension of  $\dim(\mathbb{R}^3) - \dim(\ker T) = 3 - 1 = 2$ . Therefore any two independent vectors in the image of  $T$  will form a basis for the image.

To find two such vectors, we can apply  $T$  to two vectors in the domain, then verify that they are indeed independent. We shall choose the vectors  $(1, 0, 0)$  and  $(0, 0, 1)$ . Applying  $T$  to these gives us a basis  $\{(1, 2), (0, -1)\}$ , which is indeed independent.

4. Consider the transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  whose matrix is the following.

$$[T] = \begin{bmatrix} 1 & 1 & 3 & 4 \\ -2 & 2 & 1 & 1 \end{bmatrix}$$

- (a)  $T$  cannot be one to one, since we shall show in part (c) that it is onto. And if it would be both one to one and onto, then it would be invertible, which is impossible since the range and domain have different dimensions.
- (b) Since the transformation is onto, the dimension of the image is two. Therefore the dimension of the kernel is  $4 - 2 = 2$ . So it suffices to find two independent vectors in  $\ker T$ , and those will form a basis.

To find these, we can treat the matrix as a homogeneous one, to indicate that  $T(x_1, x_2, x_3, x_4) = (0, 0)$ , then row reduce it to reduced row echelon form.

$$\begin{bmatrix} 1 & 0 & \frac{5}{4} & \frac{7}{4} \\ 0 & 1 & \frac{7}{4} & \frac{9}{4} \end{bmatrix}$$

This gives us the following system of equations which holds when  $[T]\vec{x} = (0, 0)$ .

$$\begin{aligned} x &= -\frac{5}{4}x_3 - \frac{7}{4}x_4 \\ y &= -\frac{7}{4}x_3 - \frac{9}{4}x_4 \end{aligned}$$

In matrix form, this equation can be written like this.

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -\frac{5}{4}x_3 - \frac{7}{4}x_4 \\ -\frac{7}{4}x_3 - \frac{9}{4}x_4 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -\frac{5}{4} \\ -\frac{7}{4} \\ 1 \\ 0 \end{bmatrix} x_3 + \begin{bmatrix} -\frac{7}{4} \\ -\frac{9}{4} \\ 0 \\ 1 \end{bmatrix} x_4$$

This means that for all  $x_3, x_4 \in \mathbb{R}$ , we can find some vector  $\vec{x} \in \ker T$  according to the previous equation. Therefore if we choose  $x_3 = 1, x_4 = 0$  we get one vector in the kernel,  $(-\frac{5}{4}, -\frac{7}{4}, 1, 0)$ . And choosing  $x_3 = 0, x_4 = 1$  gives us another vector in the kernel,  $(-\frac{7}{4}, -\frac{9}{4}, 0, 1)$ . Since these are independent, they form a basis for the kernel.

- (c)  $T$  must be onto, since two of the vectors in its image, for example  $(1, -2)$  and  $(1, 2)$ , span the entire range  $\mathbb{R}^2$ , since they are linearly independent.
5. (b) Given the transformation  $T : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  over the vector space  $\mathbb{C}$  represented by the matrix below, we are tasked with proving that it is invertible, and we seek the inverse transformation matrix.

$$[T] = \begin{bmatrix} i & 2i \\ 2 & i \end{bmatrix}$$

A linear transformation is invertible if and only if its transformation matrix is invertible. And a transformation matrix is invertible if and only if its determinant is not zero.

$$\det[T] = i \times i - 2 \times 2i = i^2 - 4i = -1 - 4i \neq 0$$

$$[T^{-1}] = [T]^{-1} = \frac{1}{-1 - 4i} \begin{bmatrix} i & -2i \\ -2 & i \end{bmatrix} = \frac{4i - 1}{17} \begin{bmatrix} i & -2i \\ -2 & i \end{bmatrix}$$