

2 **Draft: Constraints on Simplified Dark Matter Models** 3 **using Mono-X Collider Searches**

4 **A. J. Brennan,^{a,1} M. F. McDonald,^a J. Gramling^b and T. D. Jacques^b**

5 ^a*The University of Melbourne, Parkville 3010, Australia*

6 ^b*SISSA/ISAS, via Bonomea 265, 34136 Trieste, Italy*

E-mail: a.brennan@student.unimelb.edu.au,
7 milliem@student.unimelb.edu.au, johanna.gramling@cern.ch,
thomas.jacques@sissa.it

8 ABSTRACT: The use of simplified models of dark matter is becoming increasingly prevalent
9 in collider searches, and while early Run II results are beginning to appear, we look to see
10 what further information can be extracted from Run I results. We consider three ‘standard’
11 simplified models that couple quarks to fermionic singlet dark matter: an s -channel vector
12 mediator with vector or axial-vector couplings, and a t -channel scalar mediator. Upper lim-
13 its on the couplings are calculated, and compared across three alternate channels, namely,
14 mono-jet, mono- Z (leptonic) and mono- W/Z (hadronic). The strongest limits are observed
15 in the mono-jet channel, however the computational simplicity, and absence of significant
16 t -channel model width effects, in the mono-boson channels make these a straightforward
17 and fast alternative.

¹Corresponding author.

18	Contents	
19	1 Introduction	2
20	2 Simplified Model Phenomenology	3
21	2.1 Model Descriptions	3
22	2.2 The Mono- $X + E_{\text{T}}^{\text{miss}}$ Signature	4
23	2.3 Mass and Coupling Points	5
24	2.4 Treatment of the width	6
25	3 Recasting mono-X constraints	9
26	3.1 Signal Simulation	9
27	3.1.1 Parton Matching Scheme	9
28	3.2 Mono-jet Constraints	10
29	3.3 Mono- Z Constraints	10
30	3.4 Mono- W/Z Constraints	11
31	4 Limits on the coupling $\sqrt{g_q g_\chi}$	13
32	4.1 Mono-jet channel	13
33	4.2 Mono- Z channel	14
34	4.3 Mono- W/Z channel	14
35	5 Conclusion	15
36	5.1 Comparison with Relic Density Constraints	15
37	5.2 Comparison with Direct Detection Constraints	21
38	5.3 Discussion	21
39	6 Acknowledgements	21
40	A Limit setting strategy	21
41	A.1 Nominal Values	21
42	A.2 Uncertainty Estimation	22
43	B Validation of signal simulation and event selection procedures	22
44	B.1 Monojet Channel	22
45	B.2 Mono- Z Channel	23
46	B.3 Mono- W/Z Channel	25

1 Introduction

Simplified models have emerged as a powerful tool for the interpretation of collider, direct and indirect detection signals of dark matter (DM). Previously, ATLAS and CMS searches for DM were conducted within the context of both Effective Field Theories (EFTs) [1, 5, 32, 33] and full UV-complete theories such as Supersymmetry [10–12, 40]. I suggest we just list official ATLAS and CMS studies here to avoid ‘opening the floodgates’, so I’ve removed a few references - TJ. The latter approach, though well-motivated, is typified by a broad parameter space and generally yields results which are insensitive to the wider class of DM models. EFT constraints, in comparison, are applicable to a broad range of models and rely on the specification of only a small set of parameters, namely the suppression scale, M_* , and the DM mass, m_{DM} [25]. In the EFT framework, interactions between the dark and Standard Model (SM) sector are parametrised by a set of higher-dimensional effective operators, that arise when the mass of the mediating particle is assumed to be significantly larger than the momentum transferred in a given interaction. Where this is not the case, the EFT prescription can produce constraints which detour dramatically from those of the associated UV-complete model [24–28]. This is not so important in direct detection experiments where the momentum transferred in the scattering of DM particles with heavy nuclei is generally of the order of tens of MeV [13, 14], or in indirect searches where the annihilations of non-relativistic DM particles in the galactic halo occur with momentum transfers of order m_{DM} []. However, for hadron collider searches - where the accessible center of mass energy of two colliding baryons may be sufficient to produce the mediator on-shell - the range of validity of the EFT prescription is significantly diminished. Indeed, recent works have shown the EFT approach to be problematic in certain cases for the interpretation of data collected during the $\sqrt{s} = 8$ TeV Run I of the Large Hadron Collider (LHC) [15–17]. In light of this, simplified models have become the preferred tool for the interpretation of collider DM searches [18, 31, 53, 55, 56].

In a nutshell, a simplified model (SiM) arises when the heavy mediator which was integrated out in the EFT framework is reintroduced. Like EFTs, SiMs admit the comparison of results obtained in the different avenues of dark matter study [] and are defined by a relatively small set of parameters - namely m_{DM} , the mass of the mediator M_{med} , and the SM-mediator and DM-mediator coupling strengths, g_q and g_χ (or $g_{q\chi}$ in the case of a single, SM-DM-mediator coupling). Unlike EFTs, constraints calculated within the context of a SiM are valid across a broad energy range.

In this paper, we examine a phenomenologically distinct set of SiMs. In particular, we place constraints on the SiMs corresponding to the simplest UV-completions of the D5 (vector) and D8 (axial-vector) effective operators in the s -channel¹. We also include a case

¹The D5 and D8 operators form a nice starting point in the analysis of simplified models as they have been studied exhaustively in the past (see refs. [1, 4, 5, 7, 15–17, 32, 33] among others). This attention is motivated by the fact that collider limits for the D5 (D8) operator can be readily transformed into limits on spin-independent (spin-dependent) DM-nucleon scattering and vice versa. With the exception of D1 (see sec. ??), and D9 and D11 (which have no simple simplified model counterparts []), the remaining effective operators induce elastic scattering which is suppressed by powers of the DM velocity or the momentum transferred [19]. Hence, these operators are not often considered in the literature.

in which a scalar mediator is exchanged in the t -channel, motivated by its analog of squark exchange in Supersymmetry. In the heavy mediator limit, this model can be expressed as a combination of operators D5 to D8 via a Fierz transformation.

The models are constrained using public results from mono- X + missing transverse energy (E_T^{miss}) searches conducted by the ATLAS Collaboration. Specifically we focus on searches where X is either a parton (manifesting in the detector as a narrow-radius jet), a leptonically-decaying Z boson, or a hadronically-decaying W or Z boson (manifesting as a large-radius jet). The purpose of this work is to strengthen existing SiM limits [23, 57] using the full 20.3 fb^{-1} of Run I ATLAS data, and to explore an enhanced phase space with respect to the mediator and DM masses and the relative strength of the couplings to the visible and dark sectors. We choose to treat the mediator width as the minimal value naturally arising, as an alternative to many past analyses which handle this as a fixed value []. We extend the study by providing a cross-check and comparison of the performance of the three targeted collider detection channels, and compare against relic density and direct detection constraints.

The remainder of the paper is organised as follows. Section 2 contains a compendium of the SiMs chosen for analysis and the associated collider phenomenology. Section 3 outlines the techniques used to recast mono- X + E_T^{miss} limits on the visible cross-section for any new physics process into constraints on SiMs, and specifically on the couplings g_q and g_χ . Lastly, our results are presented in Section 4 along with a discussion of the implications of this work. Appendices A and B include details of the limit setting and analysis validation procedures.

2 Simplified Model Phenomenology

2.1 Model Descriptions

We begin with a short set of assumptions: that the DM particle, χ , is a weakly interacting Dirac fermion, that it is a singlet under the SM, and that it is the lightest stable new particle. Additionally the new sector is assumed to couple only to the SM quarks; while possible coupling to SM leptons e.g. [?] or gluons e.g. [?] has been studied elsewhere, it is beyond the scope of this paper. The nature of the mediating particle then results from these assumptions: in the s -channel it is chosen to be a vector particle and must also be a SM singlet, denoted ξ , while in the t -channel it is identified as scalar and is necessarily charged and coloured, and labelled ϕ .

The s -channel models chosen for analysis are characterised by vector (sV) or axial-vector (sA) couplings to both the dark and SM sectors. They are described by the following interaction Lagrangians:

$$\mathcal{L}_{sV} \supset -\xi_\mu \left[\sum_q g_q \bar{q} \gamma^\mu q - g_\chi \bar{\chi} \gamma^\mu \chi \right], \quad (2.1)$$

$$\mathcal{L}_{sA} \supset \xi_\mu \left[\sum_q g_q \bar{q} \gamma^\mu \gamma_5 q - g_\chi \bar{\chi} \gamma^\mu \gamma_5 \chi \right], \quad (2.2)$$

where the sum is over all quarks. For the couplings g_q and g_χ to remain within the perturbative regime, they are required to satisfy $g_q, g_\chi \leq 4\pi$, though stronger perturbativity requirements do exist [15].

The t -channel model considered in this paper (abbreviated tS) is characterised by a scalar mediator and is motivated by analogy with a common aspect of Supersymmetric models: neutralino DM interacting with the SM sector via t -channel exchange of a squark [20]. Note that in this Supersymmetric scenario the DM particle is a Majorana fermion. SiMs in which χ is of Majorana type are kinematically identical to the corresponding Dirac cases (requiring multiplication of the cross-section by a simple factor in order to compute limits) and so are not covered here². The exception to this rule is the s -channel vector mediator model, which vanishes if χ is a Majorana fermion [21].

In the tS model, the mediator is allowed to couple to either the left or right-handed quarks as an SU(2) doublet or singlet respectively. Since the LHC is insensitive to the chirality of the quarks, we assume for simplicity that ϕ couples to the left-handed quarks only, and is itself an SU(2) doublet, allowing radiation of a W boson. To avoid different couplings to quarks of different generations, and to remain in step with the DM forum recommendations [], we include three generations of mediator ϕ , with equal masses and couplings. The interaction Lagrangian for this model is then:

$$\mathcal{L}_{tS} \supset \sum_Q g_{q\chi} \bar{Q} P_R \phi \chi + \text{h.c.}, \quad (2.3)$$

where the sum is over the three quark doublets, $g_{q\chi}$ is the scalar coupling of the incoming quark, ϕ and χ , and P_R is the usual chiral projection operator.

2.2 The Mono- $X + E_T^{\text{miss}}$ Signature

The mono- $X + E_T^{\text{miss}}$ (abbreviated to mono- X) is a popular collider signal in the search for new physics, particularly in the search for dark matter. Since DM particles are not expected to interact with detector material, they appear as missing transverse energy when balanced against a visible object, X , that is radiated from the initial or intermediate state (where the latter is permitted in the t -channel model). For the s -channel SiMs discussed above, only initial-state radiation is permitted; see figs. 1a and 1b for examples. For the tS model, radiation of a gluon or electroweak (EW) boson is permitted both from initial state partons (fig. 1c) or from the mediator (fig. 1d).

To fully exploit the potential of the ATLAS detector to record and identify a vast array of particle types, we consider for analysis three scenarios: 1.) X is a parton which hadronises to form a jet, 2.) X is a Z boson which decays into an $\ell^+\ell^-$ pair, and 3.) X is a hadronically-decaying W or Z boson. In the context of DM searches, these signatures are referred to as mono-jet, mono- Z , and mono- W/Z respectively. Note that the mono-jet scenario is the most likely scenario at the LHC owing to the strength of the strong coupling yet the mono- Z channel has the advantage of a relative simple and clean signature. OR: Leptons are in general much cleaner and simpler than jets, so this channel is included here

²The exception being in the validation of the mono- Z channel, see Sec. B.2.




Figure 1: Representative dark matter pair-production processes with a gluon or W/Z boson in the final state for the s -channel (a,b) and t -channel (c,d) models.

to investigate whether the reduction in systematic uncertainties can provide easily-obtained results that are comparable to the more complicated mono-jet channel. The inclusion of the mono-WZ channel is motivated by its fundamentally different signature structure wherein event selection is based on large R jets that are consistent with having come from an EW boson.

NEW: The most likely scenario at the LHC is production of a jet alongside the invisible χ pair, as a result of the strong coupling and prevalence of partons in the initial state. However, to fully exploit the potential of the ATLAS detector to record and identify a vast array of particle types, we can include the alternative mono-W/Z channels to potentially glean further information. We can take advantage of the relative cleanliness and simplicity of leptons in the leptonically-decaying mono-Z channel, or the large hadronic branching fraction, and developing techniques to identify jets resulting from EW bosons, in the hadronically-decaying mono-W/Z channel. In both cases, the large multi-jet background is reduced, and complications in jet production such as parton-matching can be ignored, making these an interesting alternative where speed, efficiency and a reduction in jet-associated uncertainties may make up for the loss in sensitivity.

2.3 Mass and Coupling Points

A representative set of dark matter and mediator masses, listed in table 1, are chosen for study in each detection channel. DM masses of 3, 30 and 300 GeV were also included in the

m_χ [GeV]	M_{med} [GeV]	s-channel		t-channel
		g_q	g_χ	$g_{q\chi}$
1, (3), 10, (30), 100, (300), 1000	1, 2, 10, 20, 100, 200, 1000, 2000	1	0.2, 0.5, 1, 2, 5	1

Table 1: Mass and coupling points chosen for the analysis of simplified dark matter models. Values in brackets are only included in the mono- Z channel. The mediator masses are primarily representative of three regimes: (near-)degenerate ($M_{\text{med}} \approx m_\chi$), on-shell ($M_{\text{med}} \geq 2m_\chi$), and EFT-like ($\sqrt{\hat{s}} \ll M_{\text{med}}$). For the t -channel model, $M_{\text{med}} > m_\chi$ is also required.

mono- Z channel, where ease of production can afford higher granularity. All (m_χ, M_{med}) combinations are permitted in the sV and sA models, while in the tS model M_{med} must be greater than m_χ to ensure stability of the DM particle. The couplings g_q and $g_{q\chi}$ are set to unity, while the DM-mediator coupling in the s -channel models, g_χ , was varied from 0.2 to 5. The mediator masses are chosen to cover a broad range of parameter space and to coincide with predominantly three regimes: (near-)degenerate ($M_{\text{med}} \approx m_\chi$), on-shell ($M_{\text{med}} \geq 2m_\chi$), and EFT-like ($\sqrt{\hat{s}} \ll M_{\text{med}}$)³. We also allow for the possibility of a light mediator/heavy WIMP scenario ($M_{\text{med}} < m_\chi$) in the sV and sA models.

2.4 Treatment of the width

An important factor when considering simplified models is to ensure the mediator width is treated appropriately, as it impacts both the cross-section calculation and, in some cases, the kinematic behaviour of the model.

Following the DM Forum recommendations [?], we use the minimal width, allowing coupling to all kinematically accessible quarks. We assume minimal flavour violation, which implies a universal coupling to all quark flavours. The minimum width for each model is given by:

$$\begin{aligned} \Gamma_{sV} = & \frac{g_\chi^2 M}{12\pi} \left(1 + \frac{2m_\chi^2}{M^2}\right) \left(1 - \frac{4m_\chi^2}{M^2}\right)^{\frac{1}{2}} \Theta(M - 2m_\chi) \\ & + \sum_q \frac{g_q^2 M}{4\pi} \left(1 + \frac{2m_q^2}{M^2}\right) \left(1 - \frac{4m_q^2}{M^2}\right)^{\frac{1}{2}} \Theta(M - 2m_q) \end{aligned} \quad (2.4)$$

³A recent study by Alves et al. found that EFT results do not apply to mediators with a mass less than 2.5 TeV at the LHC during Run I [35]. I can't find this, could someone check? Maybe this is covered by the validity issues citation anyway

$$\begin{aligned}\Gamma_{sA} = & \frac{g_\chi^2 M}{12\pi} \left(1 - \frac{4m_\chi^2}{M^2}\right)^{\frac{3}{2}} \Theta(M - 2m_\chi) \\ & + \sum_q \frac{g_q^2 M}{4\pi} \left(1 - \frac{4m_q^2}{M^2}\right)^{\frac{3}{2}} \Theta(M - 2m_q)\end{aligned}\tag{2.5}$$

$$\begin{aligned}\Gamma_{tS} = & \sum_q \frac{g_{q\chi}^2 M}{16\pi} \left(1 - \frac{m_q^2}{M^2} - \frac{m_\chi^2}{M^2}\right) \\ & \times \sqrt{\left(1 - \frac{m_q^2}{M^2} + \frac{m_\chi^2}{M^2}\right)^2 - 4\frac{m_\chi^2}{M^2}} \Theta(M - m_q - m_\chi)\end{aligned}\tag{2.6}$$

191 It is possible that the mediator may decay to other SM or BSM particles [?], but this
 192 is not expected to have a large effect on the kinematic distribution as long as the width
 193 remains relatively small [?]. The generator treatment of the mediator as a Breit-Wigner
 194 propagator, rather than a true kinetic propagator, breaks down for large widths [57]. This
 195 ref. also showed that for $\Gamma/M \lesssim 0.5$ the following scaling relations approximately hold:

$$\sigma \propto \begin{cases} g_q^2 g_\chi^2 / \Gamma & \text{if } M_{\text{med}} \geq 2m_{\text{DM}} \\ g_q^2 g_\chi^2 & \text{if } M_{\text{med}} < 2m_{\text{DM}} \end{cases}\tag{2.7}$$

196 in the sV and sA models, and

$$\sigma \propto g_{q\chi}^4\tag{2.8}$$

197 in the tS model. When valid, these approximations allow us to greatly simplify our limit
 198 calculations, and for this reason, we restrict our primary results to regions of parameter
 199 space where $\Gamma/M < 0.5$.

200 To simplify the procedure further, we can take advantage of the fact that (with the
 201 notable exception of the tS model in the mono-jet channel) for each point in (m_χ, M_{med})
 202 phase space, the mediator width (and therefore the couplings) do not greatly affect each
 203 model's kinematic behaviour. This is demonstrated in fig. 2, where for the sV (representing
 204 both the sV and sA models) and tS models, we plot a simplified $E_{\text{T}}^{\text{miss}}$ distribution, as a
 205 proxy for the full selection in each analysis, for two mass points and a demonstrative set of
 206 couplings such that $\Gamma < M_{\text{med}}/2$. The $E_{\text{T}}^{\text{miss}}$ distribution is predominantly independent of
 207 the mediator width for the s -channel models in the mono-jet channel, and all models in the
 208 mono- Z ⁴ channel. However, there is a clear variation in the kinematic behaviour of the tS
 209 model in the mono-jet channel, which can be attributed to additional diagrams (accessible
 210 only in this channel) featuring a gluon in the initial state and subsequently allowing the
 211 mediator to go on-shell. In this scenario, when the resulting quark and DM particle are

⁴In this discussion, the mono- W/Z channel can be assumed to follow the same logic as for the mono- Z channel.




Figure 2: The E_T^{miss} distribution of the sV and tS models in the mono-jet and mono- Z channels, for some demonstrative masses. The parameter μ is defined as Γ/M_{med} , and is used to demonstrate the impact of a changing width; in particular, the tS model in the mono-jet channel shows clear width-dependence. The widths are obtained with couplings of 0.1, 1, and 5 where $\mu < 0.5$ remains true.

both small compared to the mediator mass, they share equally its energy leading to a peak in the E_T^{miss} distribution at approximately half the mediator mass.

In the cases where the model behaviour is independent of the width, we can greatly simplify the calculations by assuming the impact of the selection cuts in each channel is unchanged for each masspoint; that is, independent of the couplings. In this case, a simple rearrangement of eqns. 2.7 and 2.8 allows us to obtain upper limits on the model couplings (see App. A for further details of this calculation).

A proper study of the tS model within the mono-jet channel, where altering the coupling can lead to changed kinematic behaviour, has been performed elsewhere [23], and requires the production of individual samples for each coupling point. This, combined with the challenges associated with including differing orders of α_s , make the generation process computationally expensive compared to the mono- Z and mono- W/Z channels. We therefore exclude an analysis of the tS model in the mono-jet channel in this work.

3 Recasting mono- X constraints

The procedure for recasting existing mono- X analyses to obtain SiM constraints follows a simple cut-and-count methodology. Firstly, signal events are simulated (described below in section 3.1) with object p_T smearing applied to approximate the detection efficiency of the ATLAS detector, ϵ . The event selection criteria of the mono- X analysis of interest is then applied to the simulated signal samples. Events surviving the selection criteria are counted to determine the likelihood of a dark matter event being observed (referred to as the acceptance, \mathcal{A}), which is then used in combination with channel-specific model-independent limits on new physics events to limit the parameter phase space of a given model. For a comprehensive description of the recasting procedure, see appendix A.

In this paper, mono-jet constraints are derived from a search for new phenomena conducted by the ATLAS Collaboration using pp collisions at $\sqrt{s} = 8$ TeV as described in ref. [41]. Similarly, the leptonic mono- Z and hadronic mono- W/Z constraints are derived from ATLAS dark matter searches originally optimised for the D1, D5 and D9 effective operators [47?]. These analyses are described in further detail in sections 3.2, 3.3 and 3.4 respectively.

3.1 Signal Simulation

Monte Carlo simulated event samples are used to model the expected signal for each channel and for each simplified model. Leading order matrix elements for the process $pp \rightarrow X + \chi\bar{\chi}$ (where X is specifically one or two jets⁵, a $Z(\rightarrow \ell^+\ell^-)$ boson or a $W/Z(\rightarrow jj)$ boson) are first simulated using MADGRAPH5_AMC@NLO v2.2.2 [49] with the MSTW2008lo68cl PDF [50]. During this stage, the renormalisation and factorisation scales are set to the default sum of $\sqrt{m^2 + p_T^2}$ for all particles in the final state. Showering and hadronisation are then performed by PYTHIA 8.201 [58] with the appropriate PDF and using the ATLAS UE Tune AU2-MSTW2008LO [51]. Reconstruction of small-radius jets (from hereon referred to just as ‘jets’) for the mono-jet channel is performed by FastJet [52] using the anti- k_T algorithm with radius parameter $R = 0.4$. Similarly, reconstruction of large-radius jets for the mono- W/Z channel is performed using the Cambridge-Aachen algorithm with $R = 1.2$. The latter channel also includes a mass-drop filtering procedure with $\mu = 0.67$ and \sqrt{y} ⁶ = 0.4 (see ref. [53] for further details), which favours large- R jets with two balanced subjets, consistent with the decay of an EW boson to a (potentially-boosted) dijet pair. Lastly, the detector response is approximated by applying a Gaussian smearing factor to the p_T of all leptons and jets.

3.1.1 Parton Matching Scheme

In the ATLAS mono-jet analysis, matching of partons generated in MADGRAPH5 to jets generated in PYTHIA 8 is performed using the MLM scheme, with two matching scales, or values of ‘QCUT’, per mass/coupling point. The QCUT values span a broad kinematic

⁵Jets are seeded by any parton excluding the (anti-)top quark.

⁶ $\sqrt{y} = \min(p_{T_{j_1}}, p_{T_{j_2}})\Delta R/m_{jet}$ is the momentum balance of the two leading subjets.

range in combination with a cut placed on the leading jet p_T per event to avoid double-counting. This treatment aims to mitigate the impact of the matching scale on the shape of the p_T and E_T^{miss} distributions; that is, to reduce the uncertainty in those areas of phase space where the mediator mass is significantly larger or smaller than the QCUT value. For the analysis of SiMs, we use instead a single matching scale of 80 GeV. Though not ideal, this approach suitably reproduces the results of the ATLAS mono-jet analysis for the masses of interest (see Sec. B.1). Importantly, it also reduces the complexity and computational expense involved in estimating limits for the mono-jet channel.

We now move to a discussion of each of the mono- X channels separately.

3.2 Mono-jet Constraints

The ATLAS mono-jet + E_T^{miss} analysis [41] was originally designed to set limits on three new physics scenarios, the most relevant of which is the production of WIMP DM within the context of seven (?) effective operators. The analysis also includes a brief study of a Z' DM model which is analogous to our sV model.

Signal selection is carried out based on at least one hard jet recoiling against missing energy. To ensure that the correct back-to-back jet + E_T^{miss} topology is selected events are required to have a leading jet, j_1 , with $p_T > 120$ GeV and $|\eta| < 2.0$ satisfying $p_T^{j_1}/E_T^{\text{miss}} > 0.5$. Surviving events must then satisfy $|\Delta\phi(j, \vec{E}_T^{\text{miss}})| > 1.0$, where j is any jet with $p_T > 30$ GeV and $|\eta| < 4.5$. This criterion reduces the multijet background contribution where the large E_T^{miss} originates mainly from jet energy mismeasurement. Note that there is no upper limit placed on the number of jets per event. The contribution from the dominant background processes, W/Z +jets, is managed with a veto on events containing muons or electrons with $p_T > 7$ GeV. A further veto is placed on events containing isolated tracks⁷ with $p_T > 10$ GeV and $|\eta| < 2.5$, to reduce the contribution from non-identified leptons (e , μ or τ) in the final state. Lastly, nine separate signal regions are defined with increasing lower thresholds on E_T^{miss} , which range from 150 GeV to 700 GeV as shown in table 2.

The ATLAS mono-jet analysis revealed no significant deviation of observed events from the expected SM backgrounds in the Run 1 8 TeV dataset. Subsequently, model-independent limits on new physics signatures were provided in terms of the visible cross-section, $\sigma \times \mathcal{A} \times \epsilon$; these are listed in table 2.

The signal simulation procedure outlined in sec. 3.1 and implementation of the selection criteria discussed above were validated for the mono-jet channel via reproduction of ATLAS limits on the suppression scale, $M_\star \equiv M_{\text{med}}/\sqrt{g_q g_\chi}$, for the Z' model. The details of this process are contained in appendix B.1. Importantly, we observe agreement within $\sim 12\%$ for all samples.

3.3 Mono- Z Constraints

The ATLAS mono- $Z(\rightarrow \ell^+ \ell^-)$ + E_T^{miss} analysis [47] was principally designed to constrain a set of EFT models of DM. As a secondary focus, it also includes a short study of a

⁷A track is considered isolated when no additional track with $p_T > 3$ GeV lies within a cone of radius 0.4 around it.

Signal Region	E_T^{miss} threshold [GeV]	$\sigma \times \mathcal{A} \times \epsilon$ [fb]
SR1	150	726 (935)
SR2	200	194 (271)
SR3	250	90 (106)
SR4	300	45 (51)
SR5	350	21 (29)
SR6	400	12 (17)
SR7	500	7.2 (7.2)
SR8	600	3.8 (3.2)
SR9	700	3.4 (1.8)

Table 2: The ATLAS mono-jet E_T^{miss} signal regions and corresponding observed (expected) model-independent upper limits on $\sigma \times \mathcal{A} \times \epsilon$ at 95% confidence level. Adapted from Ref. [41].

t -channel simplified model similar to our tS model.

The selection criteria for this analysis are summarised as follows (see the paper for a full description). Electrons (muons) are required to have a p_T greater than 20 GeV, and $|\eta|$ less than 2.47 (2.5). Two opposite-sign, same-flavour leptons are selected, and required to have invariant mass and pseudorapidity such that $m_{\ell\ell} \in [76, 106]$ GeV and $|\eta^{\ell\ell}| < 2.5$. The reconstructed Z boson should be approximately back-to-back and balanced against the E_T^{miss} , ensured with the selections $\Delta\phi(\vec{E}_T^{\text{miss}}, p_T^{\ell\ell}) > 2.5$ and $|p_T^{\ell\ell} - E_T^{\text{miss}}| / p_T^{\ell\ell} < 0.5$. Events containing a jet with $p_T > 25$ GeV and $|\eta| < 2.5$ are vetoed. Events are also vetoed if they contain a third lepton with $p_T > 7$ GeV. The signal regions are defined by increasing lower E_T^{miss} thresholds: $E_T^{\text{miss}} > 150, 250, 350, 450$ GeV.

A cut-and-count strategy is used to estimate the total observed yields and expected SM backgrounds in each signal region. The limits on $\sigma \times \mathcal{A} \times \epsilon$ are not publicly available, so we take the numbers of expected and observed events, along with the associated uncertainties, and convert these into model-dependent upper limits with a single implementation of the HistFitter framework [54] using a frequentist calculator and a one-sided profile likelihood test statistic (the LHC default). The results of this process are displayed in table 3. Note that we use signal regions 1 and 2 only, as our simplified HistFitter approach is inadequate to handle the very low statistics of signal regions 3 and 4. These upper limits are also used for the validation of the mono- Z signal generation and selection procedures (see app. B.2).

3.4 Mono- W/Z Constraints

The ATLAS mono- $W/Z + E_T^{\text{miss}}$ search [48] was aimed at constraining the spin-independent effective operators C1, D1, and D5, and the spin-dependent operator D9. The search was originally designed to exploit the constructive interference of W boson emission from opposite-sign up-type and down-type quarks, leading to DM production wherein the mono- W channel is dominant. Recent studies [] have revealed this scenario to violate unitarity and so we ignore it in this analysis. (should we not add some motivation on why we include

Signal Region	E_T^{miss} threshold [GeV]	$\sigma \times \mathcal{A} \times \epsilon$ [fb]
SR1	150	1.59 (1.71)
SR2	250	0.291 (0.335)

Table 3: The ATLAS mono- Z E_T^{miss} signal regions and corresponding observed (expected) model-independent upper limits on $\sigma \times \mathcal{A} \times \epsilon$ at 95% confidence level. Adapted using HistFitter from ref. [47].

326 it? Millie, you commented out these parts, maybe you could pick your favourite sentence
327 or so and put it back in, in case you agree?)

328 The mono- W/Z event selection is carried out as follows. Large-radius jets are selected
329 using a mass-drop filtering procedure (see sec. 3.1) to suppress non- W/Z processes. Events
330 are required to contain at least one large- R jet with $p_T > 250$ GeV, $|\eta| < 1.2$ and a mass,
331 m_{jet} , within a 30-40 GeV window of the W/Z mass (i.e. $m_{\text{jet}} \in [50, 120]$ GeV). In order
332 to reduce the $t\bar{t}$ and multijet backgrounds, a veto removes events containing a small- R jet
333 with $\Delta\phi(\text{jet}, E_T^{\text{miss}}) < 0.4$, or containing more than one small- R jet with $p_T > 40$ GeV,
334 $|\eta| < 4.5$, and $\Delta R(\text{small-}R \text{ jet, large-}R \text{ jet}) > 0.9$. Electrons, muons and photons are vetoed
335 if their p_T is larger than 10 GeV and they lie within $|\eta| < 2.47$ (electrons), 2.5 (muons),
336 2.37 (photons). Two signal regions were defined with $E_T^{\text{miss}} > 350$ GeV and $E_T^{\text{miss}} > 500$
337 GeV.

338 The ATLAS analysis used a shape-fit of the mass distribution of the large- R jet to
339 estimate the background yields in the two signal regions, along with the associated sta-
340 tistical and systematic uncertainties. As in the mono- Z case, we do not take the shapes
341 into account but convert the published number of expected and observed events into upper
342 limits on the expected and observed number of new physics events using the HistFitter
343 framework. For the $E_T^{\text{miss}} > 500$ GeV signal region, we obtain the limits shown in table 4.
344 We do not consider the first signal region with $E_T^{\text{miss}} > 350$ GeV in the recasting proce-
345 dure, since the cut-and-count limits extracted could not be convincingly validated. The
346 high E_T^{miss} signal region was found to be optimal for most operators studied by the ATLAS
347 analysis. (add that in other cases the two perform similar?)

Signal Region	E_T^{miss} threshold [GeV]	$\sigma \times \mathcal{A} \times \epsilon$ [fb]
SR2	500	1.35 (1.34)

Table 4: The ATLAS mono- W/Z E_T^{miss} signal region considered in this work and corresponding observed (expected) model-independent upper limits on $\sigma \times \mathcal{A} \times \epsilon$ at 95% confidence level. Adapted using HistFitter from ref. [47].

348 4 Limits on the coupling $\sqrt{g_q g_\chi}$

349 The 95% confidence level upper limits on the sV and sA model coupling combination
 350 $\sqrt{g_q g_\chi}$, and the tS model coupling $g_{q\chi}$, obtained from each of the mono- X channels, are
 351 presented in figs. 3-8. These quantities were evaluated as described in appendix A and
 352 correspond to the best limits of each signal region tested.

353 In each plot, limits are shown ranging from <0.01 to the upper perturbative limit
 354 for each coupling, 4π ⁸; where a limit was calculated to be larger than this, that limit
 355 is considered meaningless and the region is coloured grey. The white (hatched) regions
 356 coincide with those mass points which yield an initial (final) value of $\sqrt{g_q g_\chi}$ or $g_{q\chi}$ which
 357 fails to satisfy our requirement that $\Gamma < M_{\text{med}}/2$. (We observe that values for which the
 358 width is just within our upper validity bound of $M_{\text{med}}/2$ may be pushed over into the
 359 invalid range with the addition of new particles, not considered here, which would serve
 360 to increase the mediator width.) When $g_\chi/g_q = 0.2$, only the mono-jet channel produces
 361 limits which survive this requirement, and so these are shown separately in fig. 7.

362 Detailed comments specific to each channel are below, however some trends are channel-
 363 independent. For the sV model, strong limits exist when $M_{\text{med}} > 2m_\chi$ as the mediator
 364 can go on-shell, thereby enhancing the cross-section. The sA model limits show similar
 365 behaviour to the sV model except in the region where $m_\chi \gtrsim 10 \times M_{\text{med}}$, where **TO BE**
 366 **CONTINUED**. The upper limit on $\sqrt{g_q g_\chi}$ is relatively constant across values of g_χ/g_q , as
 367 is expected when the coupling (and hence the width) has been demonstrated to have little
 368 effect on kinematic behaviour, and using the assumptions of eq. 2.7. As the ratio increases,
 369 points in the region $M_{\text{med}} > m_\chi$ disappear as the initial condition, $g_q = 1$, leads to failure
 370 of the width condition. However, one could easily have chosen a smaller initial value of g_q
 371 to recover these points, and we suggest that the limits in this region would be quite similar
 372 to those seen in the $g_\chi/g_q = 0.2$ and 0.5 cases.

373 The results are limited when m_χ or M_{med} is large due to the smaller cross-sections in
 374 these cases, and are expected to improve at higher centre-of-mass energies. At small DM
 375 masses, the $E_{\text{T}}^{\text{miss}}$ distribution peaks at low values (with the exception of the tS model in
 376 the mono-jet channel), so results in this region of phase space are limited by statistical
 377 uncertainties from being in the tail of the distribution; this region would benefit from
 378 further optimisation of event selection in analyses aimed at SiM study, as we expect to see
 379 from the upcoming Run II results.

380 The results are discussed according to channel below.

381 4.1 Mono-jet channel

382 The upper limits on the coupling combination $\sqrt{g_q g_\chi}$ of the sV and sA models, obtained in
 383 the mono-jet channel, are displayed in the left-hand column of figs. 3-6, for $g_q/g_\chi = 0.5, 1,$
 384 2 and 5 respectively. Additional results for the $g_q/g_\chi = 0.2$ case are also shown separately
 385 in fig. 7, as these limits are only meaningful within this channel.

⁸We note that perturbativity is questionable for couplings greater than $\sim 2 - 3$, however 4π is an oft-quoted hard upper limit.

As expected, the mono-jet channel produces the strongest coupling limits in both s -channel models, which are better than those from the next-best mono- Z channel by a factor of a few. For the sV model, the weakest limits result for large m_χ or large M_{med} , and in fact are so weak that they are pushed into the region of invalidity where $\Gamma > M_{\text{med}}/2$; this is because although the acceptance is considerably higher in these regions compared to low masses, the cross section is sufficiently small that this effect prevails. Within the valid region ($m_\chi \in [1, 100]$ GeV and $M_{\text{med}} \in [1, 200]$ GeV), the limit on $\sqrt{g_q g_\chi}$ generally ranges from 0.1 to 0.7, with a handful of on-shell masses reaching a limit of ~ 0.05 in the large g_χ/g_q case. In the large g_χ/g_q scenario, limits for $m_\chi = 1000$ GeV start to become valid. This is because if $\sqrt{g_q g_\chi}$ remains constant but the g_χ/g_q increases then the value of g_q is pushed downward and so the width, which is dominated by decays to SM particles, decreases.

The sA model displays similar behaviour to the sV model except for the $m_\chi \gtrsim 10 \times M_{\text{med}}$ region, as described above.

The uncertainties on the limits in the sV model generally range from ??% to ??%, and are dominated by **X**. The uncertainties for the sA model are dominated by **X** and are within the range $X \sim Y\%$.

4.2 Mono- Z channel

The simplicity of the mono- Z channel relative to the mono-jet channel, and the speed of its production within MADGRAPH5, allowed us to study a finer granularity of points in the mass phase space. The resulting limits on the sV and sA models are shown in the central column of figs. 3-6. The behaviour of the limits as g_χ/g_q varies is similar to that within the mono-jet channel, and overall the limits are weaker compared to that channel by a factor of a few.

The total relative uncertainties on $\sqrt{g_q g_\chi}$ in the s -channel models are generally within 10%, but can range up to 80% in a few cases; they are dominated by **X**.

The advantage of the mono-boson channels is in the study of the tS model; since this was not included in the mono-jet channel the strongest limits are obtained with the mono- Z analysis, and are shown in the left-hand side of fig. 8. In general the tS model limits are weaker than the corresponding s -channel points (why?), note that the scale here is increased by a factor of 10. We find stronger limits for smaller m_χ and M_{med} masses, where larger cross sections compensate for lower acceptances at these points.

4.3 Mono- W/Z channel

The limits on the couplings of the sV , sA and tS models, obtained within the mono- W/Z channel, are shown in the right-hand column of figs. 3-8. This channel was studied to compare with the leptonic mono- Z channel in particular, but a coarser selection of masses was chosen as the limits were initially found to be somewhat weaker. Additionally, further estimates were made: a) as the kinematic behaviour is reasonable independent of the couplings, a single acceptance was found for each (m_χ, M_{med}) combination and applied to each value of g_χ/g_q , and b) complete systematic uncertainties were generated for a subset of masses and compared to those from the mono- Z channel; from this comparison the mono- Z

systematic uncertainties were multiplied by 3 and then applied to the mono- W/Z limits. As a result, the limits obtained in this channel are not intended to be rigorously quantitative; rather, they are used more to indicate qualitatively how the channel compares.

The ATLAS mono- W/Z analysis (and in particular the higher E_T^{miss} signal region) was not optimised for a simplified model interpretation, and much of the phase space produced insignificant numbers of events passing the event selection, with up to 200 thousand events generated. Generally, the limits are a factor of a few weaker again than those from the mono- Z channel, which is both consistent with the limits on the EFT models studied in the ATLAS analyses, and expected following our use of a cut-and-count interpretation of the mono- W/Z public results (can we say this? Point is to compare with them probably doing a shape analysis to improve their limits).

In some cases - most notably the $m_\chi \gtrsim 10 \times M_{\text{med}}$ region - the limits become comparable with the mono- Z channel, suggesting that more statistics and an improved treatment of systematic uncertainties would bring these closer in line with that channel.

Overall, the uncertainties from this channel lie within the range XXX and are dominated by Y.

5 Conclusion

5.1 Comparison with Relic Density Constraints


In Figs. ?? we show lines where the constraint on the coupling corresponds to the coupling strength that would reproduce the correct DM density if DM is a thermal relic of the early universe. For points diagonally above and to the left of the dashed line, the LHC constraints naively rule out the couplings leading to the correct relic density. Below and to the right of this line the relic density coupling is still allowed.

In this scenario, the measured abundance is approximately related to the unknown self-annihilation cross-section via


$$\Omega_{\text{DM}} h^2 \simeq \frac{2 \times 2.4 \times 10^{-10} \text{ GeV}^{-2}}{\langle \sigma v \rangle_{\text{ann}}}. \quad (5.1)$$

This is used with measurements of the DM abundance by Planck, $\Omega_{\text{DM}}^{\text{obs}} h^2 = 0.1199 \pm 0.0027$ [36], to find $\langle \sigma v \rangle_{\text{ann}} \simeq 4.0 \times 10^{-9} \text{ GeV}^{-2}$ for thermal relic DM. This relation is only approximately accurate, and so we use the Micromegas code [52] to determine the coupling strength leading to the correct relic density for each model. We verified this technique against the semi-analytic technique outlined in e.g. Ref. [37].


If the DM mass lies at the electroweak scale, the thermal relic scenario provides a natural explanation for the observed DM density, and so the coupling strengths leading to the correct relic density are a natural benchmark with which to compare constraints from other DM searches, indicating the scale at which we expect the couplings may lie. However the relic density couplings should by no means be treated as a constraint. If the DM was not produced thermally or if there is some unknown effect which modifies the evolution of the density with temperature, then these relations break down. Further, even if DM is a thermal relic, then the relationship no longer holds if there are other annihilation channels




(a) sV model, $g_q/g_\chi = 0.5$, mono-jet channel.




(b) sV model, $g_q/g_\chi = 0.5$, mono-Z channel.




(c) sV model, $g_q/g_\chi = 0.5$, mono- W/Z channel.



(d) sA model, $g_q/g_\chi = 0.5$, mono-jet channel.




(e) sA model, $g_q/g_\chi = 0.5$, mono-Z channel.




(f) sA model, $g_q/g_\chi = 0.5$, mono- W/Z channel.


Figure 3: Upper limits on the coupling for the s -channel models in the mono-jet (left), mono- Z (centre) and mono- W/Z (right) channels, for $g_\chi/g_q = 0.5$. The grey region represents the phase space where no meaningful limit was obtained. The hatched region represents a limit which leads to a width greater than $M_{\text{med}}/2$, so the validity of the calculation begins to fail. The dotted region represents phase space where insufficient statistics were available.




(a) sV model, $g_q/g_\chi = 1$, mono-jet channel.




(b) sV model, $g_q/g_\chi = 1$, mono-Z channel.




(c) sV model, $g_q/g_\chi = 1$, mono-W/Z channel.



(d) sA model, $g_q/g_\chi = 1$, mono-jet channel.




(e) sA model, $g_q/g_\chi = 1$, mono-Z channel.




(f) sA model, $g_q/g_\chi = 1$, mono-W/Z channel.


Figure 4: Upper limits on the couplings for the s-channel models in the mono-jet (left), mono-Z (centre) and mono-W/Z (right) channels, for $g_\chi/g_q = 1$. Refer to fig. 3 for details.




(a) sV model, $g_q/g_\chi = 2$, mono-jet channel.




(b) sV model, $g_q/g_\chi = 2$, mono- Z channel.




(c) sV model, $g_q/g_\chi = 2$, mono- W/Z channel.



(d) sA model, $g_q/g_\chi = 2$, mono-jet channel.




(e) sA model, $g_q/g_\chi = 2$, mono- Z channel.




(f) sA model, $g_q/g_\chi = 2$, mono- W/Z channel.


Figure 5: Upper limits on the coupling for the s -channel models in the mono-jet (left), mono- Z (centre) and mono- W/Z (right) channels, for $g_\chi/g_q = 2$. Refer to fig. 3 for details.




(a) sV model, $g_q/g_\chi = 5$, mono-jet channel.




(b) sV model, $g_q/g_\chi = 5$, mono- Z channel.




(c) sV model, $g_q/g_\chi = 5$, mono- W/Z channel.



(d) sA model, $g_q/g_\chi = 5$, mono-jet channel.



(e) sA model, $g_q/g_\chi = 5$, mono- Z channel.



(f) sA model, $g_q/g_\chi = 5$, mono- W/Z channel.

Figure 6: Upper limits on the coupling for the s -channel models in the mono-jet (left), mono- Z (centre) and mono- W/Z (right) channels, for $g_\chi/g_q = 5$. Refer to fig. 3 for details.




Figure 7: Upper limits on the coupling for the s -channel models in the mono-jet channel, for $g_\chi/g_q = 0.2$. Refer to fig. 3 for details.




Figure 8: Upper limits on the coupling $g_{q\chi}$ for the t -channel model in the mono- Z (left) and mono- W/Z (right) channels. Refer to fig. 3 for details.

not taken into account, or if there are other beyond-SM particles contributing to the DM abundance.

5.2 Comparison with Direct Detection Constraints

In Figs. ?? we also show the intercept line where constraints from direct detection experiments are equally as strong as the LHC constraint. Below and to the right of the dotted line, direct detection constraints are stronger than the LHC constraint, while above and to the left, the LHC gives the stronger constraint. We use the toolset from Ref. [?] to convert the strongest available direct detection constraints, which are from the LUX 2013 dataset [?], onto constraints on our models.

Compared to direct detection, the LHC performs relatively better for the SAD model than for the SVD model. This is because the axial-vector coupling leads to a suppressed scattering rate in direct detection experiments while the LHC is relatively insensitive to the difference between the vector and axial-vector couplings. In the non-relativistic limit, the TSD model leads to a mix of both suppressed and unsuppressed operators.

The direct detection constraints assume that the DM candidate under consideration contributes 100% of the local DM density, while the LHC constraints make no assumptions about either the local DM density or overall abundance. In this sense the LHC constraints remain useful even in the region where they are not as strong as those from direct detection.

5.3 Discussion

- Comparison to direct mediator searches: dijet gives strongest constraints on mediator especially for small r . Missing ET still good for large M but in this region EFT is fine
- Comparison to non-grid searches, e.g. McCullough et al
- Comparison to grid searches e.g. Zurek et al, Jacques and Nordstrom

MonoX searches dominate.

6 Acknowledgements

A Limit setting strategy

In this appendix we present a summary of the procedure employed to calculate the 95% confidence level (CL) limits on the coupling parameter $\sqrt{g_q g_\chi}$, where this parameter can be replaced with $g_{q\chi}$ for the tS model, and M_\star in the validation of the mono-jet analysis.

A.1 Nominal Values

For each SiM, the nominal limit is calculated by taking the model-independent upper limit on $\sigma \times \mathcal{A} \times \epsilon$ from each analysis, dividing by the value of $\mathcal{A} \times \epsilon$ (which is taken as a single parameter for each point) to obtain the limiting cross section σ_{lim} , and rearranging eq. 2.7 to convert to a limit on the couplings. In the s -channel, on-shell case, the width can be

expressed as a function of g_q and the ratio g_χ/g_q , which simplifies the calculation. We arrive at

$$\sqrt{g_q g_\chi} \lim = \begin{cases} \sqrt{g_q g_\chi} \times (\sigma_{\text{lim}}/\sigma_{\text{gen}})^{\frac{1}{2}} & \text{if } M_{\text{med}} \geq 2m_{\text{DM}} \text{ (} s\text{-channel)} \\ \sqrt{g_q g_\chi} \times (\sigma_{\text{lim}}/\sigma_{\text{gen}})^{\frac{1}{4}} & \text{if } M_{\text{med}} < 2m_{\text{DM}} \end{cases} \quad (\text{A.1})$$

where $\sqrt{g_q g_\chi}$ and σ_{gen} are the input couplings and resulting cross-section (taken from PYTHIA 8), respectively.

The signal region in each case is chosen based on where the best ‘expected’ limit exists, where that limit is calculated assuming that exactly the expected SM background is observed.

A.2 Uncertainty Estimation

Our nominal limits rely on both σ_{gen} and $\mathcal{A} \times \epsilon$ and so are subject to systematic uncertainties which derive from our choice of MC generation procedure. For our MC samples, there are three key sources of systematic uncertainty: the factorisation and renormalisation scales, the strong coupling constant (α_s) and the choice of parton distribution function (PDF). *Should we say something about the actual choice of generator here? And what about LO vs NLO?*

The factorisation and renormalisation default scales are straightforward to test in MADGRAPH5 where they varied simultaneously by factors of 2 (‘up’) and 0.5 (‘down’). The systematic effects of the strong coupling constant and the PDF are difficult to separate and so are treated in tandem. We assume that the systematic uncertainty introduced by α_s at matrix-element level is negligible when compared to the PDF uncertainties, as demonstrated to be valid in ref. [42]. The variation of α_s in conjunction with a PDF is done with the use of specific tunes in PYTHIA 8, which we change simultaneously with the PDF choice to estimate the uncertainty on $\Delta\sigma_{\text{gen}}$. The nominal choices of PDF and tune are varied ‘up’ to NNPDF2.1LO PDF + Monash tune, and ‘down’ to CTEQ6L1 PDF and ATLAS UE AU2-CTEQ6L1 tune. *Millie: put discussion of matching scale systematic here.* These systematic uncertainty sources are summarised in table 5.

The average variation from the nominal value of σ_{lim} , as a fraction of σ_{lim} , resulting from each systematic source is added in quadrature and propagated to $\sqrt{g_q g_\chi}$ to obtain the total systematic uncertainty. This process is adjusted slightly to account for the inclusion of statistical uncertainties, which are estimated rather conservatively by taking the 95% CL *lower* limit on $\mathcal{A} \times \epsilon$ as calculated with the Wald approximation, i.e. $\mathcal{A} \times \epsilon \rightarrow (\mathcal{A} \times \epsilon) - \Delta(\mathcal{A} \times \epsilon)$. The uncertainty on the luminosity is less than 3%, so is considered to be negligible in comparison to other systematic sources.

B Validation of signal simulation and event selection procedures

B.1 Monojet Channel

The MC generation and signal selection procedures for the mono-jet channel are validated via reproduction of the ATLAS limits on $M_\star \equiv M_{\text{med}}/\sqrt{g_q g_\chi}$, for the s -channel vector

main systematic sources	PDF/tune	factorisation and renormalisation scales	matching scale (mono-jet only)
variation ‘up’	NNPDF2.1LO + Monash tune	2	160 GeV
nominal	MSTW2008lo68cl + ATLAS UE AU2-MSTW2008LO	1	80 GeV
variation ‘down’	CTEQ6L1 + ATLAS UE AU2-CTEQ6L1	0.5	40 GeV

Table 5: The sources of systematic uncertainty considered in this analysis. Each point in phase space is varied up or down by one of these sources, and the systematic uncertainty is taken to be the average difference in modified acceptance from the nominal value.

simplified model. A comparison of SR7 limits for a representative sample of mediator masses with $m_\chi = 50$ GeV, $\Gamma = M/8\pi$ and $\sqrt{g_q g_\chi} = 1$ is presented in table 6. In general, good agreement is observed between the ATLAS and reproduced limits, with a maximum difference (with respect to the ATLAS limit) of $<23\%$. We note that a discrepancy such as this is expected and allowed for three reasons. Firstly, the MC generation procedure employed in this analysis does not include a full simulation of the ATLAS detector. Instead, reconstruction effects are simulated by applying a Gaussian smearing of the jet p_T . Secondly, the matching procedure employed in this analysis (and discussed in detail in Section 3.1.1) is largely simplified. This introduces a substantial uncertainty when compared to the matching procedure utilised by the ATLAS mono-jet group. For example, where the ATLAS group observe a maximum matching scale uncertainty of 5% for events with E_T^{miss} above 350 GeV, we observe an uncertainty of $\sim 30\%$. Lastly, the 95% CL uncertainties on M_\star for this work are estimated as described in appendix A, while the ATLAS analysis used a more complex HistFitter treatment (confirm). As our results are consistently more conservative than those of the ATLAS analysis, we consider this approach acceptable.

B.2 Mono-Z Channel

The ATLAS mono- Z analysis result includes an upper limit on the coupling $g_{q\chi}$ for a t -channel simplified model that is very similar to the tS model investigated here, and so is used for validating our signal generation and selection procedure. The most significant differences are in the number of mediating particles — the ATLAS model includes just two mediators (up - and $down$ -type) compared to our six — and in the nature of the DM particle, which is taken to be Majorana. Additionally, while we use a universal coupling $g_{q\chi}$ to all three quark generations, the analysis used a model which set $g_{t,b\chi} = 0$.

M [TeV]	M_\star^{95} [GeV] (ATLAS)	M_\star^{95} [GeV] (this work)	Difference [%]
0.05	91	89	2.16
0.3	1151	1041	7.3
0.6	1868	1535	11.8
1	2225	1732	12.0
3	1349	1072	6.8
6	945	769	8.5
10	928	724	10.6
30	914	722	9.6

Table 6: Comparison of the 95% CL upper limits on M_\star from this work and from the ATLAS mono-jet analysis [41]. The limits are compared for an s -channel model with $m_\chi = 50$ GeV and Γ set to a fixed width of $M_{\text{med}}/8\pi$, for the process $pp \rightarrow \chi\bar{\chi} + 1, 2j$ with QCUT = 80 GeV.

m_χ [GeV]	M_{med} [GeV]	$g_{q\chi}^{95\%CL}$ (ATLAS)	$g_{q\chi}^{95\%CL}$ (this work)	Difference [%]
10	200	1.9	2.0	5.3
	500	2.8	3.2	14.3
	700	3.5	4.4	25.7
	1000	4.5	5.2	15.6
200	500	3.4	4.0	17.6
	700	4.2	4.5	7.1
	1000	5.2	5.3	1.9
400	500	5.5	5.7	3.6
	700	6.1	6.5	6.6
	1000	7.2	7.4	2.8
1000	1200	23.3	24.1	3.4

Table 7: Comparison of the upper limit on $g_{q\chi}$ from the ATLAS analysis [47] and this work.

Table 7 shows the 95% CL upper limits on $g_{q\chi}$ that we calculate using the same t -channel model and our own generation procedure (using the values in table 3), compared with the limits on this same variable taken from the ATLAS analysis. The difference as a percentage of the ATLAS limit is also shown in the table. We see reasonable agreement; most of the 11 points in parameter space are within 10% of the ATLAS limits, and all are within 26%. Additionally, our results are consistently more conservative, which is to be expected due to the less sophisticated nature of our generation procedure. Similarly to the mono-jet validation, the dominant effects are due to the use of p_T smearing applied

m_χ [GeV]	$M_\star^{90\%CL}$ [GeV] (ATLAS)	$M_\star^{95\%CL}$ [GeV] (this work)	Difference [%]
1	570	499	12.5

Table 8: Comparison of the upper limit on M_\star from the ATLAS mono- W/Z analysis [48] and this work.

to the leptons, rather than considering the full reconstruction effects, and the simplified treatment of systematics; we also obtained $\sigma \times \mathcal{A} \times \epsilon$ independently.

B.3 Mono- W/Z Channel

The signal generation and event selection for the mono- W/Z channel are validated by reproducing the published limit on M_\star for the D5 EFT operator with $m_\chi = 1$ GeV. We see agreement within 12.5%, where our limit is slightly more conservative. This is to be expected for the following reasons: the D5 limits are only published for the low- E_T^{miss} signal region, whereas we use only the high- E_T^{miss} signal region in this recast; the ATLAS analysis uses a shape fit to extract the limit on the number of new physics events, whereas we use a cut-and-count approach; and the ATLAS limit is quoted at 90% CL, while ours is calculated at 95% CL.

References

- [1] ATLAS Collaboration, *Search for new phenomena with the monojet and missing transverse momentum signature using the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions*, *Phys. Lett. B* (2011), arXiv:1106.5327.
- [2] ATLAS Collaboration, *Search for New Phenomena in Monojet plus Missing Transverse Momentum Final States using 10 fb^{-1} of pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector at the LHC*, 2012, ATLAS-CONF-2012-147.
- [3] CMS Collaboration, *Search for new physics in monojet events in pp collisions at $\sqrt{s} = 8$ TeV*, 2013, CMS-PAS-EXO-12-048.
- [4] M. R. Buckley, *Using Effective Operators to Understand CoGeNT and CDMS-Si*, *Phys.Rev. D* 88, 055028 (2013), arXiv:1308.4146.
- [5] J. Abdallah et al., *Search for new phenomena with mono-jet plus missing transverse energy signature in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector*, 2012, ATL-COM-PHYS-2012-1211.
- [6] N. Bell et al., *Searching for Dark Matter at the LHC with a Mono-Z*, *Phys.Rev. D* 86, 096011 (2012), arXiv:1209.0231.
- [7] N. Zhou, D. Berge, and D. Whiteson, *Mono-everything: combined limits on dark matter production at colliders from multiple final states*, *Phys.Rev. D* 87, 095013 (2013), arXiv:1302.3619.
- [8] M. Cahill-Rowley et al., *Complementarity and Searches for Dark Matter in the $p\text{MSSM}$* , SLAC-PUB-15450 (2013), arXiv:1305.6921.

- [9] ATLAS Collaboration, *Further search for supersymmetry at $\sqrt{s} = 7$ TeV in final states with jets, missing transverse momentum and isolated leptons with the ATLAS detector*, *Phys.Rev. D* 86, 092002 (2012), arXiv:1208.4688.
- [10] ATLAS Collaboration, *Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collision data*, *Phys.Rev. D* 87, 012008 (2013), arXiv:1208.0949.
- [11] ATLAS Collaboration, *Search for pair-produced third-generation squarks decaying via charm quarks or in compressed supersymmetric scenarios in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys.Rev. D* 90, 052008 (2014), arXiv:1407.0608.
- [12] ATLAS Collaboration, *Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton-proton collision data*, *JHEP* 09 (2014) 146, arXiv:1405.7875.
- [13] H. Dreiner et al., *Contact Interactions Probe Effective Dark Matter Models at the LHC*, *Europhys.Lett.* (2013), arXiv:1303.3348.
- [14] J. Goodman et al., *Gamma Ray Line Constraints on Effective Theories of Dark Matter*, *Nucl.Phys.* (2011), arXiv:1009.0008.
- [15] G. Busoni et al., *On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC*, *Phys.Lett.* (2014), arXiv:1307.2253.
- [16] G. Busoni et al., *On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC, Part II: Complete Analysis for the s -channel*, *JCAP* 1406:060 (2014), arXiv:1402.1275.
- [17] G. Busoni et al., *On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC Part III: Analysis for the t -channel*, *JCAP* 09 (2014) 022, arXiv:1405.3101.
- [18] Oliver Buchmueller, Matthew J. Dolan, Sarah A. Malik and Christopher McCabe, *Characterising dark matter searches at colliders and direct detection experiments: Vector mediators*, 2014, arXiv:1407.8257.
- [19] J. Kumar and D. Marfatia, *Matrix element analyses of dark matter scattering and annihilation*, *Phys.Rev.* (2013), arXiv:1305.1611.
- [20] G. Jungman et al., *Supersymmetric dark matter*, *Phys.Rept.* (1996).
- [21] P. J. Fox et al., *Missing Energy Signatures of Dark Matter at the LHC*, *Phys.Rev.* (2012), arXiv:1109.4398.
- [22] P. J. Fox, R. Harnik, R. Primulando, and C-T. Yu, *Taking a Razor to Dark Matter Parameter Space at the LHC*, *Phys.Rev.* (2012), arXiv:1203.1662.
- [23] M. Papucci, A. Vichi, and K. M. Zurek, *Monojet versus rest of the world I: t -channel Models*, *JHEP* (2014), arXiv:1402.2285.
- [24] Y. Bai, P. J. Fox, and R. Harnik, *The Tevatron at the Frontier of Dark Matter Direct Detection*, *JHEP* (2010), arXiv:1005.3797.
- [25] J. Goodman et al., *Constraints on Dark Matter from Colliders*, *Phys.Rev. D* 82, 116010 (2010), arXiv:1008.1783.
- [26] P. J. Fox, R. Harnik, J. Kopp, and Y. Tsai, *LEP Shines Light on Dark Matter*, *Phys.Rev.* (2011), arXiv:1103.0240.

- [27] M. L. Graesser, I. M. Shoemaker, and L. Vecchi, *A Dark Force for Baryons*, 2011, arXiv:1107.2666.
- [28] H. An and F. Gao, *Fitting CoGeNT Modulation with an Inelastic, Isopin-Violating Z' Model*, 2011, arXiv:1108.3943.
- [29] CMS Collaboration, *Search for narrow resonances using the dijet mass spectrum in pp collisions at $\sqrt{s} = 8\text{TeV}$* , *Phys.Rev.* (2013), arXiv:1302.4794.
- [30] ATLAS Collaboration, *Search for high-mass resonances decaying to dilepton final states in pp collisions at $\sqrt{s} = 7\text{-TeV}$ with the ATLAS detector*, *JHEP* (2012), arXiv:1209.2535.
- [31] P. Harris, V. V. Khoze, M. Spannowsky and C. Williams, *Constraining Dark Sectors at Colliders: Beyond the Effective Theory Approach*, *Phys.Rev.* (2015), arXiv:1411.0535.
- [32] CMS Collaboration. *Search for new physics in monojet events in pp collisions at $\sqrt{s} = 8\text{TeV}$* , 2013, CMS-PAS-EXO-12-048.
- [33] ATLAS Collaboration. *Search for New Phenomena in Monojet plus Missing Transverse Momentum Final States using 10 fb^{-1} of pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector at the LHC*, 2012, ATLAS-CONF-2012-147.
- [34] J. Kumar and D. Marfatia, *Matrix element analyses of dark matter scattering and annihilation*, *Phys.Rev.* (2013), arXiv:1305.1611.
- [35] D. Alves et al., *Simplified Models for LHC New Physics Searches*, *J.Phys.* (2012), arXiv:1105.2838.
- [36] P. A. R. Ade *et al.* [Planck Collaboration], *Astron. Astrophys.* **571**, A16 (2014) [arXiv:1303.5076 [astro-ph.CO]].
- [37] G. Busoni, A. De Simone, T. Jacques, E. Morgante and A. Riotto, *Making the Most of the Relic Density for Dark Matter Searches at the LHC 14 TeV Run*, *JCAP* 03 (2015) 022, arXiv:1410.7409.
- [38] CMS Collaboration. *Search for new physics in monojet events in pp collisions at $\sqrt{s} = 8\text{TeV}$* , 2013, CMS-PAS-EXO-12-048.
- [39] ATLAS Collaboration. *Search for New Phenomena in Monojet plus Missing Transverse Momentum Final States using 10 fb^{-1} of pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector at the LHC*, 2012, ATLAS-CONF-2012-147.
- [40] ATLAS Collaboration. *Further search for supersymmetry at $\sqrt{s} = 7\text{ TeV}$ in final states with jets, missing transverse momentum and isolated leptons with the ATLAS detector*, *Phys.Rev.* (2012), arXiv:1208.4688.
- [41] ATLAS Collaboration. *Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector*, 2015, arXiv:1502.01518
- [42] S. Schramm, *Searching for Dark Matter with the ATLAS Detector in Events with an Energetic Jet and Large Missing Transverse Momentum*, 2015, CERN-THESIS-2015-038.
- [43] A. Cooper-Sarkar. *PDFs for the LHC*, 2011, arXiv:1107.5170.
- [44] ATLAS Collaboration. *Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the ATLAS detector*, 2013, CERN-PH-EP-2012-210, arXiv:1210.4491.
- [45] P. J. Fox et al. *Missing Energy Signatures of Dark Matter at the LHC*, *Phys. Rev.*, 2012.

- [46] N. Bell, J. Dent, T. Jacques, and T. Weiler. *W/Z Bremsstrahlung as the Dominant Annihilation Channel for Dark Matter*, *Phys. Rev.*, 2011.
- [47] ATLAS Collaboration. *Search for dark matter in events with a Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys.Rev.D* **90** (2014) 012004, arXiv:1404.0051.
- [48] ATLAS Collaboration. *Search for dark matter in events with a hadronically decaying W or Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. Lett.* **112** (2014) 041802, arXiv:1309.4017.
- [49] J. Alwall /emphet al.. *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP07* (2014) 079, arXiv:1405.0301.
- [50] A. D. Martin, W. J. Stirling, R. S. Thorne, G. Watt, *Parton distributions for the LHC*, *Eur.Phys.J.C63*, (2009), 189-285, arXiv:0901.0002.
- [51] ATLAS Collaboration. *Summary of ATLAS Pythia8 tunes*, 2012, ATL-PHYS-PUB-2012-003.
- [52] G. Blanger, F. Boudjema, A. Pukhov and A. Semenov, *Comput. Phys. Commun.* **192**, 322 (2015) doi:10.1016/j.cpc.2015.03.003 [arXiv:1407.6129 [hep-ph]].
- [53] D. Abercrombie et al. *Dark Matter Benchmark Models for Early LHC Run-2 Searches: Report of the ATLAS/CMS Dark Matter Forum*, arXiv:1507.00966.
- [54] M. Baak et al. *HistFitter software framework for statistical data analysis*, *Eur.Phys.J.C* (2015), arXiv:1410.1280.
- [55] J. Abdallah et al. *Simplified Models for Dark Matter Searches at the LHC*, *Phys. Dark Uni.* 9-10 (2015) 8-23, arXiv:1506.03116.
- [56] J. Abdallah et al. *Simplified Models for Dark Matter and Missing Energy Searches at the LHC*, (2014), arXiv:1409.2893.
- [57] T. Jacques and K. Nordstrom, *Mapping monojet constraints onto Simplified Dark Matter Models*, *JHEP* 06 (2015) 142, arXiv:1502.05721.
- [58] T. Sjstrand et al. *An Introduction to PYTHIA 8.2*, *Comput. Phys. Comm.* 191 (2015) 159-177, arXiv:1410.3012.