

3 Collide and Conquer: Constraints on Simplified Dark 4 Matter Models using Mono- X Collider Searches

5 **A. J. Brennan,^{a,1} M. F. McDonald,^a J. Gramling^b and T. D. Jacques^c**

6 ^a*The University of Melbourne, Parkville 3010, Australia*

7 ^b*Université de Genève, Quai E. Ansermet 24, 1211 Genève 4, Switzerland*

8 ^c*SISSA/ISAS, via Bonomea 265, 34136 Trieste, Italy*

9 *E-mail:* amelia.brennan@coepp.org.au, millie.mcdonald@coepp.org.au,
johanna.gramling@cern.ch, thomas.jacques@sissa.it

10 ABSTRACT: The use of simplified models as a tool for interpreting dark matter collider
11 searches has become increasingly prevalent, and while early Run II results are beginning to
12 appear, we look to see what further information can be extracted from the Run I dataset.
13 We consider three ‘standard’ simplified models that couple quarks to fermionic singlet dark
14 matter: an s -channel vector mediator with vector or axial-vector couplings, and a t -channel
15 scalar mediator. Upper limits on the couplings are calculated and compared across three
16 alternate channels, namely mono-jet, mono- Z (leptonic) and mono- W/Z (hadronic). The
17 strongest limits are observed in the mono-jet channel, however the computational simplicity
18 and absence of significant t -channel model width effects in the mono-boson channels make
19 these a straightforward and competitive alternative. We also include a comparison with
20 relic density and direct detection constraints.

¹Corresponding author.

21	Contents	
22	1 Introduction	2
23	2 Simplified model phenomenology	3
24	2.1 Model descriptions	3
25	2.2 The mono- $X + E_{\text{T}}^{\text{miss}}$ signatures	4
26	2.3 Mass and coupling points	5
27	2.4 Treatment of the width	5
28	3 Recasting mono-X constraints	7
29	3.1 Signal simulation	9
30	3.1.1 Parton matching scheme	9
31	3.2 Mono-jet constraints	9
32	3.3 Mono- $Z(\text{lep})$ constraints	10
33	3.4 Mono- $W/Z(\text{had})$ constraints	11
34	4 Results and discussion	12
35	4.1 Limits on the couplings $\sqrt{g_q g_\chi}$	12
36	4.1.1 Mono-jet channel	13
37	4.1.2 Mono- $Z(\text{lep})$ channel	14
38	4.1.3 Mono- $W/Z(\text{had})$ channel	14
39	4.2 Comparison with relic density constraints	15
40	4.3 Comparison with direct detection constraints	15
41	5 Conclusion	21
42	6 Acknowledgements	22
43	A Limit setting strategy	22
44	A.1 Nominal values	22
45	A.2 Uncertainty estimation	22
46	B Validation of signal simulation and event selection procedures	24
47	B.1 Mono-jet channel	24
48	B.2 Mono- $Z(\text{lep})$ channel	24
49	B.3 Mono- $W/Z(\text{had})$ channel	25

1 Introduction

Simplified models have emerged as a powerful tool for the interpretation of collider, direct and indirect detection signals of dark matter (DM). Previously, ATLAS and CMS searches for DM were conducted within the context of both Effective Field Theories (EFTs) [1–4] and full UV-complete theories such as Supersymmetry [5–8]. The latter approach, though well-motivated, is typified by a broad parameter space and generally yields results which are insensitive to the wider class of DM models. EFT constraints, in comparison, are applicable to a broad range of models and rely on the specification of only a small set of parameters, namely the suppression scale, M_\star , and the DM mass, m_{DM} [9]. In the EFT framework, interactions between the dark and Standard Model (SM) sector are parametrised by a set of higher-dimensional effective operators, that arise when the mass of the mediating particle is assumed to be significantly larger than the momentum transferred in a given interaction. Where this condition is not fulfilled, the EFT prescription can produce constraints which detour dramatically from those of the associated UV-complete model [9–13]. This is less important in direct detection experiments where the momentum transferred in the scattering of DM particles with heavy nuclei is generally of the order of tens of MeV [14, 15], or in indirect searches where the annihilations of non-relativistic DM particles in the galactic halo occur with momentum transfers of order m_{DM} . However, for hadron collider searches, where the accessible center of mass energy of two colliding baryons may be sufficient to produce the mediator on-shell, the range of validity of the EFT prescription is significantly diminished. Indeed, recent works have quantitatively shown that the EFT approach can break down as a valid interpretation of data collected during the $\sqrt{s} = 8$ TeV Run I of the Large Hadron Collider (LHC) [16–19]. In light of this, simplified models have become the preferred tool for the interpretation of collider DM searches [20–24].

In a nutshell, a simplified model (SiM) arises when the heavy mediator which was integrated out in the EFT framework is reintroduced. Like EFTs, SiMs admit the comparison of results obtained in the different avenues of dark matter study [25, 26] and are defined by a relatively small set of parameters - namely m_{DM} , the mass of the mediator M_{med} , and the SM-mediator and DM-mediator coupling strengths, g_q and g_χ (or $g_{q\chi}$ in the case of a single, SM-DM-mediator coupling). Unlike EFTs, constraints calculated within the context of a SiM are valid across a broad energy range.

In this paper, we examine a phenomenologically distinct set of SiMs. In particular, we place constraints on the SiMs corresponding to the simplest UV-completions of the D5 (vector) and D8 (axial-vector) effective operators in the s -channel¹. We also include a case in which a scalar mediator is exchanged in the t -channel, motivated by its analog of squark exchange in Supersymmetry. In the heavy mediator limit, the operator describing this model can be rearranged via a Fierz transformation into a combination of operators D5 to D8.

¹The D5 and D8 operators form a nice starting point in the analysis of SiMs as they have been studied exhaustively in the past (see refs. [1–4, 17–19, 27, 28] among others). This attention is motivated by the fact that collider limits for the D5 (D8) operator can be readily transformed into limits on spin-independent (spin-dependent) DM-nucleon scattering and vice versa.

88 The models are constrained using public results from mono- X + missing transverse
 89 energy (E_T^{miss}) searches conducted by the ATLAS Collaboration. Specifically, we focus on
 90 searches where X is either a parton (manifesting in the detector as a narrow-radius jet),
 91 a leptonically-decaying Z boson (manifesting as two opposite-sign same-flavor leptons), or
 92 a hadronically-decaying W or Z boson (manifesting as a large-radius jet). The purpose of
 93 this work is to strengthen existing SiM limits using the full 20.3 fb^{-1} of Run I ATLAS
 94 data, and to explore an enhanced phase space with respect to the mediator and DM masses
 95 and the relative strength of the couplings to the visible and dark sectors. We choose to
 96 treat the mediator width as the minimal value naturally arising, which is more realistic
 97 than a fixed width. We extend the study by providing a cross-check and comparison of the
 98 performance of the three targeted collider detection channels, and compare against relic
 99 density and direct detection constraints.

100 The remainder of the paper is organised as follows. Section 2 contains a compendium of
 101 the SiMs chosen for analysis and the associated collider phenomenology. Section 3 outlines
 102 the techniques used to recast mono- X + E_T^{miss} limits on the visible cross-section for any
 103 new physics process into constraints on SiMs, and specifically on the couplings g_q and g_χ .
 104 Lastly, our results are presented in section 4 along with a discussion of the implications of
 105 this work. Appendices A and B include details of the limit setting and analysis validation
 106 procedures.

107 2 Simplified model phenomenology

108 2.1 Model descriptions

109 We begin with a short set of assumptions: that the DM particle, χ , is a weakly interacting
 110 Dirac fermion, that it is a singlet under the SM, and that it is the lightest stable new
 111 particle. Additionally, the new sector is assumed to couple only to the SM quarks. While
 112 possible couplings to SM leptons (e.g. [11]) or gluons (e.g. [29]) have been studied elsewhere,
 113 they are beyond the scope of this paper. The nature of the mediating particle then results
 114 from these assumptions: in the s -channel it is chosen to be a vector particle which must also
 115 be a SM singlet, denoted ξ , while in the t -channel it is a scalar particle which is necessarily
 116 charged and coloured, and labelled ϕ .

117 The s -channel models chosen for analysis are Z' -type models characterised by vector
 118 (sV) or axial-vector (sA) couplings to both the dark and SM sectors. They are described
 119 by the following interaction Lagrangians:

$$\mathcal{L}_{sV} \supset -\xi_\mu \left[\sum_q g_q \bar{q} \gamma^\mu q - g_\chi \bar{\chi} \gamma^\mu \chi \right], \quad (2.1)$$

$$\mathcal{L}_{sA} \supset \xi_\mu \left[\sum_q g_q \bar{q} \gamma^\mu \gamma_5 q - g_\chi \bar{\chi} \gamma^\mu \gamma_5 \chi \right], \quad (2.2)$$

121 where the sum is over all quarks. This is a simple extension of the standard model and has
 122 been studied extensively [23, 24, 30–47]. For the couplings g_q and g_χ to remain within the

123 perturbative regime, they are required to satisfy $g_q, g_\chi \leq 4\pi$, though stronger perturbativity
 124 requirements do exist [17].

125 The t -channel model (abbreviated tS) is primarily motivated by analogy with a com-
 126 mon aspect of Supersymmetric models: neutralino DM interacting with the SM sector via
 127 t -channel exchange of a squark [48], and has been studied in the context of the LHC by
 128 a number of groups [25, 45, 49–55]. Note that in this Supersymmetric scenario the DM
 129 particle is a Majorana fermion. The collider phenomenology of a t -channel SiM in which
 130 χ is of Majorana type is kinematically identical to the corresponding Dirac case (requiring
 131 multiplication of the cross-section by a simple factor in order to compute limits) and so
 132 Majorana DM is not covered here².

133 In the tS model, the mediator is allowed to couple to either the left or right-handed
 134 quarks as an SU(2) doublet or singlet respectively. Since the LHC is insensitive to the
 135 chirality of the quarks, we assume for simplicity that ϕ couples to the left-handed quarks
 136 only, and is itself an SU(2) doublet, allowing radiation of a W boson. To avoid different
 137 couplings to quarks of different generations, and to remain in step with the DM forum
 138 recommendations [22], we include three generations of mediator doublets ϕ_i , with equal
 139 masses and couplings. The interaction Lagrangian for this model is then:

$$\mathcal{L}_{tS} \supset \sum_i g_{q\chi} \bar{Q}_i P_R \phi_i \chi + \text{h.c.}, \quad (2.3)$$

140 where the sum is over the three quark doublets, $g_{q\chi}$ is the DM-quark coupling (equal for
 141 each generation), and P_R is the usual chiral projection operator.

142 2.2 The mono- $X + E_T^{\text{miss}}$ signatures

143 The mono- $X + E_T^{\text{miss}}$ signal (abbreviated to mono- X) is a popular collider signal in the
 144 search for new physics, particularly in the search for dark matter. Since DM particles are
 145 not expected to interact with detector material, they appear as missing transverse energy
 146 when balanced against a visible object, X , that is radiated from the initial or intermediate
 147 state. For the s -channel SiMs discussed above, only initial-state radiation is permitted; see
 148 figs. 1a and 1b for examples. For the tS model, radiation of a gluon or electroweak (EW)
 149 boson is permitted both from initial state partons (fig. 1c) or from the mediator (fig. 1d).

150 The most likely scenario at the LHC is production of a jet alongside the invisible χ pair,
 151 as a result of the strong coupling and prevalence of partons in the initial state. However,
 152 to fully exploit the potential of the ATLAS detector to record and identify a vast array of
 153 particle types, we also consider two additional channels. Firstly, we take advantage of the
 154 relative cleanliness and simplicity of leptons in the leptonically-decaying mono- $(Z \rightarrow \ell^+ \ell^-)$
 155 channel. We also take advantage of the large hadronic branching fraction, and developing
 156 jet-identification techniques for boosted EW bosons, in the hadronically-decaying mono-
 157 $(W/Z \rightarrow jj)$ channel³. In both cases, the large multi-jet background is reduced, and
 158 complications in jet production such as parton-matching can be ignored, making these an

²The exception being in the validation of the mono- $Z(\text{lep})$ channel, see Sec. B.2.

³In addition, one of the first Run II dark matter search results from ATLAS was from this channel [56], released during the preparation of this paper.




Figure 1: Representative dark matter pair-production processes with a gluon or W/Z boson in the final state for the s -channel (a,b) and t -channel (c,d) models.

interesting alternative to the mono-jet channel where speed, efficiency and a reduction in jet-associated uncertainties may make up for a loss in sensitivity.

2.3 Mass and coupling points

A representative set of dark matter and mediator masses, listed in table 1, are chosen for study in each detection channel. DM masses of 3, 30 and 300 GeV are also included in the mono- Z (lep) channel, where ease of production permits higher granularity. All (m_χ, M_{med}) combinations are allowed in the sV and sA models, while in the tS model M_{med} must be greater than m_χ to ensure stability of the DM particle. The couplings g_q and $g_{q\chi}$ are set to unity, while the DM-mediator coupling in the s -channel models, g_χ , is varied from 0.2 to 5. The mediator masses are chosen to cover a broad range of parameter space and to coincide with predominantly three regimes: (near-)degenerate ($M_{\text{med}} \approx m_\chi$), on-shell ($M_{\text{med}} \geq 2m_\chi$) and off-shell ($M_{\text{med}} < 2m_\chi$).

2.4 Treatment of the width

An important factor when considering SiMs is to ensure that the mediator width is treated appropriately, as it impacts both the cross-section calculation and, in some cases, the kinematic behaviour of the model.

Following the DM Forum recommendations [22], we use the minimal width, allowing coupling to all kinematically accessible quarks. We assume minimal flavour violation, which

m_χ [GeV]	M_{med} [GeV]	s -channel		t -channel
		g_q	g_χ	$g_{q\chi}$
1, (3), 10, (30), 100, (300), 1000	1, 2, 10, 20, 100, 200, 1000, 2000	1	0.2, 0.5, 1, 2, 5	1

Table 1: Mass and coupling points chosen for the analysis of simplified dark matter models. Values in brackets are only included in the mono- $Z(\text{lep})$ channel. The mediator masses are primarily representative of three regimes: (near-)degenerate ($M_{\text{med}} \approx m_\chi$), on-shell ($M_{\text{med}} \geq 2m_\chi$) and off-shell ($M_{\text{med}} < 2m_\chi$). For the t -channel model, $M_{\text{med}} > m_\chi$ is required to ensure stability of the DM particle.

implies a universal coupling to all quark flavours. The minimum width for each model is given by:

$$\begin{aligned} \Gamma_{sV} = & \frac{g_\chi^2 M}{12\pi} \left(1 + \frac{2m_\chi^2}{M^2}\right) \left(1 - \frac{4m_\chi^2}{M^2}\right)^{\frac{1}{2}} \Theta(M - 2m_\chi) \\ & + \sum_q \frac{g_q^2 M}{4\pi} \left(1 + \frac{2m_q^2}{M^2}\right) \left(1 - \frac{4m_q^2}{M^2}\right)^{\frac{1}{2}} \Theta(M - 2m_q) \end{aligned} \quad (2.4)$$

$$\begin{aligned} \Gamma_{sA} = & \frac{g_\chi^2 M}{12\pi} \left(1 - \frac{4m_\chi^2}{M^2}\right)^{\frac{3}{2}} \Theta(M - 2m_\chi) \\ & + \sum_q \frac{g_q^2 M}{4\pi} \left(1 - \frac{4m_q^2}{M^2}\right)^{\frac{3}{2}} \Theta(M - 2m_q) \end{aligned} \quad (2.5)$$

$$\begin{aligned} \Gamma_{tS} = & \sum_q \frac{g_{q\chi}^2 M}{16\pi} \left(1 - \frac{m_q^2}{M^2} - \frac{m_\chi^2}{M^2}\right) \\ & \times \sqrt{\left(1 - \frac{m_q^2}{M^2} + \frac{m_\chi^2}{M^2}\right)^2 - 4 \frac{m_\chi^2}{M^2}} \Theta(M - m_q - m_\chi) \end{aligned} \quad (2.6)$$

It is possible that the mediator may decay to other SM or BSM particles [23], but this is not expected to have a large effect on the kinematic distribution as long as the width remains relatively small [22]. The generator treatment of the mediator as a Breit-Wigner propagator, rather than a true kinetic propagator, breaks down for large widths [36, 44].

We can take advantage of the fact that for each point in (m_χ, M_{med}) phase space, the mediator width (and therefore the couplings) do not greatly affect a model's kinematic behaviour (with the notable exception of the tS model in the mono-jet channel). This is demonstrated in fig. 2, where we plot a simplified E_T^{miss} distribution (as a proxy for the full selection in each analysis) for the sV (representing both the sV and sA models) and tS models for two mass points and a demonstrative set of couplings such that $\Gamma < M_{\text{med}}/2$. The E_T^{miss} distribution is predominantly independent of the mediator width for the s -

channel models in the mono-jet channel, and all models in the mono- $Z(\text{lep})$ ⁴ channel. However, there is a clear variation in the kinematic behaviour of the tS model in the mono-jet channel, which can be attributed to additional diagrams (accessible only in this channel) featuring a gluon in the initial state and subsequently allowing the mediator to go on-shell. In this scenario, when the resulting quark and DM particle are both small compared to the mediator mass, they share equally its energy leading to a peak in the E_T^{miss} distribution at approximately half the mediator mass.

In the cases where the kinematic distribution is independent of the width, we assume that the impact of the selection cuts in each channel is unchanged by the couplings. In this case, the following relations approximately hold:

$$\sigma \propto \begin{cases} g_q^2 g_\chi^2 / \Gamma & \text{if } M_{\text{med}} \geq 2m_{\text{DM}} \\ g_q^2 g_\chi^2 & \text{if } M_{\text{med}} < 2m_{\text{DM}} \end{cases} \quad (2.7)$$

in the sV and sA models [44], and

$$\sigma \propto g_{q\chi}^4 \quad (2.8)$$

in the tS model. When valid, these approximations allow us to greatly simplify our limit calculations, and for this reason, we restrict our primary results to regions of parameter space where $\Gamma/M_{\text{med}} < 0.5$ (see app. A for further details of the limit-setting calculation).

More problematically, it was noted by refs. [36, 44] that the Breit-Wigner propagator breaks down in the $m_{\text{DM}} \gg M_{\text{med}}$ region even if Γ/M_{med} is small. To correct for this we follow ref. [44], and rescale the cross section in the $m_{\text{DM}} > M_{\text{med}}$ region by a factor which takes into account the error introduced by using the Breit-Wigner propagator in the generator. The factor is found by convolving the PDF with both the kinetic and Breit-Wigner propagators in turn and taking the ratio at each mass point. We approximate the kinetic propagator by making the substitution $M_{\text{med}}\Gamma(M_{\text{med}}) \rightarrow s\Gamma(\sqrt{s})/M_{\text{med}}$ in the Breit-Wigner propagator.

A full study of the tS model within the mono-jet channel, where altering the coupling can lead to changed kinematic behaviour, has been performed elsewhere [52], and requires the production of individual samples for each coupling point. This, combined with the challenges associated with including differing orders of α_s , make the generation process computationally expensive compared to the mono- $Z(\text{lep})$ and mono- $W/Z(\text{had})$ channels. We therefore exclude an analysis of the tS model in the mono-jet channel in this work.

3 Recasting mono- X constraints

The procedure for recasting existing mono- X analyses to obtain SiM constraints follows a simple cut-and-count methodology. Firstly, signal events are simulated (described below in section 3.1) with object p_T smearing applied to approximate the detection efficiency of the ATLAS detector, ϵ . The event selection criteria of the mono- X analysis of interest

⁴In this discussion, the mono- $W/Z(\text{had})$ channel can be assumed to follow the same logic as for the mono- $Z(\text{lep})$ channel.



Figure 2: The E_T^{miss} distribution of the sV and tS models in the mono-jet and mono- $Z(\text{lep})$ channels, for some exemplary masses. The parameter μ is defined as Γ/M_{med} , and is used to demonstrate the impact of a changing width; the tS model in the mono-jet channel shows a clear width-dependence. The widths are obtained with couplings of 0.1, 1 and 5 where $\mu < 0.5$ remains true.

is then applied to the simulated signal samples. Events surviving the selection criteria are counted to determine the likelihood of a dark matter event being observed (referred to as the acceptance, \mathcal{A}), which is then used in combination with channel-specific model-independent limits on new physics events to limit the parameter space of a given model. For a comprehensive description of the recasting procedure, see appendix A.

In this paper, mono-jet constraints are derived from a search for new phenomena conducted by the ATLAS Collaboration using pp collisions at $\sqrt{s} = 8$ TeV as described in ref. [57]. Similarly, the leptonic mono- Z and hadronic mono- W/Z constraints are derived from ATLAS dark matter searches that were optimised for the D1, D5 and D9 effective operators [58, 59]. These analyses are described in further detail in sections 3.2, 3.3 and 3.4 respectively.

3.1 Signal simulation

Monte Carlo simulated event samples are used to model the expected signal for each channel and for each SiM. Leading order matrix elements for the process $pp \rightarrow X + \chi\bar{\chi}$ (where X is specifically one or two jets⁵, a $Z(\rightarrow \ell^+\ell^-)$ boson or a $W/Z(\rightarrow jj)$ boson) are first simulated using MADGRAPH5_aMC@NLO v2.2.2 [60] with the MSTW2008lo68cl PDF [61]. During this stage, the renormalisation and factorisation scales are set to the default sum of $\sqrt{m^2 + p_T^2}$ for all particles in the final state. Showering and hadronisation are then performed by PYTHIA 8.201 [62] with the appropriate PDF and using the ATLAS UE Tune AU2-MSTW2008LO [63]. Reconstruction of small-radius jets (hereon referred to just as ‘jets’) for the mono-jet channel is performed by FASTJET [64] using the anti- k_T algorithm with radius parameter $R = 0.4$. Similarly, reconstruction of large-radius jets for the mono- $W/Z(\text{had})$ channel is performed using the Cambridge-Aachen algorithm with $R = 1.2$. The latter channel also includes a mass-drop filtering procedure with $\mu = 0.67$ and \sqrt{y} ⁶ = 0.4 (see ref. [65] for further details), which favours large- R jets with two balanced subjects, consistent with the decay of an EW boson to a (potentially boosted) dijet pair. Lastly, the detector response is approximated by applying a Gaussian smearing factor to the p_T of all leptons and jets.

3.1.1 Parton matching scheme

In the ATLAS mono-jet analysis, matching of partons generated in MADGRAPH5 to jets generated in PYTHIA 8 is performed using the MLM scheme [66], with two matching scales, or values of ‘QCUT’, per mass/coupling point. In combination, the QCUT values span a broad kinematic range with a cut placed on the leading jet p_T per event to avoid double-counting. This treatment aims to mitigate the impact of the matching scale on the shape of the p_T and E_T^{miss} distributions; that is, to reduce the uncertainty in those areas of phase space where the transferred momentum is significantly larger or smaller than the QCUT value. For the analysis of SiMs, we use instead a single matching scale of 80 GeV. Though not ideal, this approach suitably reproduces the results of the ATLAS mono-jet analysis for the masses of interest (see Sec. B.1). Importantly, it also reduces the complexity and computational expense involved in estimating limits for the mono-jet channel.

We now move to a discussion of each of the mono- X channels separately.

3.2 Mono-jet constraints

The ATLAS mono-jet + E_T^{miss} analysis [57] was originally designed to set limits on three new physics scenarios, the most relevant of which is the production of WIMP DM within the context of a set of effective operators. The analysis also includes a brief study of a Z' DM model which is analogous to our sV model. **mention third signal scenario? or not mention it before?**

Signal selection is carried out based on at least one hard jet recoiling against missing energy. To ensure that the correct back-to-back jet + E_T^{miss} topology is selected events are

⁵Jets are seeded by any parton excluding the (anti-)top quark.

⁶ $\sqrt{y} = \min(p_{T_{j_1}}, p_{T_{j_2}})\Delta R/m_{jet}$ is the momentum balance of the two leading subjects.

required to have a leading jet, j_1 , with $p_T > 120$ GeV and $|\eta| < 2.0$ satisfying $p_T^{j_1}/E_T^{\text{miss}} > 0.5$. Surviving events must then fulfill $|\Delta\phi(j, \vec{E}_T^{\text{miss}})| > 1.0$, where j is any jet with $p_T > 30$ GeV and $|\eta| < 4.5$. This criterion reduces the multijet background contribution where the large E_T^{miss} originates mainly from jet energy mismeasurements. Note that there is no upper limit placed on the number of jets per event. The contribution from the dominant background processes, W/Z +jets, is managed with a veto on events containing muons or electrons with $p_T > 7$ GeV. Lastly, nine separate signal regions are defined with increasing lower thresholds on E_T^{miss} , which range from 150 GeV to 700 GeV as shown in table 2.

The ATLAS mono-jet analysis revealed no significant deviation of observed events from the expected SM backgrounds in the 8 TeV dataset of Run I. Subsequently, model-independent limits on new physics signatures were provided in terms of the visible cross-section, $\sigma \times \mathcal{A} \times \epsilon$; these are listed in table 2.

Signal Region	E_T^{miss} threshold [GeV]	$\sigma \times \mathcal{A} \times \epsilon$ [fb]
SR1	150	726 (935)
SR2	200	194 (271)
SR3	250	90 (106)
SR4	300	45 (51)
SR5	350	21 (29)
SR6	400	12 (17)
SR7	500	7.2 (7.2)
SR8	600	3.8 (3.2)
SR9	700	3.4 (1.8)

Table 2: The ATLAS mono-jet E_T^{miss} signal regions and corresponding observed (expected) model-independent upper limits on $\sigma \times \mathcal{A} \times \epsilon$ at 95% confidence level. Adapted from ref. [57].

The signal simulation procedure outlined in sec. 3.1 and implementation of the selection criteria discussed above were validated for the mono-jet channel via reproduction of ATLAS limits on the suppression scale, $M_\star \equiv M_{\text{med}}/\sqrt{g_q g_\chi}$, for the Z' model. The details of this process are contained in appendix B.1. Importantly, we observe agreement within $\sim 12\%$ for all samples.

3.3 Mono- $Z(\text{lep})$ constraints

The ATLAS mono- $Z(\text{lep}) + E_T^{\text{miss}}$ analysis [58] was principally designed to constrain a set of EFT models of DM. As a secondary focus, it also includes a short study of a t -channel SiM similar to our tS model.

The selection criteria for this analysis are summarised as follows (see the paper for a full description). Electrons (muons) are required to have a p_T greater than 20 GeV, and $|\eta|$ less than 2.47 (2.5). Two opposite-sign, same-flavour leptons are selected, and required to have invariant mass and pseudorapidity such that $m_{\ell\ell} \in [76, 106]$ GeV and $|\eta^{\ell\ell}| < 2.5$. The reconstructed Z boson should be approximately back-to-back and balanced against

the E_T^{miss} , ensured with the selections $\Delta\phi(\vec{E}_T^{\text{miss}}, p_T^{\ell\ell}) > 2.5$ and $|p_T^{\ell\ell} - E_T^{\text{miss}}|/p_T^{\ell\ell} < 0.5$. Events containing a jet with $p_T > 25$ GeV and $|\eta| < 2.5$ are vetoed. Events are also vetoed if they contain a third lepton with $p_T > 7$ GeV. The signal regions are defined by increasing lower E_T^{miss} thresholds: $E_T^{\text{miss}} > 150, 250, 350, 450$ GeV.

A cut-and-count strategy is used to estimate the total observed yields and expected SM backgrounds in each signal region. The limits on $\sigma \times \mathcal{A} \times \epsilon$ are not publicly available, so we take the numbers of expected and observed events, along with the associated uncertainties, and convert these into model-dependent upper limits with a single implementation of the HistFitter framework [67] using a frequentist calculator and a one-sided profile likelihood test statistic (the LHC default). The results of this process are displayed in table 3. Note that we use signal regions 1 and 2 only, as our simplified HistFitter approach is inadequate to handle the very low statistics of signal regions 3 and 4. These upper limits are also used for the validation of the mono- $Z(\text{lep})$ signal generation and selection procedures (see app. B.2).

Signal Region	E_T^{miss} threshold [GeV]	$\sigma \times \mathcal{A} \times \epsilon$ [fb]
SR1	150	1.59 (1.71)
SR2	250	0.291 (0.335)

Table 3: The ATLAS mono- $Z(\text{lep}) + E_T^{\text{miss}}$ signal regions and corresponding observed (expected) model-independent upper limits on $\sigma \times \mathcal{A} \times \epsilon$ at 95% confidence level. Adapted using HistFitter from ref. [58].

3.4 Mono- $W/Z(\text{had})$ constraints

The ATLAS mono- $W/Z(\text{had}) + E_T^{\text{miss}}$ search [59] was aimed at constraining the spin-independent effective operators C1, D1, and D5, and the spin-dependent operator D9. The search was originally designed to exploit the constructive interference of W boson emission from opposite-sign up-type and down-type quarks, leading to DM production wherein the mono- W channel is dominant. Recent studies [68] have revealed this scenario to violate gauge invariance and so we ignore it in this analysis.

The mono- $W/Z(\text{had})$ event selection is carried out as follows. Large-radius jets are selected using a mass-drop filtering procedure (see sec. 3.1) to suppress non- W/Z processes. Events are required to contain at least one large- R jet with $p_T > 250$ GeV, $|\eta| < 1.2$ and a mass, m_{jet} , within a 30-40 GeV window of the W/Z mass (i.e. $m_{\text{jet}} \in [50, 120]$ GeV). In order to reduce the $t\bar{t}$ and multijet backgrounds, a veto removes events containing a small- R jet with $\Delta\phi(\text{jet}, E_T^{\text{miss}}) < 0.4$, or containing more than one small- R jet with $p_T > 40$ GeV, $|\eta| < 4.5$, and $\Delta R(\text{small-}R \text{ jet}, \text{large-}R \text{ jet}) > 0.9$. Electrons, muons and photons are vetoed if their p_T is larger than 10 GeV and they lie within $|\eta| < 2.47$ (electrons), 2.5 (muons), 2.37 (photons). Two signal regions were defined with $E_T^{\text{miss}} > 350$ GeV and $E_T^{\text{miss}} > 500$ GeV.

329 The ATLAS analysis used a shape-fit of the mass distribution of the large- R jet to
 330 estimate the background yields in the two signal regions, along with the associated statis-
 331 tical and systematic uncertainties. As in the mono- $Z(\text{lep})$ case, we do not take the shapes
 332 into account but convert the published number of expected and observed events into upper
 333 limits on the expected and observed number of new physics events using the HistFitter
 334 framework. For the $E_T^{\text{miss}} > 500$ GeV signal region, we obtain the limits shown in table 4.
 335 We do not consider the first signal region with $E_T^{\text{miss}} > 350$ GeV in the recasting proce-
 336 dure, since the cut-and-count limits extracted could not be convincingly validated. The
 337 high E_T^{miss} signal region was found to be optimal for most operators studied by the ATLAS
 338 analysis.

Signal Region	E_T^{miss} threshold [GeV]	$\sigma \times \mathcal{A} \times \epsilon$ [fb]
SR2	500	1.35 (1.34)

Table 4: The ATLAS mono- $W/Z(\text{had})$ E_T^{miss} signal region considered in this work and corresponding observed (expected) model-independent upper limits on $\sigma \times \mathcal{A} \times \epsilon$ at 95% confidence level. Adapted using HistFitter from ref. [59].

339 4 Results and discussion

340 4.1 Limits on the couplings $\sqrt{g_q g_\chi}$

341 The 95% confidence level upper limits on the sV and sA model coupling combination
 342 $\sqrt{g_q g_\chi}$, and the tS model coupling $g_{q\chi}$, obtained from each of the mono- X channels, are
 343 presented in figs. 3-8. These quantities were evaluated as described in appendix A (in-
 344 cluding statistical and systematic uncertainties), and correspond to the best limits of each
 345 signal region tested.

346 In each plot, limits are shown ranging from <0.01 to the upper perturbative limit⁷ for
 347 each coupling, 4π ; where a limit was calculated to be larger than this, the limit is considered
 348 meaningless and the region is coloured grey. The white (hatched) regions coincide with
 349 those mass points which yield an initial (final) value of $\sqrt{g_q g_\chi}$ or $g_{q\chi}$ which fails to satisfy
 350 $\Gamma < M_{\text{med}}/2$. (We observe that values for which the width is just within our upper validity
 351 bound of $M_{\text{med}}/2$ may be pushed over into the invalid range with the addition of new
 352 particles, not considered here, which would serve to increase the mediator width.) When
 353 $g_\chi/g_q = 0.2$, only the mono-jet channel produces limits which survive this requirement,
 354 and so these are shown separately in fig 7.

355 Detailed comments specific to each channel are provided below, however some trends
 356 are channel-independent. For the sV model, strong limits exist when $M_{\text{med}} > 2m_\chi$ as the
 357 mediator can go on-shell, thereby enhancing the cross-section. The sA model limits are
 358 generally similar to the sV model limits except in the region corresponding to $m_{\text{DM}} \gtrsim$

⁷We note that perturbativity is questionable for couplings greater than $\sim 2 - 3$, however 4π is an oft-quoted hard upper limit.

359 $\sqrt{4\pi}M_{\text{med}}/g_{\chi}^{\text{gen}}$ where g_{χ}^{gen} is the DM coupling used at the generator level. We remove
 360 this region in the sA model to avoid violating perturbative unitarity, which can lead to
 361 an unphysical enhancement of the cross-section when the DM mass is much larger than
 362 the mediator mass [45, 46]. The upper limit on $\sqrt{g_q g_{\chi}}$ is relatively constant across values
 363 of g_{χ}/g_q , as is expected when the coupling (and hence the width) has been demonstrated
 364 to have little effect on kinematic behaviour (see sec 2.4), and using the assumptions of
 365 eq. 2.7. As the ratio increases, points in the region $M_{\text{med}} > m_{\chi}$ disappear as the initial
 366 value, $g_q = 1$, leads to a failure of the width condition. However, one could easily choose
 367 a smaller initial value of g_q to recover these points, and we suggest that the limits in this
 368 region would be quite similar to those seen in the $g_{\chi}/g_q = 0.2$ and 0.5 cases.

369 The constraints on the coupling strength are weaker when m_{χ} or M_{med} is large (>100
 370 GeV) owing to suppression of the cross-section. In this region, the constraints are expected
 371 to improve at higher centre-of-mass energies. For small DM masses with an off-shell me-
 372 diator, the $E_{\text{T}}^{\text{miss}}$ distribution is softer, therefore results in this region of phase space are
 373 limited by statistical uncertainties associated with the tail-end of the distribution. This
 374 region of phase space would benefit from further optimisation of event selection in analyses
 375 aimed at the study of simplified models, as we expect to see in the upcoming Run II results.

376 These mono- X searches are complementary to direct searches for the mediator via dijet
 377 resonances [69–72]. These have been used to study SiMs in, for example, [36, 46, 52]. Dijet
 378 studies search for the signature of a direct mediator decay into standard model particles,
 379 generally assuming a narrow resonance. These constraints can be stronger than mono- X
 380 constraints, particularly when the width is small and when the coupling to quarks is large
 381 relative to the coupling to DM. Mono- X searches however have the advantage for larger
 382 values of g_{χ}/g_q and smaller mediator masses.

383 We now examine channel-specific trends.

384 4.1.1 Mono-jet channel

385 The mono-jet channel upper limits on the coupling combination $\sqrt{g_q g_{\chi}}$ for the sV and
 386 sA models are displayed in the left-hand column of figs. 3-6, for $g_{\chi}/g_q = 0.5, 1, 2$ and
 387 5 respectively (where the ratio of 5 is only shown for the sV model, due to a lack of
 388 meaningful results in the sA model). The $g_{\chi}/g_q = 0.2$ case is shown separately in fig. 7, as
 389 these limits are only meaningful within this channel.

390 As expected, the mono-jet channel produces the strongest coupling limits for both
 391 s -channel models, which are better than those from the next-best mono- $Z(\text{lep})$ channel
 392 by a factor of a few. For these models, the weakest limits result for large m_{χ} or large
 393 M_{med} , and in fact are so weak that they are pushed into the region of invalidity where
 394 $\Gamma > M_{\text{med}}/2$. Although the acceptance is considerably higher when both m_{DM} and M_{med}
 395 are large compared to low masses, the cross-section is sufficiently small so as to nullify any
 396 gain. Within the valid region ($m_{\chi} \in [1, 100]$ GeV and $M_{\text{med}} \in [1, 200]$ GeV), the limit
 397 on $\sqrt{g_q g_{\chi}}$ generally ranges from 0.1 to 0.7, with a handful of on-shell masses reaching a
 398 limit of ~ 0.05 in the large g_{χ}/g_q case. In the large g_{χ}/g_q scenario, limits for $m_{\chi} = 1000$
 399 GeV start to become valid; where $\sqrt{g_q g_{\chi}}$ remains constant but g_{χ}/g_q increases, the value

of g_q is pushed downward and so the width, which is dominated by decays to SM particles, decreases with respect to m_{DM} .

The uncertainties on the limits for both s -channel models are dominated by contributions from the matching scale at acceptance-level, and generally range from $\sim 5\%$ to 46% .

4.1.2 Mono- $Z(\text{lep})$ channel

The simplicity of the mono- $Z(\text{lep})$ channel relative to the mono-jet channel, and the ease of signal simulation at MADGRAPH5 level allowed us to study a finer granularity of points in the mass phase space. The resulting limits on the sV and sA models are shown in the central column of figs. 3-6. While the behaviour of the limits as g_χ/g_q is varied is similar to that within the mono-jet channel, the mono- $Z(\text{lep})$ limits are overall weaker by a factor of a few.

The total relative uncertainties on $\sqrt{g_q g_\chi}$ for the s -channel models are generally within 10% , but can range up to 80% in a few cases where m_χ is small; they are in general split equally between statistical and systematic contributions.

The advantage of the mono-boson channels is in the study of the tS model; since this was not included in the mono-jet channel the strongest limits are obtained with the mono- $Z(\text{lep})$ analysis, and are shown in the left-hand side of fig. 8. Note that, in comparison to the s -channel models, the limits have weakened by a factor of 10. This is the result of an orders-of-magnitude weaker cross-section and the inability of the mediator to go on-shell in this channel. We find stronger limits for smaller m_χ and M_{med} masses, where larger cross-sections compensate for lower acceptances at these points. Overall, the uncertainties contribute less than 10% .

4.1.3 Mono- $W/Z(\text{had})$ channel

The limits on the couplings of the sV , sA and tS models, obtained within the mono- $W/Z(\text{had})$ channel, are shown in the right-hand column of figs. 3-8. This channel was included for comparison with the leptonic mono- $Z(\text{lep})$ channel in particular, but a coarser selection of masses was chosen as the limits were initially found to be somewhat weaker. Additionally, further estimates were made: a) as the kinematic behaviour is reasonably independent of the couplings, a single acceptance was found for each (m_χ, M_{med}) combination and applied to each value of g_χ/g_q , and b) complete systematic uncertainties were generated for a subset of masses and compared to those from the mono- $Z(\text{lep})$ channel; from this comparison the mono- $Z(\text{lep})$ systematic uncertainties were multiplied by 2 and then applied to the mono- $W/Z(\text{had})$ limits. As a result, the limits obtained in this channel are not intended to be rigorously quantitative; rather, they are used to indicate qualitatively how the channel compares.

The ATLAS mono- $W/Z(\text{had})$ analysis (and in particular the higher $E_{\text{T}}^{\text{miss}}$ signal region) was not optimised for a SiM interpretation, and much of the phase space produced insignificant numbers of events passing the event selection, with up to 200 thousand events generated. Generally, the limits are a factor of a few weaker than those from the mono- $Z(\text{lep})$ channel, which is both consistent with the limits on the EFT models studied in the

ATLAS analyses, and expected following our use of a cut-and-count interpretation, rather than a shape analysis, of the mono- W/Z (had) public results.

In some cases the limits become comparable with the mono- Z (lep) channel, suggesting that more statistics and an improved treatment of systematic uncertainties would bring these closer in line with that channel.

Overall, the uncertainties from this channel lie within 5 to 50%, most of the time being between 10 and 30%. Generally, both statistical and systematic uncertainties contribute in a similar manner. A few points are clearly limited by the generated statistics, resulting in a statistical error of up to 90%. Points with high m_χ and low M_{med} tend to have larger systematic uncertainties.

4.2 Comparison with relic density constraints

In figs. 3-8 we show lines where the constraint on the coupling corresponds to the coupling strength that would reproduce the correct DM relic density if DM is a thermal relic of the early universe. For points diagonally above and to the left of the solid purple line, the LHC constraints naively rule out the couplings leading to the correct relic density. Below and to the right of this line the relic density coupling is still allowed. In some cases the intercept does not pass through a significant number of data points surviving the quality criteria outlined in previous sections. In these cases the line is not shown.

In this scenario, the measured abundance is approximately related to the unknown self-annihilation cross-section via:


$$\Omega_{\text{DM}} h^2 \simeq \frac{2 \times 2.4 \times 10^{-10} \text{ GeV}^{-2}}{\langle \sigma v \rangle_{\text{ann}}}. \quad (4.1)$$

This is used with measurements of the DM abundance by Planck, $\Omega_{\text{DM}}^{\text{obs}} h^2 = 0.1199 \pm 0.0027$ [73], to find $\langle \sigma v \rangle_{\text{ann}} \simeq 4.0 \times 10^{-9} \text{ GeV}^{-2}$ for thermal relic DM. This relation is only approximately accurate, and so we use the micrOMEGAs code [74] to determine the coupling strength leading to the correct relic density for each model. This technique was verified against the semi-analytic technique outlined in e.g. ref. [75].


If the DM mass lies at the electroweak scale, the thermal relic scenario provides a natural explanation for the observed DM density. The coupling strengths leading to the correct relic density are therefore a natural benchmark with which to compare constraints from collider (and indeed direct detection) searches. However the relic density couplings should by no means be regarded as serious constraints. If DM is not produced thermally or there is an unknown effect which modifies the evolution of the density with temperature, then eq. 4.1 breaks down. Additionally, in the scenario where we assume DM to be a thermal relic, we ignore the possibility of there being other annihilation channels and other beyond-SM particles contributing to the DM abundance, which, if taken into account, would also invalidate eq. 4.1.

4.3 Comparison with direct detection constraints


In figs 3-8 we also show the intercept line where constraints from direct detection experiments are equivalent to our mono- X constraints. Below and to the right of the dashed




(a) sV model, $g_\chi/g_q = 0.5$, mono-jet channel.




(b) sV model, $g_\chi/g_q = 0.5$, mono- Z channel.




(c) sV model, $g_\chi/g_q = 0.5$, mono- W/Z channel.



(d) sA model, $g_\chi/g_q = 0.5$, mono-jet channel.




(e) sA model, $g_\chi/g_q = 0.5$, mono- Z channel.




(f) sA model, $g_\chi/g_q = 0.5$, mono- W/Z channel.


Figure 3: Upper limits on the coupling for the s -channel models in the mono-jet (left), mono- Z (lep) (centre) and mono- W/Z (had) (right) channels, for $g_\chi/g_q = 0.5$. The grey region represents the phase space where no meaningful limit was obtained. The hatched region represents a limit which leads to a width greater than $M_{\text{med}}/2$, so the validity of the calculation begins to fail. The dotted region represents phase space where insufficient statistics were available. The purple dotted line shows the threshold where direct detection constraints become stronger than the mono- X constraint. Similarly the solid purple line shows where relic density constraints become stronger than mono- X constraints. See the text for further details.




(a) sV model, $g_\chi/g_q = 1$, mono-jet channel.




(b) sV model, $g_\chi/g_q = 1$, mono-Z channel.




(c) sV model, $g_\chi/g_q = 1$, mono- W/Z channel.



(d) sA model, $g_\chi/g_q = 1$, mono-jet channel.




(e) sA model, $g_\chi/g_q = 1$, mono-Z channel.




(f) sA model, $g_\chi/g_q = 1$, mono- W/Z channel.


Figure 4: Upper limits on the couplings for the s -channel models in the mono-jet (left), mono- Z (lep) (centre) and mono- W/Z (had) (right) channels, for $g_\chi/g_q = 1$. Refer to fig. 3 for details.




(a) sV model, $g_\chi/g_q = 2$, mono-jet channel.




(b) sV model, $g_\chi/g_q = 2$, mono- Z channel.




(c) sV model, $g_\chi/g_q = 2$, mono- W/Z channel.



(d) sA model, $g_\chi/g_q = 2$, mono-jet channel.




(e) sA model, $g_\chi/g_q = 2$, mono- Z channel.



(f) sA model, $g_\chi/g_q = 2$, mono- W/Z channel.

Figure 5: Upper limits on the coupling for the s -channel models in the mono-jet (left), mono- Z (lep) (centre) and mono- W/Z (had) (right) channels, for $g_\chi/g_q = 2$. Refer to fig. 3 for details.



(a) sV model, $g_\chi/g_q = 5$, mono-jet channel.

(b) sV model, $g_\chi/g_q = 5$, mono- Z channel.

(c) sV model, $g_\chi/g_q = 5$, mono- W/Z channel.

Figure 6: Upper limits on the coupling for the sV model in the mono-jet (left), mono- Z (lep) (centre) and mono- W/Z (had) (right) channels, for $g_\chi/g_q = 5$. Refer to fig. 3 for details.




Figure 7: Upper limits on the coupling for the s -channel models in the mono-jet channel, for $g_\chi/g_q = 0.2$. Refer to fig. 3 for details.




Figure 8: Upper limits on the coupling $g_{q\chi}$ for the t -channel model in the mono- Z (lep) (left) and mono- W/Z (had) (right) channels. Refer to fig. 3 for details.

purple line, direct detection constraints are stronger while above and to the left of this line, the LHC gives the stronger limit. As with the relic density contours, we do not show the intercept where it does not pass through sufficient valid data points. We use the toolset from ref. [76] to convert the strongest available direct detection constraints, which are from the LUX 2013 dataset [77], onto constraints on our models.

Compared to direct detection, the mono- X collider limits perform relatively better for the sA model than for the sV model. This is because the axial-vector coupling leads to a suppressed scattering rate in direct detection experiments while collider searches are relatively insensitive to the difference between the vector and axial-vector couplings. In the non-relativistic limit, the tS model leads to a mix of both suppressed and unsuppressed operators.

The direct detection constraints assume that the DM candidate under consideration contributes 100% of the local DM density, while the mono- X constraints make no assumptions about either the local DM density or overall abundance. In this sense the mono- X limits remain useful even in those regions of phase space where they are not as strong as those from direct detection.

5 Conclusion

In this paper we have examined a set of three simplified dark matter models, extracting constraints from ATLAS Run I mono- X plus missing energy searches featuring the associated production of a mono-jet, $Z(\rightarrow \text{leptons})$, or $W/Z(\rightarrow \text{hadrons})$. We explored a phase space where both the DM and mediator masses span $\mathcal{O}(\text{GeV})$ to $\mathcal{O}(\text{TeV})$, and considered ratios of g_χ/g_q of 0.2, 0.5, 1, 2 and 5 in the s -channel models. Rather than setting limits in the $M_{\text{med}} - m_{\text{DM}}$ plane for a fixed value of the coupling strength, we instead constrained the coupling strength as a function of both M_{med} and m_{DM} in a 3D plane. Whilst this approach necessitates the introduction of some approximations, it also allows for a thorough examination of the interplay between the DM production cross-section and the free parameters of the models.

As expected, the mono-jet channel is found to yield the strongest limits on vector and axial-vector SM and DM couplings to a vector mediator exchanged in the s -channel. This channel is also found to perform well for small values of g_χ . The limits obtained in the mono- $Z(\text{lep})$ channel, in comparison, are generally weaker by a factor of a few, while the mono- $W/Z(\text{had})$ results are weaker again. This is partly due to our conservative estimations of the systematic uncertainties and partly due to limited statistics resulting from a harder $E_{\text{T}}^{\text{miss}}$ selection cut. The width effects associated with the t -channel exchange of an $\text{SU}(2)$ doublet scalar mediator are observed to vanish in both the mono- $Z(\text{lep})$ and mono- $W/Z(\text{had})$ channels, greatly simplifying the analysis and confirming these as straightforward and competitive channels for future collider DM detection.

Where the axial-vector model is not excluded by perturbative unitarity requirements, we find the coupling limits to be on par with those of the vector model within each analysis channel. Weaker limits are found for the t -channel model, a result of cross-section suppression not present in the s -channel models.

Finally, we compared our limits to constraints from relic density and direct detection; although each search has a different set of assumptions, this demonstrates the complementarity and impressive reach of simplified models as a tool for the interpretation of collider DM searches. We eagerly await the improved constraints expected from Run II of the LHC.

6 Acknowledgements

A.J.B. and M.F.M. were supported by the Australian Research Council. J.G. was supported by SNF (grant 200020_156083). We thank Karl Nordström for discussions on the cross-section reweighting, Brian Peterson and Steven Schramm for helpful discussions, and Sean Crosby for technical support.

A Limit setting strategy

In this appendix we present a summary of the procedure employed to calculate the 95% confidence level (CL) limits on the coupling parameter $\sqrt{g_q g_\chi}$, where this parameter can be replaced with $g_{q\chi}$ for the tS model, and M_\star in the validation of the mono-jet analysis.

A.1 Nominal values

For each SiM, the nominal limit is calculated by taking the model-independent upper limit on $\sigma \times \mathcal{A} \times \epsilon$ from each analysis, dividing by the value of $\mathcal{A} \times \epsilon$ (which is taken as a single parameter for each point) to obtain the limiting cross section σ_{lim} , and rearranging eq. 2.7 to convert to a limit on the couplings. In the s -channel on-shell case, the width can be expressed as a function of g_q and the ratio g_χ/g_q , which simplifies the calculation. We arrive at

$$\sqrt{g_q g_\chi}_{\text{lim}} = \begin{cases} \sqrt{g_q g_\chi}_{\text{gen}} \times (\sigma_{\text{lim}}/\sigma_{\text{gen}})^{\frac{1}{2}} & \text{if } M_{\text{med}} \geq 2m_{\text{DM}} \text{ (} s\text{-channel)} \\ \sqrt{g_q g_\chi}_{\text{gen}} \times (\sigma_{\text{lim}}/\sigma_{\text{gen}})^{\frac{1}{4}} & \text{if } M_{\text{med}} < 2m_{\text{DM}} \end{cases} \quad (\text{A.1})$$

where $\sqrt{g_q g_\chi}_{\text{gen}}$ and σ_{gen} are the input couplings and cross-section (taken from PYTHIA 8), respectively.

The signal region in each case is chosen based on where the best ‘expected’ limit lies, where that limit is calculated assuming that exactly the expected SM background is observed.

A.2 Uncertainty estimation

Our nominal limits rely on both σ_{gen} and $\mathcal{A} \times \epsilon$ and so are subject to systematic uncertainties which derive from our choice of signal generation procedure. For our signal samples, there are three key sources of systematic uncertainty: the factorisation and renormalisation scales, the strong coupling constant (α_s) and the choice of parton distribution function (PDF).

We assess the impact of the factorisation and renormalisation default scales in a straightforward manner; by varying them simultaneously by factors of 2 (‘up’) and 0.5

main systematic sources	PDF/tune	factorisation and renormalisation scales	matching scale (mono-jet only)
variation ‘up’	NNPDF2.1LO + Monash tune	2	160 GeV
nominal	MSTW2008lo68cl + ATLAS UE AU2-MSTW2008LO	1	80 GeV
variation ‘down’	CTEQ6L1 + ATLAS UE AU2-CTEQ6L1	0.5	40 GeV

Table 5: Reading left to right, the sources of systematic uncertainty considered in this analysis. Each point in phase space is varied up or down by one of these sources, and the systematic uncertainty is then taken from the resultant changes to the acceptance and cross-section in comparison to their nominal values.

(‘down’). The systematic effects of the strong coupling constant and PDF are difficult to separate and so are treated in tandem. We assume that the systematic uncertainty introduced by α_s at matrix-element level is negligible when compared to the PDF uncertainties, as demonstrated to be valid in ref. [78]. The variation of α_s in conjunction with a change of PDF is done with the use of specific tunes in PYTHIA 8, which we change simultaneously with the PDF choice to estimate the uncertainty on $\Delta\sigma_{gen}$. The nominal choices of PDF and tune are varied ‘up’ to NNPDF2.1LO PDF + Monash tune, and ‘down’ to CTEQ6L1 PDF and ATLAS UE AU2-CTEQ6L1 tune. For the mono-jet channel, the impact of the matching scale (QCUT) is assessed in a manner similar to that of the factorisation and renormalisation scales. That is, we vary the QCUT by factors of 2 (‘up’ to 160 GeV) and 0.5 (‘down’ to 40 GeV). These systematic uncertainty sources are summarised in table 5.

The average variation in the nominal value of σ_{lim} (measured as a fraction of σ_{lim}) resulting from each systematic source is added in quadrature and propagated to $\sqrt{g_q g_\chi}$ to obtain the total systematic uncertainty. This process is adjusted slightly to account for the inclusion of statistical uncertainties, which are estimated conservatively by taking the 95% CL *lower* limit on $\mathcal{A} \times \epsilon$ as calculated with the Wald approximation, i.e. $\mathcal{A} \times \epsilon \rightarrow (\mathcal{A} \times \epsilon) - \Delta(\mathcal{A} \times \epsilon)$. Note that the uncertainty on the luminosity is less than 3%, so is considered to be negligible in comparison to other systematic sources.

B Validation of signal simulation and event selection procedures

B.1 Mono-jet channel

The signal generation and selection procedures for the mono-jet channel are validated via reproduction of the ATLAS limits on $M_\star \equiv M_{\text{med}}/\sqrt{g_q g_\chi}$, for the s -channel vector SiM. A comparison of SR7⁸ limits for a representative sample of mediator masses with $m_\chi = 50$ GeV, $\Gamma = M/8\pi$ and $\sqrt{g_q g_\chi} = 1$ is presented in Table 6. In general, good agreement is observed between the ATLAS and reproduced limits, with a maximum difference of 12%. We note that a discrepancy of a few percent is expected given the differences in signal simulation. For example, the simplified matching procedure discussed in detail in Sec 3.1.1 introduces an additional uncertainty of approximately 25% for events with $E_T^{\text{miss}} > 350$ GeV when compared to the approach utilised by the ATLAS mono-jet group. Further uncertainties are introduced by the jet smearing approximation used in place of a full detector simulation and by the 95% CL estimation procedure (outlined in Appendix A) used instead of a thorough HistFitter treatment. As our results are consistently more conservative than those of the ATLAS analysis, we consider our approach to be acceptable.

M_\star^{gen} [TeV]	$M_\star^{95\% \text{CL}}$ [GeV] (ATLAS)	$M_\star^{95\% \text{CL}}$ [GeV] (this work)	Difference [%]
0.05	91	89	2.16
0.3	1151	1041	7.3
0.6	1868	1535	11.8
1	2225	1732	12.0
3	1349	1072	6.8
6	945	769	8.5
10	928	724	10.6
30	914	722	9.6

Table 6: Comparison of the 95% CL upper limits on M_\star from this work and from the ATLAS mono-jet analysis [57]. The limits are for an s -channel vector mediator model with $m_\chi = 50$ GeV and $\Gamma = M_{\text{med}}/8\pi$, and for the process $pp \rightarrow \chi\bar{\chi} + 1, 2j$ with QCUT = 80 GeV. Note that M_\star^{gen} is the input suppression scale.

B.2 Mono- $Z(\text{lep})$ channel

The ATLAS mono- $Z(\text{lep})$ results include an upper limit on the coupling $g_{q\chi}$ for a t -channel SiM analogous to our tS model, and so it is this model which we use to validate our signal generation and selection procedures. Note that the following differences exist: the ATLAS model includes just two mediators (up - and $down$ -type) where we consider six, the DM particle is taken to be Majorana where we assume Dirac, and the couplings $g_{t,b\chi}$ are set to zero where we have universal coupling to all three quark generations.

⁸We use this signal region as it is the only one for which ATLAS limits are provided.

m_χ [GeV]	M_{med} [GeV]	$g_{q\chi}^{95\% \text{CL}}$ (ATLAS)	$g_{q\chi}^{95\% \text{CL}}$ (this work)	Difference [%]
10	200	1.9	2.0	5.3
	500	2.8	3.2	14.3
	700	3.5	4.4	25.7
	1000	4.5	5.2	15.6
200	500	3.4	4.0	17.6
	700	4.2	4.5	7.1
	1000	5.2	5.3	1.9
400	500	5.5	5.7	3.6
	700	6.1	6.5	6.6
	1000	7.2	7.4	2.8
1000	1200	23.3	24.1	3.4

Table 7: Comparison of the 95% CL upper limit on $g_{q\chi}$ from this work and from the ATLAS mono- $Z(\text{lep})$ analysis [58]. The limits are for a variant of the t -channel scalar mediator model with Majorana Dark matter for the process $pp \rightarrow \chi\bar{\chi} + Z(\rightarrow e^+e^-/\mu^+\mu^-)$.

Table 7 shows the 95% CL upper limits on $g_{q\chi}$ that we calculate using our own generation procedure (and the values in table 3), compared with the limits taken from the ATLAS analysis. Also shown is the difference as a percentage of the ATLAS limit. We see reasonable agreement; most of the 11 points in parameter space are within 10% of the ATLAS limits, and all are within 26%. Additionally, our results are consistently more conservative, which again is to be expected given the less sophisticated nature of our generation procedure. As in the case of the mono-jet validation, the differences stem from the use of p_T smearing applied to the leptons (rather than a full reconstruction simulation) and from the simplified treatment of systematics; we also obtained $\sigma \times \mathcal{A} \times \epsilon$ independently.

B.3 Mono- $W/Z(\text{had})$ channel

The event generation and selection procedures for the mono- $W/Z(\text{had})$ channel are validated via reproduction of the ATLAS limits on M_\star for the D5 and D9 effective operators with $m_\chi = 1$ GeV. We see agreement within 12.5% and 7.4% respectively, where the ATLAS limits are consistently stronger, as shown in table B.3. The relative sizes of the discrepancies are expected given that only low- E_T^{miss} limits are available for the D5 operator while we use the high- E_T^{miss} signal region in our recast. Note that a general discrepancy of a few percent is expected for both operators for the reasons discussed in sections B.1 and B.2, and also because we use a cut-and-count approach while the ATLAS limits are extracted using a shape-fit. Furthermore, the ATLAS limits are quoted at 90% CL while ours are calculated at 95% CL.

EFT operator	m_χ [GeV]	$M_\star^{90\%CL}$ [GeV] (ATLAS)	$M_\star^{95\%CL}$ [GeV] (this work)	Difference [%]
D9	1	2400	2221	7.4
D5	1	570	499	12.5

Table 8: Comparison of the 95% CL upper limits on M_\star from this work and from the ATLAS mono- W/Z (had) analysis [59]. The limits are for the process $pp \rightarrow \chi\bar{\chi} + W/Z (\rightarrow jj)$.

References

- [1] ATLAS Collaboration, *Search for new phenomena with the monojet and missing transverse momentum signature using the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions*, *Phys. Lett. B* (2011), arXiv:1106.5327.
- [2] ATLAS Collaboration. *Search for New Phenomena in Monojet plus Missing Transverse Momentum Final States using 10 fb¹ of pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector at the LHC*, 2012, ATLAS-CONF-2012-147.
- [3] CMS Collaboration. *Search for new physics in monojet events in pp collisions at $\sqrt{s} = 8$ TeV*, 2013, CMS-PAS-EXO-12-048.
- [4] ATLAS Collaboration, *Search for new phenomena with mono-jet plus missing transverse energy signature in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector*, 2012, ATL-COM-PHYS-2012-1211.
- [5] ATLAS Collaboration. *Further search for supersymmetry at $\sqrt{s} = 7$ TeV in final states with jets, missing transverse momentum and isolated leptons with the ATLAS detector*, *Phys.Rev.* (2012), arXiv:1208.4688.
- [6] ATLAS Collaboration, *Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb⁻¹ of $\sqrt{s} = 7$ TeV proton-proton collision data*, *Phys.Rev. D* 87, 012008 (2013), arXiv:1208.0949.
- [7] ATLAS Collaboration, *Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton-proton collision data*, *JHEP* 09 (2014) 146, arXiv:1405.7875.
- [8] ATLAS Collaboration, *Search for pair-produced third-generation squarks decaying via charm quarks or in compressed supersymmetric scenarios in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys.Rev. D* 90, 052008 (2014), arXiv:1407.0608.
- [9] J. Goodman et al., *Constraints on Dark Matter from Colliders*, *Phys.Rev. D* 82, 116010 (2010), arXiv:1008.1783.
- [10] Y. Bai, P. J. Fox, and R. Harnik, *The Tevatron at the Frontier of Dark Matter Direct Detection*, *JHEP* (2010), arXiv:1005.3797.
- [11] P. J. Fox, R. Harnik, J. Kopp, and Y. Tsai, *LEP Shines Light on Dark Matter*, *Phys.Rev.* (2011), arXiv:1103.0240.
- [12] M. L. Graesser, I. M. Shoemaker, and L. Vecchi, *A Dark Force for Baryons*, 2011, arXiv:1107.2666.

- [13] H. An and F. Gao, *Fitting CoGeNT Modulation with an Inelastic, Isopin-Violating Z' Model*, 2011, arXiv:1108.3943.
- [14] H. Dreiner et al., *Contact Interactions Probe Effective Dark Matter Models at the LHC*, *Europhys.Lett.* (2013), arXiv:1303.3348.
- [15] J. Goodman et al., *Gamma Ray Line Constraints on Effective Theories of Dark Matter*, *Nucl.Phys.* (2011), arXiv:1009.0008.
- [16] O. Buchmueller, M. J. Dolan and C. McCabe, *Beyond Effective Field Theory for Dark Matter Searches at the LHC*, *JHEP* 1401 (2014) 025 arXiv:1308.6799.
- [17] G. Busoni et al., *On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC*, *Phys.Lett.* (2014), arXiv:1307.2253.
- [18] G. Busoni et al., *On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC, Part II: Complete Analysis for the s-channel*, *JCAP* 1406:060 (2014), arXiv:1402.1275.
- [19] G. Busoni et al., *On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC Part III: Analysis for the t-channel*, *JCAP* 09 (2014) 022, arXiv:1405.3101.
- [20] J. Abdallah et al. *Simplified Models for Dark Matter and Missing Energy Searches at the LHC*, (2014), arXiv:1409.2893.
- [21] J. Abdallah et al. *Simplified Models for Dark Matter Searches at the LHC*, *Phys. Dark Uni.* 9-10 (2015) 8-23, arXiv:1506.03116.
- [22] D. Abercrombie et al. *Dark Matter Benchmark Models for Early LHC Run-2 Searches: Report of the ATLAS/CMS Dark Matter Forum*, arXiv:1507.00966.
- [23] P. Harris, V. V. Khoze, M. Spannowsky and C. Williams, *Constraining Dark Sectors at Colliders: Beyond the Effective Theory Approach*, *Phys.Rev.* (2015), arXiv:1411.0535.
- [24] Oliver Buchmueller, Matthew J. Dolan, Sarah A. Malik and Christopher McCabe, *Characterising dark matter searches at colliders and direct detection experiments: Vector mediators*, 2014, arXiv:1407.8257.
- [25] A. DiFranzo, K. I. Nagao, A. Rajaraman, and T. M. P. Tait *Simplified Models for Dark Matter Interacting with Quarks*, *JHEP* 11 (2013) 014, arXiv:1308.2679.
- [26] M. R. Buckley, D. Feld, and D. Goncalves, *Scalar Simplified Models for Dark Matter*, *Phys. Rev. D* 91 (2015) 015017, arXiv:1410.6497.
- [27] M. R. Buckley, *Using Effective Operators to Understand CoGeNT and CDMS-Si*, *Phys.Rev. D* 88, 055028 (2013), arXiv:1308.4146.
- [28] N. Zhou, D. Berge, and D. Whiteson, *Mono-everything: combined limits on dark matter production at colliders from multiple final states*, *Phys.Rev. D* 87, 095013 (2013), arXiv:1302.3619.
- [29] R. M. Godbole, G. Mendiratta and T. M. P. Tait, *A Simplified Model for Dark Matter Interacting Primarily with Gluons*, *JHEP* 08 (2015) 064, arXiv:1506.01408.
- [30] J. Heisig, M. Krmer, M. Pellen and C. Wiebusch, *Constraints on Majorana Dark Matter from the LHC and IceCube*, arXiv:1509.07867.
- [31] M. Blennow, J. Herrero-Garcia, T. Schwetz and S. Vogl, *Halo-independent tests of dark matter direct detection signals: local DM density, LHC, and thermal freeze-out*, *JCAP* 1508 (2015) 08, 039 arXiv:1505.05710.

- [32] O. Lebedev and Y. Mambrini, *Axial dark matter: The case for an invisible Z'* , *Phys. Lett. B* 734 (2014) 350 arXiv:1403.4837.
- [33] A. Alves, A. Berlin, S. Profumo and F. S. Queiroz, *Dark Matter Complementarity and the Z' Portal*, *Phys. Rev. D* 92 (2015) 8, 083004 arXiv:1501.03490.
- [34] A. Alves, S. Profumo and F. S. Queiroz, *The dark Z' portal: direct, indirect and collider searches*, *JHEP* 1404 (2014) 063 arXiv:1312.5281.
- [35] A. Alves, A. Berlin, S. Profumo and F. S. Queiroz, *Dirac-fermionic dark matter in $U(1)_X$ models*, *JHEP* 1510 (2015) 076 arXiv:1506.06767.
- [36] H. An, X. Ji and L. T. Wang, *Light Dark Matter and Z' Dark Force at Colliders*, *JHEP* 1207 (2012) 182, arXiv:1202.2894.
- [37] H. An, R. Huo and L. T. Wang, *Searching for Low Mass Dark Portal at the LHC*, *Phys. Dark Univ.* 2 (2013) 50 arXiv:1212.2221.
- [38] M. T. Frandsen, F. Kahlhoefer, A. Preston, S. Sarkar and K. Schmidt-Hoberg, *LHC and Tevatron Bounds on the Dark Matter Direct Detection Cross-Section for Vector Mediators*, *JHEP* 1207 (2012) 123 arXiv:1204.3839.
- [39] G. Arcadi, Y. Mambrini, M. H. G. Tytgat and B. Zaldivar, *Invisible Z' and dark matter: LHC vs LUX constraints*, *JHEP* 1403 (2014) 134 arXiv:1401.0221.
- [40] I. M. Shoemaker and L. Vecchi, *Unitarity and Monojet Bounds on Models for DAMA, CoGeNT, and CRESST-II*, *Phys. Rev. D* 86 (2012) 015023 arXiv:1112.5457.
- [41] M. T. Frandsen, F. Kahlhoefer, S. Sarkar and K. Schmidt-Hoberg, *Direct detection of dark matter in models with a light Z'* , *JHEP* 1109 (2011) 128 arXiv:1107.2118.
- [42] P. Gondolo, P. Ko and Y. Omura, *Light dark matter in leptophobic Z' models*, *Phys. Rev. D* 85 (2012) 035022 arXiv:1106.0885.
- [43] M. Fairbairn and J. Heal, *Complementarity of dark matter searches at resonance*, *Phys. Rev. D* 90 (2014) 11, 115019 arXiv:1406.3288.
- [44] T. Jacques and K. Nordstrom, *Mapping monojet constraints onto Simplified Dark Matter Models*, *JHEP* 06 (2015) 142, arXiv:1502.05721.
- [45] N. F. Bell, Y. Cai and R. K. Leane, *Mono- W Dark Matter Signals at the LHC: Simplified Model Analysis*, *JCAP* 1601 (2016) 01, 051, arXiv:1512.00476.
- [46] M. Chala, F. Kahlhoefer, M. McCullough, G. Nardini and K. Schmidt-Hoberg, *Constraining Dark Sectors with Monojets and Dijets*, *JHEP* 1507 (2015) 089 arXiv:1503.05916.
- [47] F. Kahlhoefer, K. Schmidt-Hoberg, T. Schwetz and S. Vogl, *Implications of unitarity and gauge invariance for simplified dark matter models*, *JHEP* 1602 (2016) 016, arXiv:1510.02110.
- [48] G. Jungman et al., *Supersymmetric dark matter*, *Phys.Rept.* (1996).
- [49] Y. Bai and J. Berger, *Fermion Portal Dark Matter*, *JHEP* 1311 (2013) 171 arXiv:1308.0612.
- [50] H. An, L. T. Wang and H. Zhang, *Dark matter with t -channel mediator: a simple step beyond contact interaction*, *Phys. Rev. D* 89 (2014) 11, 115014 arXiv:1308.0592.
- [51] S. Chang, R. Edezhath, J. Hutchinson and M. Luty, *Effective WIMPs*, *Phys. Rev. D* 89 (2014) 1, 015011 arXiv:1307.8120.
- [52] M. Papucci, A. Vichi, and K. M. Zurek, *Monojet versus rest of the world I: t -channel Models*, *JHEP* (2014), arXiv:1402.2285.

- [53] M. Garny, A. Ibarra and S. Vogl, *Signatures of Majorana dark matter with t -channel mediators*, *Int. J. Mod. Phys. D* 24 (2015) 07, 1530019 arXiv:1503.01500.
- [54] M. Garny, A. Ibarra, S. Rydbeck and S. Vogl, *Majorana Dark Matter with a Coloured Mediator: Collider vs Direct and Indirect Searches*, *JHEP* 1406 (2014) 169 arXiv:1403.4634.
- [55] N. F. Bell, J. B. Dent, A. J. Galea, T. D. Jacques, L. M. Krauss and T. J. Weiler, *W/Z Bremsstrahlung as the Dominant Annihilation Channel for Dark Matter, Revisited*, *Phys. Lett. B* 706 (2011) 6 arXiv:1104.3823.
- [56] ATLAS Collaboration, *Search for dark matter produced in association with a hadronically decaying vector boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector at the LHC*, ATLAS-CONF-2015-080 (2015).
- [57] ATLAS Collaboration. *Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, 2015, arXiv:1502.01518
- [58] ATLAS Collaboration. *Search for dark matter in events with a Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys.Rev.D* **90** (2014) 012004, arXiv:1404.0051.
- [59] ATLAS Collaboration. *Search for dark matter in events with a hadronically decaying W or Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. Lett.* **112** (2014) 041802, arXiv:1309.4017.
- [60] J. Alwall /emphet al.. *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP07* (2014) 079, arXiv:1405.0301.
- [61] A. D. Martin, W. J. Stirling, R. S. Thorne, G. Watt, *Parton distributions for the LHC*, *Eur.Phys.J.C* **63**, (2009), 189-285, arXiv:0901.0002.
- [62] T. Sjöstrand et al. *An Introduction to PYTHIA 8.2*, *Comput. Phys. Comm.* 191 (2015) 159-177, arXiv:1410.3012.
- [63] ATLAS Collaboration. *Summary of ATLAS Pythia8 tunes*, 2012, ATL-PHYS-PUB-2012-003.
- [64] M. Cacciari, G. P. Salam and G. Soyez, *FastJet user manual*, arXiv:1111.6097.
- [65] ATLAS Collaboration, *Performance of jet substructure techniques for large- R jets in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector*, *JHEP* 09 (2013) 076, arXiv:1306.4945.
- [66] M. L. Mangano et al. *Matching matrix elements and shower evolution for top-quark production in hadronic collisions*, *JHEP* 01 (2007), hep-ph/0611129.
- [67] M. Baak et al. *HistFitter software framework for statistical data analysis*, *Eur.Phys.J.C* (2015), arXiv:1410.1280.
- [68] N. F. Bell et al, *Dark matter at the LHC: EFTs and gauge invariance*, *Phys. Rev. D* 92 (2015) 053008, arXiv:1503.07874.
- [69] CMS Collaboration, *Search for narrow resonances using the dijet mass spectrum in pp collisions at $\sqrt{s} = 8$ TeV*, *Phys.Rev.* (2013), arXiv:1302.4794.
- [70] G. Aad et al. [ATLAS Collaboration], *Search for new phenomena in the dijet mass distribution using $p-p$ collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. D* 91 (2015) 5, 052007, arXiv:1407.1376.

- [71] T. Aaltonen *et al.* [CDF Collaboration], *Search for new particles decaying into dijets in proton-antiproton collisions at $s^{1/2} = 1.96$ -TeV*, *Phys. Rev. D* **79** (2009) 112002, arXiv:0812.4036
- [72] V. Khachatryan *et al.* [CMS Collaboration], *Search for resonances and quantum black holes using dijet mass spectra in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Phys. Rev. D* **91** (2015) 052009, arXiv:1501.04198
- [73] P. A. R. Ade *et al.* [Planck Collaboration], *Planck 2013 results. XVI. Cosmological parameters*, *Astron. Astrophys.* **571**, A16 (2014) [arXiv:1303.5076 [astro-ph.CO]].
- [74] G. Blanger, F. Boudjema, A. Pukhov and A. Semenov, *micrOMEGAs4.1: two dark matter candidates*, *Comput. Phys. Commun.* **192**, 322 (2015) [arXiv:1407.6129 [hep-ph]].
- [75] G. Busoni, A. De Simone, T. Jacques, E. Morgante and A. Riotto, *Making the Most of the Relic Density for Dark Matter Searches at the LHC 14 TeV Run*, *JCAP* **03** (2015) 022, arXiv:1410.7409.
- [76] M. Cirelli, E. Del Nobile and P. Panci, *Tools for model-independent bounds in direct dark matter searches*, *JCAP* **1310** (2013) 019, arXiv:1307.5955.
- [77] D. S. Akerib *et al.* [LUX Collaboration], *First results from the LUX dark matter experiment at the Sanford Underground Research Facility*, *Phys. Rev. Lett.* **112** (2014) 091303, arXiv:1310.8214.
- [78] S. Schramm, *Searching for Dark Matter with the ATLAS Detector in Events with an Energetic Jet and Large Missing Transverse Momentum*, 2015, CERN-THESIS-2015-038.
- [79] N. Bell *et al.*, *Searching for Dark Matter at the LHC with a Mono-Z*, *Phys.Rev. D* **86**, 096011 (2012), arXiv:1209.0231.
- [80] M. Cahill-Rowley *et al.*, *Complementarity and Searches for Dark Matter in the p MSSM*, SLAC-PUB-15450 (2013), arXiv:1305.6921.
- [81] J. Kumar and D. Marfatia, *Matrix element analyses of dark matter scattering and annihilation*, *Phys.Rev.* (2013), arXiv:1305.1611.
- [82] P. J. Fox *et al.*, *Missing Energy Signatures of Dark Matter at the LHC*, *Phys.Rev.* (2012), arXiv:1109.4398.
- [83] P. J. Fox, R. Harnik, R. Primulando, and C-T. Yu, *Taking a Razor to Dark Matter Parameter Space at the LHC*, *Phys.Rev.* (2012), arXiv:1203.1662.
- [84] ATLAS Collaboration, *Search for high-mass resonances decaying to dilepton final states in pp collisions at $s^{1/2} = 7$ -TeV with the ATLAS detector*, *JHEP* (2012), arXiv:1209.2535.
- [85] J. Kumar and D. Marfatia, *Matrix element analyses of dark matter scattering and annihilation*, *Phys.Rev.* (2013), arXiv:1305.1611.
- [86] D. Alves *et al.*, *Simplified Models for LHC New Physics Searches*, *J.Phys.* (2012), arXiv:1105.2838.
- [87] CMS Collaboration, *Search for new physics in monojet events in pp collisions at $\sqrt{s} = 8$ TeV*, 2013, CMS-PAS-EXO-12-048.
- [88] ATLAS Collaboration, *Search for New Phenomena in Monojet plus Missing Transverse Momentum Final States using 10 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector at the LHC*, 2012, ATLAS-CONF-2012-147.
- [89] A. Cooper-Sarkar, *PDFs for the LHC*, 2011, arXiv:1107.5170.

- 812 [90] ATLAS Collaboration. *Search for dark matter candidates and large extra dimensions in*
813 *events with a jet and missing transverse momentum with the ATLAS detector*, 2013,
814 CERN-PH-EP-2012-210, arXiv:1210.4491.
- 815 [91] P. J. Fox et al. *Missing Energy Signatures of Dark Matter at the LHC*, *Phys. Rev.*, 2012.
- 816 [92] N. Bell, J. Dent, T. Jacques, and T. Weiler. *W/Z Bremsstrahlung as the Dominant*
817 *Annihilation Channel for Dark Matter*, *Phys. Rev.*, 2011.