

# Probability

G V V Sharma\*

## CONTENTS

1	Axioms	1
2	Elementary Probability	4
3	Random Variables	6
4	Transformation of Variables	18
5	Independent Random Variables	19
6	Binomial Distribution	28
7	Poisson Distribution	30
8	Gaussian Distribution	32
9	Geometric Distribution	36
10	Two Dimensions	36
11	Integral Transforms	42
12	Markov Chain	45
13	Inequalities	53
14	Convergence	55
15	Statistics	64

**Abstract**—This book provides solved examples on Probability

## 1 AXIOMS

1.1. The probability that a given positive integer lying between 1 and 100 (both inclusive) is NOT divisible by 2,3 or 5 is ...

**Solution:** Table 1.1.1 summarizes the given information.

Event	Definition	Probability
A	$n \equiv 0 \pmod{2}$	$\frac{50}{100}$
B	$n \equiv 0 \pmod{3}$	$\frac{33}{100}$
C	$n \equiv 0 \pmod{5}$	$\frac{20}{100}$
AB	$n \equiv 0 \pmod{6}$	$\frac{16}{100}$
BC	$n \equiv 0 \pmod{15}$	$\frac{6}{100}$
AC	$n \equiv 0 \pmod{10}$	$\frac{10}{100}$
ABC	$n \equiv 0 \pmod{30}$	$\frac{3}{100}$

TABLE 1.1.1:  $1 \leq n \leq 100$

\*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

$$\begin{aligned}
 \therefore \Pr(A + B + C) &= \Pr(A) + \Pr(B) + \Pr(C) \\
 &\quad - \Pr(AB) - \Pr(BC) \\
 &\quad - \Pr(AC) + \Pr(ABC) \quad (1.1.1)
 \end{aligned}$$

Substituting from Table 1.1.1 in (1.1.1),

$$\begin{aligned}\Pr(A + B + C) &= \frac{50}{100} + \frac{33}{100} + \frac{20}{100} \\ &\quad - \frac{16}{100} - \frac{6}{100} - \frac{10}{100} + \frac{3}{100} \\ &= \frac{74}{100} \quad (1.1.2)\end{aligned}$$

Thus, the required probability is

$$1 - \Pr(A + B + C) = \frac{26}{100} \quad (1.1.3)$$

- 1.2. P and Q are considering to apply for a job. The probability that P applies for the job is  $\frac{1}{4}$ , the probability that P applies for the job given that Q applies for the job is  $\frac{1}{2}$ , and the probability that Q applies for the job given that P applies for the job is  $\frac{1}{3}$ . Then the probability that P does not apply for the job given that Q does not apply for the job is

- a)  $\frac{4}{5}$       b)  $\frac{5}{6}$       c)  $\frac{7}{8}$       d)  $\frac{11}{12}$

**Solution:** The given information can be expressed as

$$\Pr(P) = \frac{1}{4} \quad (1.2.1)$$

$$\Pr(P|Q) = \frac{1}{2} = \frac{\Pr(PQ)}{\Pr(Q)} \quad (1.2.2)$$

$$\Pr(Q|P) = \frac{1}{3} = \frac{\Pr(PQ)}{\Pr(P)} \quad (1.2.3)$$

which yields

$$\begin{aligned}\Pr(PQ) &= \frac{1}{3} \times \frac{1}{4} = \frac{1}{12} \\ \Pr(Q) &= \frac{\frac{1}{12}}{\frac{1}{2}} = \frac{1}{6}\end{aligned} \quad (1.2.4)$$

The objective is to find

$$\Pr(P'|Q') \quad (1.2.5)$$

(1.2.1) can be expressed as

$$\Pr(P'|Q') = \frac{\Pr(P'Q')}{\Pr(Q')} \quad (1.2.6)$$

$$= \frac{\Pr(1 - (P + Q)')}{\Pr(Q')} \quad (1.2.7)$$

$$= \frac{1 - \Pr(P) - \Pr(Q) + \Pr(PQ)}{1 - \Pr(Q)} \quad (1.2.8)$$

Substituting from (1.2.4) and (1.2.1) in (1.2.8),

$$\Pr(P'|Q') = \frac{4}{5} \quad (1.2.9)$$

- 1.3. Out of 6 unbiased coins, 5 are tossed independently and they all result in heads. If the 6th coin is now independently tossed, the probability of getting head is:

- (a) 1  
(b) 0  
(c)  $\frac{1}{2}$   
(d)  $\frac{1}{6}$

**Solution:** Define a random variable  $X = \{0, 1\}$  denoting the outcome of the toss of 6th coin with  $X = 0$  and  $X = 1$  representing tails and head respectively. Therefore,

$$\Pr(X = 0) + \Pr(X = 1) = 1 \quad (1.3.1)$$

$$\Pr(X = 1) = \frac{1}{2} \quad (1.3.2)$$

Hence the correct answer is option (c).

- 1.4. Two students are solving the same problem independently, if the probability of first one solves the problem is  $\frac{3}{5}$  and the probability that the second one solves the problem is  $\frac{4}{5}$ , what is the probability that atleast one of them solves the problem?

- a)  $\frac{17}{25}$   
b)  $\frac{19}{25}$   
c)  $\frac{21}{25}$   
d)  $\frac{23}{25}$

**Solution:** Let X, Y be two events representing solving the problem by students A, B respectively.

Given

$$\Pr(X) = \frac{3}{5} \quad (1.4.1)$$

$$\Pr(Y) = \frac{4}{5} \quad (1.4.2)$$

Since students solve the problem independently, So events X and Y are independent, For independent events

$$\Pr(XY) = \Pr(X) \times \Pr(Y) \quad (1.4.3)$$

from (1.4.1) and (1.4.2)

$$\Pr(XY) = \frac{3}{5} \times \frac{4}{5} \quad (1.4.4)$$

$$\Pr(XY) = \frac{12}{25} \quad (1.4.5)$$

Now we have to find probability of solving the problem by atleast one of them i.e  $\Pr(X + Y)$ . As,

$$\Pr(X + Y) = \Pr(X) + \Pr(Y) - \Pr(XY) \quad (1.4.6)$$

from (1.4.1), (1.4.2), (1.4.5)

$$\Pr(X + Y) = \frac{3}{5} + \frac{4}{5} - \frac{12}{25} \quad (1.4.7)$$

$$\Pr(X + Y) = \frac{23}{25} \quad (1.4.8)$$

Hence the required probability is  $\frac{23}{25}$

1.5. Three types of components are used in electrical circuits 1, 2, 3 as shown below in the figure

**Solution:** For  $q_1$ , the truth table Multiplying

A	B	C	$(AB) + C$
1	1	0	1
1	1	1	1
0	1	1	1
0	0	1	1
1	0	1	1

TABLE 1.5.1: Circuit 1 working

and adding probability for each case of  $q_1$  gives us the value of  $q_1$  as

$$q_1 = p^3 - 2p^2 + 1 \quad (1.5.1)$$

For  $q_2$ , the truth table Multiplying and adding probability for each case of  $q_2$  gives us the




Fig. 1.5.1: Figure

A	B	C	$(A + B)C$
1	1	1	1
1	0	1	1
0	1	1	1

TABLE 1.5.2: Circuit 2 working

value of  $q_2$  as

$$q_2 = p^3 - p^2 - p + 1 \quad (1.5.2)$$

For  $q_3$ , the truth table Multiplying and adding


A	B	C	$A + B + C$
1	0	0	1
0	1	0	1
0	0	1	1
1	1	0	1
1	0	1	1
0	1	1	1
1	1	1	1

TABLE 1.5.3: Circuit 3 working

probability for each case of  $q_3$  gives us the

value of  $q_3$  as

$$q_3 = 1 - p^3 \quad (1.5.3)$$



$$\therefore q_3 > q_1 > q_2 \quad (1.5.4)$$

Hence **Option 1** is correct

## 2 ELEMENTARY PROBABILITY

2.1. There are 3 red socks, 4 green socks and 3 blue socks. You choose 2 socks. The probability that they are of the same colour is

- a)  $\frac{1}{5}$       b)  $\frac{7}{30}$       c)  $\frac{1}{4}$       d)  $\frac{4}{15}$

**Solution:** Let  $X_i \in \{1, 2, 3\}$  represent the  $i^{th}$  draw, where 1, 2, 3 correspond to the colour of socks drawn as Red, Blue and Green respectively

TABLE 2.1.1

	$X_1 = 1$	$X_1 = 2$	$X_1 = 3$
$X_2 = 1$	6/90	12/90	9/90
$X_2 = 2$	12/90	12/90	12/90
$X_2 = 3$	9/90	12/90	6/90

TABLE 2.1.1 represents all the possibilities of choosing socks one by one.

The probability that the two socks drawn are of the same colour (by substituting values from

table 2.1.1)

$$= \Pr(X_1 = X_2) \quad (2.1.1)$$

$$= \sum_{i=1}^3 \Pr(X_2 = i | X_1 = i) \Pr(X_1 = i) \quad (2.1.2)$$

$$= \frac{6}{90} + \frac{12}{90} + \frac{6}{90} \quad (2.1.3)$$

$$= \frac{4}{15} \quad (2.1.4)$$

So the correct option is (D)

2.2. A box contains 40 numbered red balls and 60 numbered black balls. From the box, balls are drawn one by one at random without replacement till all the balls are drawn. The probability that the last ball drawn is black equals ... Now, this problem is equivalent to the problem where we have to arrange 40 distinct R's and 60 distinct B's such that, a B should come at last. So, the desired probability is given by

$$\frac{(\text{placing a B at last}) \times (\text{arranging other letters})}{\text{arranging 100 letters}} = \frac{60 \times 99!}{100!} = \frac{3}{5} \quad (2.2.1)$$

2.3. An experiment consists of two papers. paper1 and paper2. The probability of failing in paper 1 is .3 and that in paper 2 is .2. Given that a student has failed in paper 2, the probability of failing in paper 1 is .6. The probability of student failing in both is

a) .5

b) .18

c) .12

d) .06

**Solution:** Table 2.3.1 summarises the given information. The desired probability is

$$\Pr(X = 0, Y = 0) = \Pr(X = 0 | Y = 0) \Pr(Y = 0) \quad (2.3.1)$$

$$= .12 \quad (2.3.2)$$

2.4. An urn contains 5 red balls and 5 black balls. In

	Description	Probability
0	failure	$\Pr(X = 0) = 0.3$
1	success	$\Pr(Y = 0) = 0.2$
X	Paper 1	$\Pr(X = 0 Y = 0) = 0.6$
Y	Paper 2	

TABLE 2.3.1: Description

the first draw, one ball is picked at random and discarded without noticing its colour. The probability to get a red ball in the second draw is

- a)  $\frac{1}{2}$       b)  $\frac{4}{9}$       c)  $\frac{5}{9}$       d)  $\frac{6}{9}$

**Solution:** Let  $X_i \in \{0, 1\}$  represent the  $i^{\text{th}}$  draw where 1 denotes a red ball being drawn.

	$X_1 = 0$	$X_1 = 1$
$X_2 = 0$	4/18	5/18
$X_2 = 1$	5/18	4/18

TABLE 2.4.1: The probabilities of all possible cases when two balls are drawn one by one from the urn.

From Table 2.4.1,

$$\Pr(X_2 = 1) = \Pr(X_2 = 1, X_1 = 0) + \Pr(X_2 = 1, X_1 = 1) \quad (2.4.1)$$

$$= \frac{5}{18} + \frac{4}{18} \quad (2.4.2)$$

$$= \frac{1}{2} \quad (2.4.3)$$

The required option is (A).

- 2.5. A sample of size  $n = 2$  is drawn from a population of size  $N = 4$  using probability proportional to size without replacement scheme, Where the probabilities proportional to size are The probability of inclusion of unit (1) in

i:	1	2	3	4
$P_i$	0.4	0.2	0.2	0.2

Table : Probability vs Size

the sample is

- a) 0.4      b) 0.6      c) 0.7      d) 0.75

**Solution:** Let  $P_i(j)$  represent the probability for selecting unit (j) as second unit after selecting unit (i)

$$P_i(j) = \frac{p_j}{1 - p_i} \quad (2.5.1)$$

Let  $\Pr(i, j)$  be probability of selecting sample  $\{i, j\}$ , using (2.5.1) is

$$\Pr(i, j) = P_i(j) + P_j(i) \quad (2.5.2)$$

$$= \left( p_i \times \frac{p_j}{1 - p_i} \right) + \left( p_j \times \frac{p_i}{1 - p_j} \right) \quad (2.5.3)$$

Total samples (Size  $n = 2$ ) are Let  $P_i$  be

Case	1	2	3	4	5	6
Sample (size $n = 2$ )	(1,2)	(1,3)	(1,4)	(2,3)	(2,4)	(3,4)

TABLE 2.5.1: list of samples

the probability of inclusion of unit (i) in the sample (size  $n = 2$ ), Now i will calculate  $P_1$ , Favourable cases for inclusion of unit (1) are case (1,2,3), So

$$P_1 = \Pr(1, 2) + \Pr(1, 3) + \Pr(1, 4) \quad (2.5.4)$$

using (2.5.3) and  $p_i$  from question ,

$$P_1 = \frac{7}{30} + \frac{7}{30} + \frac{7}{30} \quad (2.5.5)$$

$$= 0.7 \quad (2.5.6)$$

Therefore Option (3) is correct.

- 2.6. There are two boxes. Box-1 contains 2 red balls and 4 green balls. Box-2 contains 4 red balls and 2 green balls. A box is selected at random and a ball is chosen randomly from the selected box. If the ball turns out to be red, what is the probability that Box-1 had been selected?

**Solution:** Box-1 has 2 red balls and 4 green balls.

Box-2 has 4 red balls and 2 green balls.

Let  $B \in \{1, 2\}$  represent a random variable where 1 represents selecting box-1 and 2 represents selecting box-2. From Baye's theorem

$$\begin{aligned} \Pr(R = 1) &= \Pr(R = 1|B = 1) \times \Pr(B = 1) \\ &+ \Pr(R = 1|B = 2) \times \Pr(B = 2) \end{aligned} \quad (2.6.1)$$

Event	definition	value
$\Pr(B = 1)$	Probability of selecting Box-1	$\frac{1}{2}$
$\Pr(B = 2)$	Probability of selecting Box-2	$\frac{1}{2}$
$\Pr(R = 1 B = 1)$	Probability of drawing red ball from Box-1	$\frac{1}{3}$
$\Pr(G = 1 B = 1)$	Probability of drawing green ball from Box-1	$\frac{2}{3}$
$\Pr(R = 1 B = 2)$	Probability of drawing red ball from Box-2	$\frac{2}{3}$
$\Pr(G = 1 B = 2)$	Probability of drawing green ball from Box-2	$\frac{1}{3}$

TABLE 2.6.1: Table 1

Substituting values from table (2.6.1) in (2.6.1)

$$\Pr(R = 1) = \frac{1}{2} \quad (2.6.2)$$

$$\Pr((R = 1)(B = 1)) = \Pr(R = 1|B = 1) \times \Pr(B = 1) \quad (2.6.3)$$

$$= \frac{1}{6} \quad (2.6.4)$$

We need to find  $\Pr(B = 1|R = 1)$

$$\Pr(B = 1|R = 1) = \frac{\Pr((R = 1)(B = 1))}{\Pr(R = 1)} \quad (2.6.5)$$

$$= \frac{1}{3} \quad (2.6.6)$$

$\therefore$  The desired probability that box-1 is selected  $= \frac{1}{3}$

2.7. An urn has 3 red and 6 black balls. Balls are drawn at random one by one without replacement. The probability that second red ball appears on fifth draw is:

- a)  $\frac{1}{9!}$
- b)  $\frac{4!}{9!}$
- c)  $4 \left( \frac{6!4!}{9!} \right)$
- d)  $\frac{6!4!}{9!}$

**Solution:** To obtain a second red ball at the fifth draw, the first 4 trials should involve drawing only 1 red ball out of the 3 and 3 black

balls out of the 6. Probability of this happening:

$$\frac{{}^3C_1 {}^6C_3}{{}^9C_4} \quad (2.7.1)$$

The probability of the fifth ball turning out to be red is:

$$\frac{{}^2C_1}{{}^5C_1} \quad (2.7.2)$$

By Multiplication rule, total probability:

$$\frac{{}^3C_1 {}^6C_3 {}^2C_1}{{}^5C_1 {}^9C_4} = \frac{3! \times 6! \times 2! \times 4! \times 4! \times 5!}{2! \times 3! \times 3! \times 5! \times 9!} \quad (2.7.3)$$

$$= 4 \left( \frac{4!6!}{9!} \right) \quad (2.7.4)$$

### 3 RANDOM VARIABLES

3.1. Consider the function  $f(x)$  defined as  $f(x) = ce^{-x^4}$ ,  $x \in R$ . For what value of  $c$  is  $f$  a probability density function?

- a)  $\frac{2}{\Gamma(1/4)}$
- b)  $\frac{4}{\Gamma(1/4)}$
- c)  $\frac{3}{\Gamma(1/3)}$
- d)  $\frac{1}{4\Gamma(4)}$

**Solution:**

3.2. Let  $X$  and  $Y$  be i.i.d random variables uniformly distributed on  $(0,4)$ . Then  $\Pr(X > Y|X < 2Y)$  is

- a)  $1/3$
- b)  $5/6$
- c)  $1/4$
- d)  $2/3$

**Solution:**

The PDF is given by

$$f_X(x) = f_Y(x) = \begin{cases} \frac{1}{4}, & \text{if } 0 < x < 4 \\ 0, & \text{otherwise} \end{cases}$$

The CDF is given by

$$F(x) = \int_{-\infty}^x f(x)dx$$

$$F_X(x) = F_Y(x) = \begin{cases} 0, & x \leq 0 \\ \frac{x}{4}, & \text{if } 0 < x < 4 \\ 1, & x \geq 4 \end{cases}$$

Using definition of conditional probability

$$\Pr(X > Y | X < 2Y) = \frac{\Pr(Y < X < 2Y)}{\Pr(X < 2Y)} \quad (3.2.1)$$

Now finding  $\Pr(X < 2Y)$

$$\Pr(X < 2y) = F_X(2y) \quad (3.2.2)$$

$$\Rightarrow \Pr(X < 2Y) = \int_{-\infty}^{\infty} f_Y(x) \times F_X(2x)dx \quad (3.2.3)$$

$$\Rightarrow \Pr(X < 2Y) = \int_0^2 \frac{x}{8}dx + \int_2^4 \frac{1}{4}dx \quad (3.2.4)$$

$$\Rightarrow \Pr(X < 2Y) = \frac{3}{4} = 0.75 \quad (3.2.5)$$

Now to find  $\Pr(Y < X < 2Y)$

$$\Pr(y < X < 2y) = F_X(2y) - F_X(y) \quad (3.2.6)$$

$$\Rightarrow \Pr(Y < X < 2Y) \quad (3.2.7)$$

$$= \int_{-\infty}^{\infty} f_Y(x)(F_X(2x) - F_X(x))dx$$

$$\Rightarrow \int_0^2 \frac{1}{4} \left( \frac{x}{2} - \frac{x}{4} \right) dx + \int_2^4 \frac{1}{4} \left( 1 - \frac{x}{4} \right) dx \quad (3.2.8)$$

$$\Rightarrow \Pr(Y < X < 2Y) = \frac{1}{4} = 0.25 \quad (3.2.9)$$

Now using (3.2.1), (3.2.5) and (3.2.9)

$$\Pr(X > Y | X < 2Y) = \frac{1/4}{3/4} = \frac{1}{3} \quad (3.2.10)$$

Hence final solution is option 1) or 1/3

3.3. Suppose  $X$  is a positive random variable with the following probability density function,

$$f(x) = (\alpha x^{\alpha-1} + \beta x^{\beta-1})e^{-x^\alpha - x^\beta}; x > 0$$

for  $\alpha > 0, \beta > 0$ . Then the hazard function of  $X$  for some choices of  $\alpha$  and  $\beta$  can be

a) an increasing function.

b) a decreasing function.

c) a constant function.

d) a non monotonic function

**Solution:**

CDF of  $X$ ,

$$F(x) = \int_{-\infty}^x f(t)dt \quad (3.3.1)$$

$$= \int_0^x f(t)dt \quad \text{as } x > 0 \quad (3.3.2)$$

$$= \int_{-\infty}^x ((\alpha t^{\alpha-1} + \beta t^{\beta-1}) \times e^{-t^\alpha - t^\beta}) dt \quad (3.3.3)$$

$$= -e^{-t^\alpha - t^\beta} \Big|_0^x \quad (3.3.4)$$

$$= 1 - e^{-x^\alpha - x^\beta} \quad (3.3.5)$$

Hazard function,


$$h(x) = \frac{f(x)}{1 - F(x)} \quad (3.3.6)$$

$$= \alpha x^{\alpha-1} + \beta x^{\beta-1} \quad (3.3.7)$$

$$h'(x) = \alpha(\alpha - 1)x^{\alpha-2} + \beta(\beta - 1)x^{\beta-2} \quad (3.3.8)$$

$$h'(x) = \begin{cases} 0 & \alpha = \beta = 1 \\ > 0 & \text{otherwise} \end{cases} \quad (3.3.9)$$

Thus  $h(x)$  can be either constant function or an increasing function.



From the above figure, it is verified that  $h(x)$  can be either constant function or an increasing function.

Correct options are 1,3.

3.4. Suppose the random variable  $X$  has the follow-

ing probability density function

$$f(x) = \begin{cases} \alpha (x - \mu)^{\alpha-1} e^{-(x-\mu)^\alpha}; & x > \mu \\ 0 & x \leq \mu \end{cases}$$

where  $\alpha > 0, -\infty < \mu < \infty$ . Which of the following are correct? The hazard function of  $X$  is

- a) an increasing function for all  $\alpha > 0$
- b) a decreasing function for all  $\alpha > 0$
- c) an increasing function for some  $\alpha > 0$
- d) a decreasing function for some  $\alpha > 0$

**Solution:**

For the random variable  $X$ , the CDF is

$$F(x) = \int_0^x f(y) dy \quad (3.4.1)$$

$$= \int_0^\mu 0 dy + \int_\mu^x \alpha (y - \mu)^{\alpha-1} e^{-(y-\mu)^\alpha} dy \quad (3.4.2)$$

$$= 0 - e^{-(y-\mu)^\alpha} \Big|_\mu^x \quad (3.4.3)$$

$$= 1 - e^{-(x-\mu)^\alpha} \quad (3.4.4)$$

For  $X$ , the hazard function  $H(y)$  is defined as

$$\begin{aligned} H(y) &= \frac{f(y)}{1 - F(y)} \\ \Rightarrow H(y) &= \begin{cases} \frac{\alpha (y-\mu)^{\alpha-1} e^{-(y-\mu)^\alpha}}{1 - (1 - e^{-(y-\mu)^\alpha})}; & y > \mu \\ 0 & y \leq \mu \end{cases} \\ &= \begin{cases} \alpha (y - \mu)^{\alpha-1}; & y > \mu \\ 0 & y \leq \mu \end{cases} \end{aligned}$$

Differentiating  $H(y)$  w.r.t.  $y$

$$H'(y) = \begin{cases} \alpha (\alpha - 1) (y - \mu)^{\alpha-2}; & y > \mu \\ 0 & y \leq \mu \end{cases}$$

When  $y \leq \mu$  then  $H'(y)$  is 0. When  $y > \mu$  then  $(y - \mu)^{\alpha-2}$  is positive. This implies that the sign for  $H'(y)$  for  $y > \mu$  is decided by the sign of  $\alpha (\alpha - 1)$ .

$$\alpha (1 - \alpha) < 0 \Rightarrow 0 < \alpha < 1$$

$$\alpha (1 - \alpha) > 0 \Rightarrow \alpha > 1 \quad (\text{ignoring } \alpha < 0)$$

$\therefore$  The Hazard function of  $X$  is decreasing when  $\alpha \in (0, 1)$  and increasing when

$\alpha \in (1, \infty)$  **Solution:** Options 3, 4




Fig. 3.4.1: Decreasing Hazard Function




Fig. 3.4.2: Increasing Hazard Function

3.5. Let  $X$  be a random variable with a certain non-degenerate distribution. Then identify the correct statements

- a) If  $X$  has an exponential distribution then  $median(X) < E(X)$
- b) If  $X$  has a uniform distribution on an interval  $[a, b]$ , then  $E(X) < median(X)$
- c) If  $X$  has a Binomial distribution then  $V(X) < E(X)$
- d) If  $X$  has a normal distribution, then  $E(X) < V(X)$

**Solution:** Expected value( $E(X)$ ): It is nothing but weighted average Median( $median(X)$ ): It is the value separating the higher half from the lower half of a data sample

Variance( $V(X)$ ): It is the expectation of the squared deviation of a random variable from its mean

- a) Let's consider  $X$  has an exponential distribution



bution.

$$X \sim \text{Exp}(\lambda) \quad (3.5.1)$$

where  $\lambda$  is rate parameter.

Probability function of exponential distribution,

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & x < 0 \end{cases} \quad (3.5.2)$$

The expected value of  $X \sim \text{Exp}(\lambda)$ ,

$$E(X) = \frac{1}{\lambda} \quad (3.5.3)$$

The median of  $X \sim \text{Exp}(\lambda)$ ,

$$\text{median}(X) = \frac{\ln 2}{\lambda} \quad (3.5.4)$$

$$\ln 2 < 1 \quad (3.5.5)$$

$$\frac{\ln 2}{\lambda} < \frac{1}{\lambda} \quad (3.5.6)$$

$$\text{median}(X) < E(X) \quad (3.5.7)$$

Hence, option 1 is correct.

- b) Let's consider  $X$  has a uniform distribution in interval  $[a, b]$ ,

$$X \sim U(a, b) \quad (3.5.8)$$

where,  $a$  = lower limit

$b$  = upper limit

Probability function of uniform distribution,

$$f_X(k) = \begin{cases} \frac{1}{b-a} & a \leq x \leq b \\ 0 & x < a, x > b \end{cases} \quad (3.5.9)$$

The expected value of  $X \sim U(a, b)$ ,

$$E(X) = \frac{1}{2}(a + b) \quad (3.5.10)$$

The median of  $X \sim U(a, b)$ ,

$$\text{median}(X) = \frac{1}{2}(a + b) \quad (3.5.11)$$

$$E(X) = \text{median}(X) \quad (3.5.12)$$

Hence, option 2 is incorrect.

- c) Let's consider  $X$  has a binomial distribution,

$$X \sim B(n, p) \quad (3.5.13)$$

where,  $n$  = no. of trials

$p$  = success parameter

Probability function of binomial distribution,

$$f_X(k) = \begin{cases} {}^nC_k p^k (1-p)^{n-k} & 0 \leq k \leq n \\ 0 & \text{otherwise} \end{cases} \quad (3.5.14)$$

The expected value of  $X \sim B(n, p)$ ,

$$E(X) = np \quad (3.5.15)$$

The variance of  $X \sim B(n, p)$ ,

$$V(X) = \sigma^2 = np(1-p) \quad (3.5.16)$$

$$1-p \leq 1 \quad (3.5.17)$$

$$np(1-p) \leq np \quad (3.5.18)$$

$$V(X) \leq E(X) \quad (3.5.19)$$

Hence, option 3 is incorrect.

- d) Let's consider  $X$  has a normal distribution,

$$X \sim N(\mu, \sigma^2) \quad (3.5.20)$$

where,  $\mu$  = mean of distribution

$\sigma^2$  = variance

Probability function of normal distribution,

$$f_X(k) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\left(\frac{x-\mu}{\sigma}\right)^2} \quad (3.5.21)$$

The expected value of  $X \sim N(\mu, \sigma^2)$ ,

$$E(X) = \mu \quad (3.5.22)$$

The variance of  $X \sim N(\mu, \sigma^2)$ ,

$$V(X) = \sigma^2 \quad (3.5.23)$$

$E(X)$  and  $V(X)$  are user defined. So, they can take any value.

Hence, option 4 is incorrect.

- 3.6. A fair coin is tossed repeatedly. Let  $X$  be the number of tails before the first heads occurs. Let  $Y$  denote the number of tails between the first and second heads. Let  $X + Y = N$ . Then which of the following are true?

- a)  $X$  and  $Y$  are independent random variables with

$$\Pr(X = k) = \Pr(Y = k) = \begin{cases} 2^{-(k+1)} & k = 0, 1, 2, \dots \\ 0 & \text{otherwise} \end{cases} \quad (3.6.1)$$

- b)  $N$  has a probability mass function given by

$$\Pr(N = k) = \begin{cases} (k-1)2^{-k} & k = 2, 3, 4, \dots \\ 0 & \text{otherwise} \end{cases} \quad (3.6.2)$$

- c) Given  $N = n$ , the conditional distribution of  $X$  and  $Y$  are independent  
d) Given  $N = n$

$$\Pr(X = k) = \begin{cases} \frac{1}{n+1} & n = 0, 1, 2, \dots \\ 0 & \text{otherwise} \end{cases} \quad (3.6.3)$$

3.7. Assume that  $X \sim \text{Binomial}(n, p)$  for some  $n \geq 1$  and  $0 < p < 1$  and  $Y \sim \text{poisson}(\lambda)$  for some  $\lambda > 0$ . Suppose  $E[X] = E[Y]$ . Then

- a)  $\text{var}(X) = \text{var}(Y)$   
b)  $\text{var}(X) < \text{var}(Y)$   
c)  $\text{var}(Y) < \text{var}(X)$   
d)  $\text{Var}(X)$  may be larger or smaller than  $\text{Var}(Y)$  depending on the values of  $n, p$  and  $\lambda$

**Solution:** For the random variable

$$X \sim \text{Binomial}(n, p)$$

As we know,

$$E[X] = np \quad (3.7.1)$$

$$\text{Var}(X) = np(1 - p) \quad (3.7.2)$$

for the random variable

$$Y \sim \text{poisson}(\lambda)$$

As we know,

$$E[Y] = \lambda \quad (3.7.3)$$

$$\text{Var}(Y) = \lambda \quad (3.7.4)$$

given that,

$$E[X] = E[Y] \quad (3.7.5)$$

$$np = \lambda \quad (3.7.6)$$

from (3.7.2),

$$\text{Var}(X) = np(1 - p) \quad (3.7.7)$$

using (3.7.6),

$$\text{Var}(X) = \lambda(1 - p) \quad (3.7.8)$$

using (3.7.4),

$$\text{Var}(X) = \text{Var}(Y)(1 - p) \quad (3.7.9)$$

$$\frac{\text{Var}(X)}{\text{Var}(Y)} = 1 - p \quad (3.7.10)$$

as,

$$1 - p < 1 \quad (3.7.11)$$

$$\frac{\text{Var}(X)}{\text{Var}(Y)} < 1 \quad (3.7.12)$$

$$\text{Var}(X) < \text{Var}(Y) \quad (3.7.13)$$

$\therefore \text{Var}(Y) > \text{Var}(X)$ , independent of  $n, p$  and  $\lambda$ .

- a)  $\text{var}(X) = \text{var}(Y)$

using TABLE 3.7.1,

$$\lambda(1 - p) = \lambda \quad (3.7.14)$$

$$1 - p = 1 \quad (3.7.15)$$

$$p = 0 \quad (3.7.16)$$

which is wrong as per the question ( $0 < p < 1$ ). hence the option is incorrect.

- b)  $\text{var}(X) < \text{var}(Y)$

using TABLE 3.7.1,

$$\lambda(1 - p) < \lambda \quad (3.7.17)$$

$$1 - p < 1 \quad (3.7.18)$$

$$p > 0 \quad (3.7.19)$$

which is true as per the question ( $0 < p < 1$ ). hence the option is correct.

- c)  $\text{var}(Y) < \text{var}(X)$

using TABLE 3.7.1,

$$\lambda(1 - p) > \lambda \quad (3.7.20)$$

$$1 - p > 1 \quad (3.7.21)$$

$$p < 0 \quad (3.7.22)$$

which is wrong as per the question ( $0 < p < 1$ ). hence the option is incorrect.

- d)  $\text{Var}(X)$  may be larger or smaller than  $\text{Var}(Y)$  depending on the values of  $n, p$  and  $\lambda$ .

Wrong, since we have shown that irrespective of the values of  $\lambda, n$ , and  $p$ ,  $\text{var}(y) > \text{var}(x)$

3.8. Let  $X$  be a non-negative integer valued random variable with probability mass function  $f(x)$  satisfying

$(x + 1)f(x + 1) = (\alpha + \beta x)f(x)$ ,  $x = 0, 1, 2, \dots$ ;  $\beta \neq 1$ . You may assume that  $E(X)$  and  $\text{Var}(X)$  exist. Then which of the following statements are true?

TABLE 3.7.1: Mean and Variance for random variables  $X$  and  $Y$

	$X$	$Y$
E	$\lambda$	$\lambda$
var	$\lambda(1 - p)$	$\lambda$

a)  $E(X) = \frac{\alpha}{1-\beta}$

b)  $E(X) = \frac{\alpha^2}{(1-\beta)(1+\alpha)}$

c)  $Var(X) = \frac{\alpha^2}{(1-\beta)^2}$

d)  $Var(X) = \frac{\alpha}{(1-\beta)^2}$

**Solution:** For a discrete random variable  $X$  with P.D.F.  $f(x)$  and which can take values from a set  $\mathbb{S}$ ,

$$E(X) = \sum_{x \in \mathbb{S}} xf(x) \quad (3.8.1)$$

And,

$$E(X^2) = \sum_{x \in \mathbb{S}} x^2 f(x) \quad (3.8.2)$$

Also, as  $f(x)$  is the P.D.F.,

$$\sum_{x \in \mathbb{S}} f(x) = 1 \quad (3.8.3)$$

Given, for  $x \in \mathbb{S} = \{0, 1, 2, \dots, n\}$ ,

$$(x+1)f(x+1) = (\alpha + \beta x)f(x) \quad (3.8.4)$$

Summing both sides for  $x \in \mathbb{S}$  we get,

$$\sum_{x=0}^n (x+1)f(x+1) = \sum_{x=0}^n (\alpha + \beta x)f(x) \quad (3.8.5)$$

Replacing  $x+1$  with  $x$  in L.H.S. we get,

$$\sum_{x=1}^{n+1} xf(x) = \sum_{x=0}^n (\alpha + \beta x)f(x) \quad (3.8.6)$$

Rewriting LHS, we get,

$$\sum_{x=0}^n xf(x) + (n+1)f(n+1) = \sum_{x=0}^n (\alpha + \beta x)f(x) \quad (3.8.7)$$

But as  $x \in \{0, 1, 2, \dots, n\}$ ,  $f(n+1) = 0$ . So the equation becomes

$$\sum_{x=0}^n xf(x) = \alpha \sum_{x=0}^n f(x) + \beta \sum_{x=0}^n xf(x) \quad (3.8.8)$$

Using (3.8.1) and (3.8.3), we get,

$$E(X) = \alpha(1) + \beta E(X) \quad (3.8.9)$$

So,

$$E(X) = \frac{\alpha}{1-\beta} \quad (3.8.10)$$

Now in (3.8.4), multiplying both sides by  $(x+1)$ , we get,

$$(x+1)^2 f(x+1) = (\alpha + \beta x)(x+1)f(x) \quad (3.8.11)$$

Summing both sides for  $x \in \mathbb{S}$  we get,

$$\sum_{x=0}^n (x+1)^2 f(x+1) = \sum_{x=0}^n (\alpha + \beta x)(x+1)f(x) \quad (3.8.12)$$

Replacing  $x+1$  with  $x$  in L.H.S. we get,

$$\sum_{x=1}^{n+1} x^2 f(x) = \sum_{x=0}^n (\beta x^2 f(x) + (\alpha + \beta)x f(x) + \alpha f(x)) \quad (3.8.13)$$

Rewriting LHS similarly as before, we get,

$$\begin{aligned} \sum_{x=0}^n x^2 f(x) &= \beta \sum_{x=0}^n x^2 f(x) + \\ &+ (\alpha + \beta) \sum_{x=0}^n x f(x) + \alpha \sum_{x=0}^n f(x) \end{aligned} \quad (3.8.14)$$

Using (3.8.1), (3.8.2) and (3.8.3), we get,

$$E(X^2) = \beta E(X^2) + (\alpha + \beta)E(X) + \alpha(1) \quad (3.8.15)$$

Using (3.8.10)

$$E(X^2)(1-\beta) = \frac{\alpha(\alpha + \beta)}{1-\beta} + \alpha \quad (3.8.16)$$

So,

$$E(X^2) = \frac{\alpha^2 + \alpha}{(1-\beta)^2} \quad (3.8.17)$$

Now,

$$Var(X) = E(X^2) - (E(X))^2 \quad (3.8.18)$$

Using (3.8.10) and (3.8.17),

$$Var(X) = \frac{\alpha^2 + \alpha}{(1-\beta)^2} - \frac{\alpha^2}{(1-\beta)^2} \quad (3.8.19)$$

So,

$$Var(X) = \frac{\alpha}{(1-\beta)^2} \quad (3.8.20)$$

So, options 1 and 4 are correct.

3.9. Let  $X$  be a random variable with probability density function,

$$f(x) = \alpha(x - \mu)^{\alpha-1} e^{-(x-\mu)^\alpha} \quad (3.9.1)$$

such that  $-\infty < \mu < \infty$ ;  $\alpha > 0$ ;  $x > \mu$ , The hazard function is:

- a) constant for all  $\alpha$
- b) an increasing function for some  $\alpha$
- c) independent of  $\alpha$
- d) independent of  $\mu$  when  $\alpha = 1$

**Solution:** Given PDF of  $X$ ,

$$f(x) = \alpha(x - \mu)^{\alpha-1} e^{-(x-\mu)^\alpha} \quad (3.9.2)$$

**Important property**(using in (3.9.8) as  $x > \mu$ ): Given  $x - y > 0$  and  $-\infty < y < \infty$ , then

$$\lim_{x \rightarrow -\infty} x - y = 0 \quad (3.9.3)$$

CDF of  $X$ ,

$$F(x) = \int_{-\infty}^x f(x) dx \quad (3.9.4)$$

$$= \int_{-\infty}^x \alpha(x - \mu)^{\alpha-1} e^{-(x-\mu)^\alpha} dx \quad (3.9.5)$$

$$= \int_{-\infty}^x e^{-(x-\mu)^\alpha} d(x - \mu)^\alpha \quad (3.9.6)$$

$$= \left[ \frac{e^{-(x-\mu)^\alpha}}{-1} \right]_{-\infty}^x \quad (3.9.7)$$

$$= -e^{-(x-\mu)^\alpha} - \lim_{x \rightarrow -\infty} \frac{e^{-(x-\mu)^\alpha}}{-1} \quad (3.9.8)$$

$$= -e^{-(x-\mu)^\alpha} + e^{-(0)^\alpha} \quad (3.9.9)$$

$$F(x) = 1 - e^{-(x-\mu)^\alpha} \quad (3.9.10)$$

Hazard function  $\beta(x)$ , (using (3.9.2) and (3.9.10))

$$\beta(x) = \frac{f(x)}{1 - F(x)} \quad (3.9.11)$$

$$= \frac{\alpha(x - \mu)^{\alpha-1} e^{-(x-\mu)^\alpha}}{1 - (1 - e^{-(x-\mu)^\alpha})} \quad (3.9.12)$$

$$= \frac{\alpha(x - \mu)^{\alpha-1} e^{-(x-\mu)^\alpha}}{e^{-(x-\mu)^\alpha}} \quad (3.9.13)$$

$$\beta(x) = \alpha(x - \mu)^{\alpha-1} \quad (3.9.14)$$

- a)  $\beta(x)$  is not constant for all  $\alpha$
- b)  $\beta(x) = \alpha(x - \mu)^{\alpha-1}$  is an increasing function for  $\alpha < 0$  or  $\alpha > 1$  as given  $x - \mu > 0$  for all  $x$ .

**Proof:** Using first derivative test, A function is increasing iff its first derivative is positive for all  $x$ .

$$\frac{d}{dx} \beta(x) = \frac{d}{dx} \alpha(x - \mu)^{\alpha-1} \quad (3.9.15)$$

$$= \alpha(\alpha - 1)(x - \mu)^{\alpha-2} \quad (3.9.16)$$

For (3.9.16) to be positive, (As given  $x - \mu > 0$ )

$$\alpha(\alpha - 1)(x - \mu)^{\alpha-2} > 0 \quad (3.9.17)$$

$$\alpha(\alpha - 1) > 0 \quad (3.9.18)$$

$$\Rightarrow \alpha \in (-\infty, 0) \cup (1, \infty) \quad (3.9.19)$$

$\therefore \beta(x)$  an increasing function for some  $\alpha$

c)  $\beta(x)$  is dependent of  $\alpha$


d) when  $\alpha = 1$ ,

$$\beta(x) = \alpha(x - \mu)^0 = \alpha \quad (3.9.20)$$

Therefore the hazard function is independent of  $\mu$  when  $\alpha = 1$ .

**ANSWER: (2) and (4)**

3.10. A point is chosen at random from a circular disc shown below. What is the probability that the point lies in the sector OAB?



( where  $\angle AOB = x$  radians )

- a)  $\frac{2x}{\pi}$
- b)  $\frac{x}{\pi}$
- c)  $\frac{x}{2\pi}$
- d)  $\frac{x}{4\pi}$

**Solution:**

Let  $X \in \{0, 1\}$  be a random variable such that  $X=0$  means we choose a point lying in sector OAB and  $X=1$  means that we choose a point lying outside sector OAB and inside the circle.

Area of a sector subtending an angle  $\theta$  at the

centre of circle with radius  $a$  is given by :

$$A = \frac{1}{2}a^2\theta \quad (3.10.1)$$

where  $\theta$  is in radians.

Let the radius of circle shown in figure be  $r$ . It is given that sector OAB subtends an angle of  $x$  radians at the centre of the circle.

Probability that the chosen point lies in sector OAB is:

$$\Pr(X = 0) = \frac{\text{Area of sector OAB}}{\text{Area of circle}} \quad (3.10.2)$$

$$= \frac{\frac{1}{2}r^2x}{\pi r^2} \quad (3.10.3)$$

$$= \frac{x}{2\pi} \quad (3.10.4)$$

$\therefore$  The correct answer is **option (3)**  $\frac{x}{2\pi}$ .

#### ALTERNATE SOLUTION

The joint pdf is given by:

$$f_{r\theta}(r, \theta) = \begin{cases} \frac{r}{\pi R^2} & \text{if } 0 < r < R, 0 < \theta < 2\pi \\ 0 & \text{otherwise} \end{cases} \quad (3.10.5)$$

Let  $A \equiv (R, \theta_2)$  and  $B \equiv (R, \theta_1)$ .

Hence,

$$(\theta_2 - \theta_1) = x \quad (3.10.6)$$

We want  $\theta \in (\theta_1, \theta_2)$  and  $r \in (0, R)$  for point to lie in the sector. Let the point to be chosen be  $(r, \theta)$ .

So, Required probability is:

$$\begin{aligned} \Pr(\theta_1 < \theta < \theta_2, 0 < r < R) \\ &= \int_{\theta_1}^{\theta_2} \int_0^R \frac{r}{\pi R^2} dr d\theta \\ &= \frac{1}{2\pi} (\theta_2 - \theta_1) = \frac{x}{2\pi} \end{aligned} \quad (3.10.7)$$

$$= \int_{\theta_1}^{\theta_2} \frac{1}{\pi R^2} \frac{r^2}{2} \Big|_0^R d\theta \quad (3.10.8)$$

$$= \int_{\theta_1}^{\theta_2} \frac{R^2}{2\pi R^2} d\theta \quad (3.10.9)$$

$$= \int_{\theta_1}^{\theta_2} \frac{1}{2\pi} d\theta \quad (3.10.10)$$

$$= \frac{\theta}{2\pi} \Big|_{\theta_1}^{\theta_2} \quad (3.10.11)$$

$$= \frac{\theta_2 - \theta_1}{2\pi} \quad (3.10.12)$$

$$= \frac{x}{2\pi} \quad (3.10.13)$$

$\therefore$  The correct answer is **option (3)**  $\frac{x}{2\pi}$ .

3.11. Let  $X$  and  $Y$  be independent random variables each following a uniform distribution on  $(0, 1)$ . Let  $W = XI_{\{Y \leq X^2\}}$ , where  $I_A$  denotes the indicator function of set  $A$ . Then which of the following statements are true?

a) The cumulative distribution function of  $W$  is given by

$$F_W(t) = t^2 I_{\{0 \leq t \leq 1\}} + I_{\{t > 1\}} \quad (3.11.1)$$

b)  $P[W > 0] = \frac{1}{3}$

c) The cumulative distribution function of  $W$  is continuous

d) The cumulative distribution function of  $W$  is given by

$$F_W(t) = \left( \frac{2 + t^3}{3} \right) I_{\{0 \leq t \leq 1\}} + I_{\{t > 1\}} \quad (3.11.2)$$

**Solution:**

Given  $X$  and  $Y$  are two independent random variables.

Given  $W = XI_{\{Y \leq X^2\}}$   
 $X \in (0, 1)$ ,  $Y \in (0, 1)$ ,  $W \in [0, 1]$

a) We need to find CDF of  $W$

i) The PDF for  $X$  is

$$p_X(x) = \begin{cases} 1 & 0 < x < 1 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.3)$$

ii) The CDF for  $X$  is

$$F_X(x) = \begin{cases} 0 & x \leq 0 \\ x & 0 < x < 1 \\ 1 & \text{otherwise} \end{cases} \quad (3.11.4)$$

iii) The PDF for  $Y$  is

$$p_Y(y) = \begin{cases} 1 & 0 < y < 1 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.5)$$

iv) The CDF for  $Y$  is

$$F_Y(y) = \begin{cases} 0 & y \leq 0 \\ y & 0 < y < 1 \\ 1 & \text{otherwise} \end{cases} \quad (3.11.6)$$

v)  $I_{\{Y \leq X^2\}}$  is defined as follows

$$I_{\{Y \leq X^2\}} = \begin{cases} 1 & y \leq x^2 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.7)$$

vi)  $W$  is defined as follows

$$W = \begin{cases} x & y \leq x^2 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.8)$$

From (3.11.8)

$$p_W(W = 0) = \Pr(I_{\{Y \leq X^2\}} = 0) \quad (3.11.9)$$

$$= \Pr(x^2 < y) \quad (3.11.10)$$

vii) Let  $Z = X^2 - Y$  be a random variable where  $Z \in (-1, 1)$

$$F_{X^2}(u) = \Pr(X^2 \leq u) \quad (3.11.11)$$

$$= \Pr(X \leq \sqrt{u}) \quad (3.11.12)$$

$$= F_X(\sqrt{u}) \quad (3.11.13)$$

A) From (3.11.4), The CDF for  $X^2$  is

$$F_{X^2}(u) = \begin{cases} 0 & u \leq 0 \\ \sqrt{u} & 0 < u < 1 \\ 1 & \text{otherwise} \end{cases} \quad (3.11.14)$$

B) The PDF for  $X^2$  is

$$p_{X^2}(u) = \begin{cases} \frac{1}{2\sqrt{u}} & 0 < u < 1 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.15)$$

$$F_{\{-Y\}}(v) = \Pr(-Y \leq v) \quad (3.11.16)$$

$$= \Pr(Y \geq -v) \quad (3.11.17)$$

$$= 1 - F_Y(-v) \quad (3.11.18)$$

C) From (3.11.6), The CDF for  $(-Y)$  is

$$F_{\{-Y\}}(v) = \begin{cases} 0 & v \leq -1 \\ 1 + v & -1 < v < 0 \\ 1 & \text{otherwise} \end{cases} \quad (3.11.19)$$

D) The PDF for  $(-Y)$  is

$$p_{\{-Y\}}(v) = \begin{cases} 1 & -1 < v < 0 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.20)$$

E)  $Z = X^2 - Y \implies z = u + v$   
Using convolution

$$p_Z(z) = \int_{-\infty}^{\infty} p_{X^2}(z - v) p_{\{-Y\}}(v) dv \quad (3.11.21)$$

Solving (3.11.21) using (3.11.20), (3.11.15) for  $z \in (-1, 1)$ , we get PDF of  $Z$  as follows

$$p_Z(z) = \begin{cases} \sqrt{z+1} & -1 < z \leq 0 \\ 1 - \sqrt{z} & 0 < z < 1 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.22)$$

F) CDF of  $Z$  as follows

$$F_Z(z) = \begin{cases} \frac{2}{3}(z+1)^{\frac{3}{2}} & -1 < z \leq 0 \\ z - \frac{2}{3}z^{\frac{3}{2}} & 0 < z < 1 \\ 1 & \text{otherwise} \end{cases} \quad (3.11.23)$$

viii) using (3.11.23) to find  $p_W(W = 0)$

$$p_W(W = 0) = \Pr(x^2 < y) \quad (3.11.24)$$

$$= F_z(0) \quad (3.11.25)$$

$$= \frac{2}{3} \quad (3.11.26)$$

ix)  $W = t \implies X = t$  where  $t \in (0, 1)$

$$p_W(t) = \int_{-\infty}^{\infty} p_X(t) I_{\{y \leq t^2\}} dy \quad (3.11.27)$$

$$0 < y < 1 \quad (3.11.28)$$

$$0 < y \leq t^2 \quad (3.11.29)$$

For  $0 < t < 1$ ,

$$p_W(t) = \int_0^{t^2} p_X(t) I_{\{y \leq t^2\}} dy \quad (3.11.30)$$

$$= t^2 \quad (3.11.31)$$

x)  $\therefore$  PDF of  $W$  is as follows

$$p_W(t) = \begin{cases} \frac{2}{3} & t = 0 \\ t^2 & 0 < t < 1 \\ 0 & \text{otherwise} \end{cases} \quad (3.11.32)$$

xi) The CDF of  $W$  is as follows:

$$F_W(t) = \begin{cases} 0 & t < 0 \\ \frac{2+t^3}{3} & 0 \leq t \leq 1 \\ 1 & \text{otherwise} \end{cases} \quad (3.11.33)$$

b) We need to find  $P[W > 0]$

$$\Pr(W > 0) = 1 - F_W(0) \quad (3.11.34)$$

$$= \frac{1}{3} \quad (3.11.35)$$

$$\therefore \Pr(W > 0) = \frac{1}{3} \quad (3.11.36)$$




c) CDF of  $W$  is discontinuous at  $W = 0$ .

$\therefore$  option 3 is incorrect.

d) The CDF in (3.11.33) can be written as

$$F_W(t) = \left( \frac{2+t^3}{3} \right) I_{\{0 \leq t \leq 1\}} + I_{\{t > 1\}} \quad (3.11.37)$$

$\therefore$  option 2 and 4 are correct.



( where  $\angle AOB = x$  radians )

3.12. A point is chosen at random from a circular disc shown below.

What is the probability that the point lies in the sector OAB?



a)  $\frac{2x}{\pi}$   
b)  $\frac{x}{\pi}$

c)  $\frac{x}{2\pi}$   
d)  $\frac{x}{4\pi}$

**Solution:**

Let  $X \in \{0, 1\}$  be a random variable such that  $X=0$  means we choose a point lying in sector OAB and  $X=1$  means that we choose a point lying outside sector OAB and inside the circle. Area of a sector subtending an angle  $\theta$  at the centre of circle with radius  $a$  is given by :

$$A = \frac{1}{2}a^2\theta \quad (3.12.1)$$

where  $\theta$  is in radians.

Let the radius of circle shown in figure be  $r$ . It is given that sector OAB subtends an angle of  $x$  radians at the centre of the circle. Probability that the chosen point lies in sector OAB is:

$$\Pr(X = 0) = \frac{\text{Area of sector OAB}}{\text{Area of circle}} \quad (3.12.2)$$

$$= \frac{\frac{1}{2}r^2x}{\pi r^2} \quad (3.12.3)$$

$$= \frac{x}{2\pi} \quad (3.12.4)$$

$\therefore$  The correct answer is **option (3)  $\frac{x}{2\pi}$** . **alternate solution** The

joint pdf is given by:

$$f_{r\theta}(r, \theta) = \begin{cases} \frac{r}{\pi R^2} & \text{if } 0 < r < R, 0 < \theta < 2\pi \\ 0 & \text{otherwise} \end{cases} \quad (3.12.5)$$

Let  $A \equiv (R, \theta_2)$  and  $B \equiv (R, \theta_1)$ .

Hence,

$$(\theta_2 - \theta_1) = x \quad (3.12.6)$$

We want  $\theta \in (\theta_1, \theta_2)$  and  $r \in (0, R)$  for point to lie in the sector.

Let the point to be chosen be  $(r, \theta)$ .

So, Required probability is:

$$\Pr(\theta_1 < \theta < \theta_2, 0 < r < R)$$

$$= \int_{\theta_1}^{\theta_2} \int_0^R \frac{r}{\pi R^2} dr d\theta \quad (3.12.7)$$

$$= \int_{\theta_1}^{\theta_2} \frac{1}{\pi R^2} \frac{r^2}{2} \Big|_0^R \frac{2}{\pi \sqrt{1-y^2}} d\theta \quad \text{Solution: From the given information,}$$

$$F_X(x) = \Pr(X \leq x) \quad (4.1.2)$$

$$= \begin{cases} 0 & x \leq 0 \\ \frac{x^2}{\pi^2} & 0 < x < \pi \\ 1 & x \geq \pi \end{cases} \quad (4.1.3)$$

$$= \int_{\theta_1}^{\theta_2} \frac{R^2}{2\pi R^2} d\theta \quad \text{after integration. Consequently,}$$

$$F_Y(y) = \Pr(Y \leq y) \quad (4.1.4)$$

$$= \Pr(\sin X \leq y) \quad (4.1.5)$$

$$= \int_{\theta_1}^{\theta_2} \frac{1}{2\pi} d\theta \quad \text{From Fig. 4.1.1,}$$

$$(3.12.10)$$

$$= \frac{\theta}{2\pi} \Big|_{\theta_1}^{\theta_2}$$


$$(3.12.11)$$

$$= \frac{\theta_2 - \theta_1}{2\pi}$$

$$(3.12.12)$$

$$= \frac{x}{2\pi}$$

$$(3.12.13)$$

Fig. 4.1.1:  $Y = \sin X$  plot

$\therefore$  The correct answer is **option**  
**(3)**  $\frac{x}{2\pi}$ .

#### 4 TRANSFORMATION OF VARIABLES

4.1. Let  $X$  be a random variable with pdf

$$f_X(x) = \begin{cases} \frac{2x}{\pi^2} & 0 < x < \pi \\ 0 & \text{otherwise} \end{cases} \quad (4.1.1)$$

Let  $Y = \sin X$ , then for  $0 < y < 1$ , the pdf of  $Y$  is given by,

- (A)  $\frac{2\pi}{\sqrt{1-y^2}}$   
 (B)  $\frac{\pi}{2} \sqrt{1-y^2}$   
 (C)  $\frac{2}{\pi} \sqrt{1-y^2}$

$$\sin X \leq y$$

$$\Rightarrow \{X \leq \sin^{-1} y \cup X \geq \pi - \sin^{-1} y\} \quad (4.1.6)$$

$$\Rightarrow F_Y(y) = \Pr(X \leq \sin^{-1} y) + \Pr(X \geq \pi - \sin^{-1} y) \quad (4.1.7)$$

$$= F_X(\sin^{-1} y) + 1 - \Pr(X \leq \pi - \sin^{-1} y) \quad (4.1.8)$$

$$\Rightarrow F_Y(y) = 1 + F_X(\sin^{-1} y) - F_X(\pi - \sin^{-1} y) \quad (4.1.9)$$

Substituting from (4.1.3) in (4.1.9)

$$F_Y(y) = \frac{(\sin^{-1} y)^2}{\pi^2} + 1 - \frac{(\pi - \sin^{-1} y)^2}{\pi^2} \quad (4.1.10)$$

$$= \frac{2 \sin^{-1} y}{\pi} \quad (4.1.11)$$

$$\therefore f_Y(y) = \frac{dF_Y(y)}{dy} \quad (4.1.12)$$

$$= \frac{2}{\pi \sqrt{1-y^2}} \quad (4.1.13)$$

Hence, option(D) is correct.

## 5 INDEPENDENT RANDOM VARIABLES

5.1. Let  $X \in \{0, 1\}$  and  $Y \in \{0, 1\}$  be two independent binary random variables. If  $\Pr(X = 0) = p$  and  $\Pr(Y = 0) = q$ , then  $\Pr(X + Y \geq 1)$  is equal to

- a)  $pq + (1-p)(1-q)$
- b)  $pq$
- c)  $p(1-q)$
- d)  $1-pq$

**Solution:**

5.2. Two independent random variables  $X$  and  $Y$  are uniformly distributed in the interval  $[-1, 1]$ . The probability that  $\max(X, Y)$  is less than  $\frac{1}{2}$  is

- a)  $\frac{3}{4}$
- b)  $\frac{9}{16}$
- c)  $\frac{1}{4}$
- d)  $\frac{2}{3}$

**Solution:**

5.3. Suppose that  $X_1, X_2, X_3, \dots, X_{10}$  are i.i.d,  $N(0,1)$ . Which of the following statements are correct ?

- (A)  $\Pr(X_1 > X_2 + X_3 + \dots + X_{10}) = \frac{1}{2}$
- (B)  $\Pr(X_1 > X_2 X_3 \dots X_{10}) = \frac{1}{2}$
- (C)  $\Pr(\sin X_1 > \sin X_2 + \sin X_3 + \dots + \sin X_{10}) = \frac{1}{2}$
- (D)  $\Pr(\sin X_1 > \sin(X_2 + X_3 + \dots + X_{10})) = \frac{1}{2}$

**Solution:**

**Lemma 5.1.** If  $X \sim N(0, 1)$  then  $Y = -X$  also follows standard normal distribution.

*Proof.*

$$P(Y \leq u) = P(-X \leq u) \quad (5.3.1)$$

$$= P(X > -u) \quad (5.3.2)$$

$$= 1 - P(X \leq -u) \quad (5.3.3)$$

$$= 1 - (1 - P(X \leq u)) \quad (5.3.4)$$

$$= P(X \leq u) \quad (5.3.5)$$

As the distribution is symmetric,

$$P(X \leq -u) = P(X \geq u) = 1 - P(X \leq u) \quad (5.3.6)$$

□

**Lemma 5.2.** If  $n$  is an even number and  $g(x)$  is an odd function, then,

a)

$$\begin{aligned} \Pr\left(g(X_1) > \sum_{k=2}^n g(X_k)\right) \\ = \Pr\left(g(X_1) < \sum_{k=2}^n g(X_k)\right) \\ = \frac{1}{2} \end{aligned} \quad (5.3.7)$$

b)

$$\begin{aligned} \Pr\left(g(X_1) > \prod_{k=2}^n g(X_k)\right) \\ = \Pr\left(g(X_1) < \prod_{k=2}^n g(X_k)\right) = \frac{1}{2} \end{aligned} \quad (5.3.8)$$

*Proof.* a)

$$\begin{aligned} \Pr\left(g(X_1) > \sum_{k=2}^n g(X_k)\right) \\ = \Pr\left(g(-X_1) < \sum_{k=2}^n g(-X_k)\right) \\ = \Pr\left(g(X_1) < \sum_{k=2}^n g(X_k)\right) \end{aligned} \quad (5.3.9)$$

As the cases

$$g(X_1) > \sum_{k=2}^n g(X_k) \quad (5.3.10)$$

and

$$g(X_1) < \sum_{k=2}^n g(X_k) \quad (5.3.11)$$

are complementary to each other,

$$\Pr\left(g(X_1) > \sum_{k=2}^n g(X_k)\right) = \frac{1}{2} \quad (5.3.12)$$

b) Similarly,

$$\begin{aligned} \Pr\left(g(X_1) > \prod_{k=2}^n g(X_k)\right) \\ = \Pr\left(g(-X_1) < \prod_{k=2}^n g(-X_k)\right) \\ = \Pr\left(g(X_1) < \prod_{k=2}^n g(X_k)\right) \end{aligned} \quad (5.3.13)$$

As they follow the same distribution, the above expression is true. Thus we have

$$\Pr\left(g(X_1) > \prod_{k=2}^n g(X_k)\right) = \Pr\left(g(X_1) < \prod_{k=2}^n g(X_k)\right) \quad (5.3.14)$$

As the cases

$$g(X_1) > \prod_{k=2}^n g(X_k) \quad (5.3.15)$$

and

$$g(X_1) < \prod_{k=2}^n g(X_k) \quad (5.3.16)$$

are complementary to each other and from (5.3.7) we have

$$\Pr\left(g(X_1) > \prod_{k=2}^n g(X_k)\right) = \frac{1}{2} \quad (5.3.17)$$

(A) From (5.3.12), taking  $g(x) = x$ ,

$$\Pr(X_1 > X_2 + \dots + X_{10}) = \frac{1}{2} \quad (5.3.18)$$

(B) From (5.3.17) taking  $g(x) = x$

$$\Pr(X_1 > X_2 X_3 \dots X_{10}) = \frac{1}{2} \quad (5.3.19)$$

(C) From (5.3.12) taking  $g(x) = \sin x$

$$\Pr(\sin X_1 > \sin X_2 + \dots + \sin X_{10}) = \frac{1}{2} \quad (5.3.20)$$

(D)

$$\begin{aligned} \Pr(\sin X_1 > \sin(X_2 + \dots + X_{10})) \\ = \Pr(\sin(-X_1) < \sin(-X_2 - \dots - X_{10})) \\ = \Pr(\sin X_1 < \sin(X_2 + \dots + X_{10})) \end{aligned} \quad (5.3.21)$$

As they follow the same distribution, the above expression is true. Thus we have

$$\begin{aligned} \Pr(\sin X_1 > \sin(X_2 + \dots + X_{10})) \\ = \Pr(\sin X_1 < \sin(X_2 + \dots + X_{10})) \end{aligned} \quad (5.3.22)$$

Also, as  $X_1$  is a continuous random variable

$$\Pr(\sin X_1 = \sin(X_2 + \dots + X_{10})) = 0 \quad (5.3.23)$$

As the cases

$$X_1 > X_2 + \dots + X_{10} \quad (5.3.24)$$

and

$$X_1 < X_2 + \dots + X_{10} \quad (5.3.25)$$

are complementary to each other

$$\Pr(\sin X_1 > \sin(X_2 + \dots + X_{10})) = \frac{1}{2} \quad (5.3.26)$$

□

5.4. Which of the following conditions imply independence of the random variables  $X$  and  $Y$ ?

a)  $\Pr(X > a | Y > a) = \Pr(X > a) \quad \forall a \in \mathbb{R}$

b)  $\Pr(X > a | Y < b) = \Pr(X > a) \quad \forall a, b \in \mathbb{R}$

c)  $X$  and  $Y$  are uncorrelated.

d)  $E[(X-a)(Y-b)] = E(X-a)E(Y-b) \quad \forall a, b \in \mathbb{R}$

**Solution:**

**Definition 1.** Two random variables  $X$  and  $Y$  are independent when the joint probability distribution of random variables is product of

their individual probability distributions i.e for all sets  $A, B$

$$\Pr(X \in A, Y \in B) = \Pr(X \in A) \Pr(Y \in B) \quad (5.4.1)$$

Alternatively,

$$F_{X,Y}(a, b) = F_X(a) F_Y(b) \quad (5.4.2)$$

**Lemma 5.3.** From (5.4.2), it follows that

$$\implies f_{X,Y}(a, b) = f_X(a) f_Y(b) \quad (5.4.3)$$

*Proof.* From (5.4.2),

$$\frac{\partial^2 F_{X,Y}(a, b)}{\partial b \partial a} = \frac{\partial F_X(a)}{\partial a} \frac{\partial F_Y(b)}{\partial b} \quad (5.4.4)$$

yielding (5.4.3).  $\square$

a) From the given information,

$$\Pr(X > a, Y > a) = \Pr(X > a) \Pr(Y > a) \quad (5.4.5)$$

$$= [1 - F_X(a)] [1 - F_Y(a)] \quad (5.4.6)$$

$$\begin{aligned} & \because \Pr(X > a) - \Pr(Y < a) \\ &= \Pr(X > a, Y > a) + \Pr(X > a, Y < a) \\ & - \Pr(X > a, Y < a) - \Pr(X < a, Y < a), \\ & \Pr(X > a, Y > a) = 1 - F_X(a) - F_Y(a) \\ & \quad + F_{X,Y}(a, a) \end{aligned} \quad (5.4.7)$$

which, upon substituting from (5.4.6) yields

$$\implies F_{X,Y}(a, a) = F_X(a) F_Y(a) \quad (5.4.8)$$

which is a special case of (5.4.1) for  $b = a$ . The spectrum of conditions for independence is hence underrepresented. Thus, the given condition does not imply independence of  $X$  and  $Y$ .

*Option 1 is incorrect.*

b) From Bayes theorem,

$$\Pr(X > a | Y < b) = \Pr(X > a) \quad (5.4.9)$$

$$\begin{aligned} & \implies \Pr(X > a, Y < b) \\ &= \Pr(X > a) \Pr(Y < b) \end{aligned} \quad (5.4.10)$$

for all  $a, b \in R$ .

$$\begin{aligned} \because F_Y(b) &= \Pr(X > a, Y < b) \\ &+ \Pr(X < a, Y < b), \end{aligned} \quad (5.4.11)$$

$$\begin{aligned} \Pr(X > a, Y < b) &= F_Y(b) - F_{X,Y}(a, b) \\ \implies F_Y(b) - F_{X,Y}(a, b) &= (1 - F_X(a)) F_Y(b) \\ \text{or, } F_{X,Y}(a, b) &= F_X(a) F_Y(b) \end{aligned} \quad (5.4.12)$$

upon substituting from (5.4.10) and simplifying. Thus,  $X$  and  $Y$  are independent.

*Option 2 is correct.*

c) We prove through a counterexample.

**Definition 2.** Two random variables  $X$  and  $Y$  are uncorrelated if their covariance is zero, i.e.,

$$\text{cov}(X, Y) = E(XY) - E(X)E(Y) = 0 \quad (5.4.13)$$

Let  $X \sim U[-1, 1]$  be a uniformly distributed random variable such that

$$f_X(x) = \begin{cases} \frac{1}{2} & -1 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad (5.4.14)$$

$$E(X) = \int_{-1}^1 x f(x) dx = 0 \quad (5.4.15)$$

Let

$$Y = X^2. \quad (5.4.16)$$

so that  $X$  and  $Y$  are dependent. Then,

$$\text{cov}(X, Y) = E(XY) - E(X)E(Y) \quad (5.4.17)$$

$$= E(X^3) - 0 \times E(Y) \quad (5.4.18)$$

$$= \int_{-1}^1 x^3 f(x) dx = 0 \quad (5.4.19)$$

$X$  and  $Y$  are uncorrelated but not independent.

*Option 3 is incorrect*

d) Given that,

$$E((X - a)(Y - b)) = E(X - a)E(Y - b) \quad (5.4.20)$$

$$\begin{aligned} \Rightarrow \text{cov}(X - a, Y - b) &= \\ &= E((X - a)(Y - b)) \\ &= E(X - a)E(Y - b) \quad (5.4.21) \end{aligned}$$

$$\text{or, } \text{cov}(X - a, Y - b) = 0 = \text{cov}(X, Y) \quad (5.4.22)$$

From option 3, it follows that  $X$  and  $Y$  are not necessarily independent.

**Option 4 is incorrect.**

5.5. Let  $X$  and  $Y$  be two independent and identically distributed (I.I.D) random variables uniformly distributed in  $(0,1)$ . Let  $Z = \max(X, Y)$  and  $W = \min(X, Y)$ , then the probability that  $[Z - W > \frac{1}{2}]$  is

(A)  $\frac{1}{2}$

(B)  $\frac{3}{4}$

(C)  $\frac{1}{4}$

(D)  $\frac{2}{3}$  **Solution:**

$X$  and  $Y$  are two independent random variables. Let

$$f_X(x) = \Pr(X = x) \quad (5.5.1)$$

$$f_Y(y) = \Pr(Y = y) \quad (5.5.2)$$

$$f_V(v) = \Pr(V = v) \quad (5.5.3)$$

be the probability densities of random variables  $X$ ,  $Y$  and  $V = X - Y$ .

The density for  $X$  is

$$f_X(x) = \begin{cases} 1 & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad (5.5.4)$$

We have ,

$$V = X - Y \iff v = x - y \iff x = v + y \quad (5.5.5)$$

The density of  $X$  can also be represented as,

$$f_X(v + y) = \begin{cases} 1 & 0 \leq v + y \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad (5.5.6)$$

and the density of  $Y$  is,

$$f_Y(y) = \begin{cases} 1 & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad (5.5.7)$$

The density of  $V$  i.e.  $V = X - Y$  is given by the convolution of  $f_X(-v)$  with  $f_Y(v)$ .

$$f_V(v) = \int_{-\infty}^{\infty} f_X(v + y)f_Y(y) dy \quad (5.5.8)$$

From 5.5.6 and 5.5.7 we have,

The integrand is 1 when,

$$0 \leq y \leq 1 \quad (5.5.9)$$

$$0 \leq v + y \leq 1 \quad (5.5.10)$$

$$-v \leq y \leq 1 - v \quad (5.5.11)$$

and zero, otherwise.

Now when  $-1 \leq v \leq 0$  we have,

$$f_V(v) = \int_{-v}^1 dy \quad (5.5.12)$$

$$= (1 - (-v)) \quad (5.5.13)$$

$$= 1 + v \quad (5.5.14)$$

For  $0 \leq v \leq 1$  we have,

$$f_V(v) = \int_0^{1-v} dy \quad (5.5.15)$$

$$= (1 - v - (0)) \quad (5.5.16)$$

$$= 1 - v \quad (5.5.17)$$

Therefore the density of  $V$  is given by

$$f_V(v) = \begin{cases} 1 + v & -1 \leq v \leq 0 \\ 1 - v & 0 < v \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad (5.5.18)$$

The plot for PDF of  $V$  can be observed at figure 5.5.1




Fig. 5.5.1: The PDF of  $V$

The CDF of  $V$  is defined as,

$$F_V(v) = \Pr(V \leq v) \quad (5.5.19)$$

Now for  $v \leq 0$ ,

$$\Pr(V \leq v) = \int_{-\infty}^v f_V(v) dv \quad (5.5.20)$$

$$= \int_{-1}^v (1 + v) dv \quad (5.5.21)$$

$$= \left( \frac{v^2}{2} + v \right) \Big|_{-1}^v \quad (5.5.22)$$

$$= \left( \left( \frac{v^2}{2} + v \right) - \left( \frac{1}{2} - 1 \right) \right) \quad (5.5.23)$$

$$= \frac{v^2 + 2v + 1}{2} \quad (5.5.24)$$

Similarly for  $v \leq 1$ ,

$$\Pr(V \leq v) = \int_{-\infty}^v f_V(v) dv \quad (5.5.25)$$

$$= \frac{1}{2} + \int_0^v (1 - v) dz \quad (5.5.26)$$

$$= \frac{-v^2 + 2v + 1}{2} \quad (5.5.27)$$

The CDF is as below:

$$F_V(v) = \begin{cases} 0 & v < -1 \\ \frac{v^2 + 2v + 1}{2} & v \leq 0 \\ \frac{-v^2 + 2v + 1}{2} & v \leq 1 \\ 1 & v > 1 \end{cases} \quad (5.5.28)$$

The plot for CDF of  $V$  can be observed at figure 5.5.2

We need  $\Pr(Z - W > \frac{1}{2})$  where  $Z = \max(X, Y)$  and  $W = \min(X, Y)$ . Now,

$$Z - W = \begin{cases} X - Y & \text{for } X \geq Y \\ Y - X & \text{for } X < Y \end{cases} \quad (5.5.29)$$




Fig. 5.5.2: The CDF of  $V$

Therefore,

$$\Pr\left(Z - W > \frac{1}{2}\right) = \Pr\left(X - Y > \frac{1}{2}, X \geq Y\right) + \Pr\left(Y - X > \frac{1}{2}, X < Y\right) \quad (5.5.30)$$

$$= \Pr\left(X - Y > \frac{1}{2}\right) + \Pr\left(Y - X > \frac{1}{2}\right) \quad (5.5.31)$$

$$= \Pr\left(V > \frac{1}{2}\right) + \Pr\left(-V > \frac{1}{2}\right) \quad (5.5.32)$$

$$= 1 - \Pr\left(V \leq \frac{1}{2}\right) + \Pr\left(V < -\frac{1}{2}\right) \quad (5.5.33)$$

$$= 1 - F_V\left(\frac{1}{2}\right) + F_V\left(-\frac{1}{2}\right) \quad (5.5.34)$$

$$= 1 - \frac{7}{8} + \frac{1}{8} \quad (5.5.35)$$

$$= \frac{1}{4} \quad (5.5.36)$$

Hence the correct answer is option (C).

5.6. Let  $X$  and  $Y$  be independent exponential random variables. If  $E[X] = 1$  and  $E[Y] = \frac{1}{2}$  then  $\Pr(X > 2Y | X > Y)$  is

$$1. \frac{1}{2} \qquad 3. \frac{2}{3}$$

$$2. \frac{1}{3} \qquad 4. \frac{3}{4}$$

**Solution:** Since  $X$  and  $Y$  are exponential random variables with means'

$$E[X] = 1 \text{ and } E[Y] = \frac{1}{2} \quad (5.6.1)$$

Marginal PDFs of  $X$  and  $Y$  are given by

$$f_X(x) = e^{-x}, x > 0 \quad (5.6.2)$$

$$f_Y(y) = 2e^{-2y}, y > 0 \quad (5.6.3)$$

CDFs for  $X$  and  $Y$  are

$$F_X(b) = \int_0^b f_X(x) dx \quad (5.6.4)$$

$$= \int_0^b e^{-x} dx \quad (5.6.5)$$

$$= 1 - e^{-b} \quad (5.6.6)$$

$$F_Y(b) = \int_0^b f_Y(y) dy \quad (5.6.7)$$

$$= \int_0^b 2e^{-2y} dy \quad (5.6.8)$$

$$= [-e^{-2y}]_0^b \quad (5.6.9)$$

$$= 1 - e^{-2b} \quad (5.6.10)$$

Now,

$$\Pr(X > 2Y | X > Y) = \frac{\Pr(X > 2Y, X > Y)}{\Pr(X > Y)} \quad (5.6.11)$$

$$= \frac{\Pr(X > 2Y)}{\Pr(X > Y)} \quad (5.6.12)$$

$$\Pr(X > Y) = \Pr(Y < X) \quad (5.6.13)$$

$$= E[F_Y(X)] \quad (5.6.14)$$

$$= \int_0^\infty F_Y(X) f_X(x) dx \quad (5.6.15)$$

$$= \int_0^\infty (1 - e^{-2x}) e^{-x} dx \quad (5.6.16)$$

$$= \left[ \frac{e^{-x}}{-1} - \frac{e^{-3x}}{-3} \right]_0^\infty \quad (5.6.17)$$

$$= (0 + 1) + \frac{1}{3}(0 - 1) \quad (5.6.18)$$

$$= \frac{2}{3} \quad (5.6.19)$$

$$\Pr(X > 2Y) = \Pr\left(Y < \frac{X}{2}\right) \quad (5.6.20)$$

$$= E[F_Y(X/2)] \quad (5.6.21)$$

$$= \int_0^\infty F_Y(X/2) f_X(x) dx \quad (5.6.22)$$

$$= \int_0^\infty (1 - e^{-x}) e^{-x} dx \quad (5.6.23)$$

$$= \left[ \frac{e^{-x}}{-1} - \frac{e^{-2x}}{-2} \right]_0^\infty \quad (5.6.24)$$

$$= (0 + 1) + \frac{1}{2}(0 - 1) \quad (5.6.25)$$

$$= \frac{1}{2} \quad (5.6.26)$$

Putting (5.6.19) and (5.6.26) in (5.6.12)

$$\Pr(X > 2Y | X > Y) = \frac{1/2}{2/3} \quad (5.6.27)$$

$$= \frac{3}{4} \quad (5.6.28)$$

$\therefore$  Option 4 is the correct answer.

5.7. A fair die is thrown two times independently. Let  $X, Y$  be the outcomes of these two throws and  $Z = X + Y$ . Let  $U$  be the remainder obtained when  $Z$  is divided by 6. Then which of the following statement(s) is/are true?

- a)  $X$  and  $Z$  are independent
- b)  $X$  and  $U$  are independent
- c)  $Z$  and  $U$  are independent
- d)  $Y$  and  $Z$  are not independent

**Solution:** Let  $X \in \{1, 2, 3, 4, 5, 6\}$  represent the random variable which represents the outcome of the first throw of a dice. Similarly,  $Y \in \{1, 2, 3, 4, 5, 6\}$  represents the random variable which represents the outcome of the second throw of a dice.

$$n(X = i) = 1, \quad i \in \{1, 2, 3, 4, 5, 6\} \quad (5.7.1)$$

$$\Pr(X = i) = \begin{cases} \frac{1}{6} & i \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases} \quad (5.7.2)$$

Similarly,

$$\Pr(Y = i) = \begin{cases} \frac{1}{6} & i \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases} \quad (5.7.3)$$



$$Z = X + Y \quad (5.7.4)$$

$$\text{Let } z \in \{1, 2, \dots, 11, 12\} \quad (5.7.5)$$

$$\Pr(Z = z) = \Pr(X + Y = z) \quad (5.7.6)$$

$$= \sum_{x=0}^z \Pr(X = x) \Pr(Y = z - x) \quad (5.7.7)$$

$$= (6 - |z - 7|) \times \frac{1}{6} \times \frac{1}{6} \quad (5.7.8)$$

$$= \frac{6 - |z - 7|}{36} \quad (5.7.9)$$

$$\Pr(Z = z) = \begin{cases} \frac{6 - |z - 7|}{36} & z \in \{1, 2, \dots, 11, 12\} \\ 0 & \text{otherwise} \end{cases} \quad (5.7.10)$$

$U$  is the remainder obtained when  $Z$  is divided by 6.

$$\text{Let } u \in \{0, 1, 2, 3, 4, 5\} \quad (5.7.11)$$

$$\Pr(U = u) = \sum_{k=0}^2 \Pr(Z = 6k + u) \quad (5.7.12)$$

$$\Pr(U = 0) = \Pr(Z = 0) + \Pr(Z = 6) + \Pr(Z = 12) \quad (5.7.13)$$

$$= 0 + \frac{5}{36} + \frac{1}{36} = \frac{1}{6} \quad (5.7.14)$$

$$\text{for } u \in \{1, 2, 3, 4, 5\} \quad (5.7.15)$$

$$\Pr(U = u) = \Pr(Z = 0 + u) + \Pr(Z = 6 + u) \quad (5.7.16)$$

$$= \frac{6 - |u - 7|}{36} + \frac{6 - |6 + u - 7|}{36} \quad (5.7.17)$$

$$= \frac{6 - (7 - u)}{36} + \frac{6 - (u - 1)}{36} \quad (5.7.18)$$

$$= \frac{u - 1 + 7 - u}{36} = \frac{6}{36} \quad (5.7.19)$$

$$= \frac{1}{6} \quad (5.7.20)$$

$$\Pr(U = u) = \begin{cases} \frac{1}{6} & u \in \{0, 1, 2, 3, 4, 5\} \\ 0 & \text{otherwise} \end{cases} \quad (5.7.21)$$

Now, for checking each option,

a) Checking if  $X$  and  $Z$  are independent

$$p_1 = \Pr(Z = z, X = x) \quad (5.7.22)$$

$$= \Pr(Y = z - x, X = x) \quad (5.7.23)$$

$$= \Pr(Y = z - x) \times \Pr(X = x) \quad (5.7.24)$$

$$= \begin{cases} \frac{1}{36} & z - x \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases} \quad (5.7.25)$$

$$\Pr(Z = z) \times \Pr(X = x) = \frac{6 - |z - 7|}{36} \times \frac{1}{6} \quad (5.7.26)$$

$$= \frac{6 - |z - 7|}{216} \quad (5.7.27)$$

$$\Pr(Z = z) \Pr(X = x) \neq \Pr(Z = z, X = x) \quad (5.7.28)$$

$X$  and  $Z$  are not independent from (5.7.28) and hence option (5.7a) is false.

b) Checking if  $X$  and  $U$  are independent

$$p_2 = \Pr(U = u, X = x) \quad (5.7.29)$$

$$p_2 = \Pr((Z = u) + (Z = 6 + u) + (Z = 12 + u), X = x) \quad (5.7.30)$$

$$p_2 = \Pr((Y = u - x) + (Y = 6 + u - x) + (Y = 12 + u - x), X = x) \quad (5.7.31)$$

$$p_2 = \frac{1}{6} \times \frac{1}{6} \quad (5.7.32)$$

$$= \frac{1}{36} \quad (5.7.33)$$

$$\Pr(U = u) \times \Pr(X = x) = \frac{1}{6} \times \frac{1}{6} \quad (5.7.34)$$

$$= \frac{1}{36} \quad (5.7.35)$$

$$\Pr(U = u) \Pr(X = x) = \Pr(U = u, X = x) \quad (5.7.36)$$

$X$  and  $U$  are independent from (5.7.36) and hence option (5.7b) is true.

c) Checking if  $Z$  and  $U$  are independent

$$p_3 = \Pr(Z = z|U = u) \quad (5.7.37)$$

$$p_3 = \begin{cases} 1 & u = 1 \text{ and } z = 7 \\ \frac{1}{2} & u = 0 \text{ and } z \in \{6, 12\} \\ \frac{1}{2} & u \in \{2, 3, 4, 5\} \text{ and } \\ & z = u \text{ or } z = 6 + u \\ 0 & \text{otherwise} \end{cases} \quad (5.7.38)$$

$$\Pr(Z = z) = \frac{6 - |z - 7|}{36} \quad (5.7.39)$$

If  $Z$  and  $U$  are independent, then

$$\Pr(Z = z|U = u) = \frac{\Pr(Z = z, U = u)}{\Pr(U = u)} \quad (5.7.40)$$

$$= \frac{\Pr(Z = z) \Pr(U = u)}{\Pr(U = u)} \quad (5.7.41)$$

$$= \Pr(Z = z) \quad (5.7.42)$$

But,

$$\Pr(Z = z|U = u) \neq \Pr(Z = z) \quad (5.7.43)$$

$X$  and  $U$  are not independent from (5.7.43) and hence option (5.7c) is false.

d) Checking if  $Y$  and  $Z$  are independent

$$p_1 = \Pr(Z = z, Y = y) \quad (5.7.44)$$

$$= \Pr(X = z - y, Y = y) \quad (5.7.45)$$

$$= \Pr(X = z - y) \times \Pr(Y = y) \quad (5.7.46)$$

$$= \begin{cases} \frac{1}{36} & z - y \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases} \quad (5.7.47)$$

$$\Pr(Z = z) \times \Pr(Y = y) = \frac{6 - |z - 7|}{36} \times \frac{1}{6} \quad (5.7.48)$$

$$= \frac{6 - |z - 7|}{216} \quad (5.7.49)$$

$$\Pr(Z = z) \Pr(Y = y) \neq \Pr(Z = z, Y = y) \quad (5.7.50)$$

$X$  and  $Z$  are not independent from (5.7.50) and hence option (5.7d) is true.

Thus, options (5.7b) and (5.7d) are true.

5.8. Suppose  $X_1, X_2, X_3$  and  $X_4$  are independent and

identically distributed random variables, having density function  $f$ . Then,

- a)  $\Pr(X_4 > \text{Max}(X_1, X_2) > X_3) = \frac{1}{6}$
- b)  $\Pr(X_4 > \text{Max}(X_1, X_2) > X_3) = \frac{1}{8}$
- c)  $\Pr(X_4 > X_3 > \text{Max}(X_1, X_2)) = \frac{1}{12}$
- d)  $\Pr(X_4 > X_3 > \text{Max}(X_1, X_2)) = \frac{1}{6}$

**Solution:** The probability density function (pdf)  $f(x)$  of a random variable  $X$  is defined as the derivative of the cdf  $F(x)$ :

$$f(x) = \frac{d}{dx} F(x).$$

It is sometimes useful to consider the cdf  $F(x)$  in terms of the pdf  $f(x)$ :

$$F(x) = \int_{-\infty}^x f(t) dt$$

The PDF of  $X$  is,

$$F_X(x) = \int_{-\infty}^x f(x) dx \quad (5.8.1)$$

a)  $\Pr(X_2 > X_1)$

$$= \int_{-\infty}^{\infty} f_X(x) \int_{-\infty}^x f_X(t) dt dx \quad (5.8.2)$$

$$= \int_{-\infty}^{\infty} f_X(x) F_X(x) dx \quad (5.8.3)$$

$$= \frac{F_X^2(x)}{2} \Big|_{-\infty}^{\infty} \quad (5.8.4)$$

$$= \frac{1}{2}. \quad (5.8.5)$$

b)  $\Pr(X_4 > \text{Max}(X_1, X_2) > X_3)$

$$= \int_{-\infty}^{\infty} f_X(x) \int_{-\infty}^x f_X(t) \cdot {}^2C_1 \cdot \left[ \int_{-\infty}^t f_X(w) dw \right] \int_{-\infty}^t f_X(z) dz dt dx \quad (5.8.6)$$

$$= \int_{-\infty}^{\infty} f_X(x) \int_{-\infty}^x 2f_X(t) F_X^2(t) dt dx \quad (5.8.7)$$

$$= \int_{-\infty}^{\infty} f_X(x) \cdot \frac{2}{3} F_X^3(x) dx \quad (5.8.8)$$

$$= \frac{2}{3} \frac{F_X^4(x)}{4} \Big|_{-\infty}^{\infty} \quad (5.8.9)$$

$$= \frac{1}{6}. \quad (5.8.10)$$

c)  $\Pr(X_4 > X_3 > \text{Max}(X_1, X_2))$

$$= \int_{-\infty}^{\infty} f_X(x) \int_{-\infty}^x f_X(t) \int_{-\infty}^t f_X(z) \cdot {}^2C_1 \cdot \left[ \int_{-\infty}^t f_X(w) dw \right] dz dt dx \quad (5.8.11)$$

$$= \int_{-\infty}^{\infty} f_X(x) \int_{-\infty}^x f_X(t) \int_{-\infty}^t 2f_X(z) F_X(t) dz dt dx \quad (5.8.12)$$

$$= \int_{-\infty}^{\infty} f_X(x) \int_{-\infty}^x f_X(t) F_X^2(t) dt dx \quad (5.8.13)$$

$$= \int_{-\infty}^{\infty} f_X(x) \cdot \frac{1}{3} F_X^3(x) dx \quad (5.8.14)$$

$$= \frac{1}{3} \frac{F_X^4(x)}{4} \Big|_{-\infty}^{\infty} \quad (5.8.15)$$

$$= \frac{1}{12}. \quad (5.8.16)$$

∴ **Option 1,3 are correct answers.**

5.9. Consider a parallel system with two components. The lifetimes of the two components are independent and identically distributed random variables each following an exponential distribution with mean 1. The expected lifetime of

the system is:

A) 1

B)  $\frac{1}{2}$

C)  $\frac{3}{2}$

D) 2

**Solution:**

Consider two random variables X and Y which represent the lifetime of the two components namely A and B.

$$X \sim \text{Exp}(\lambda_X) \quad (5.9.1)$$

$$Y \sim \text{Exp}(\lambda_Y) \quad (5.9.2)$$

Let  $f_X(x)$  denote the probability distribution function for random variable X.

$$f_X(x) = \begin{cases} \lambda_X e^{-\lambda_X x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases} \quad (5.9.3)$$

Let  $f_Y(y)$  denote the probability distribution function for random variable Y.

$$f_Y(y) = \begin{cases} \lambda_Y e^{-\lambda_Y y} & y \geq 0 \\ 0 & \text{otherwise} \end{cases} \quad (5.9.4)$$

Let  $F_X(x)$  denote the cumulative distribution function for random variable X.

$$F_X(x) = \begin{cases} 1 - e^{-\lambda_X x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases} \quad (5.9.5)$$

Let  $F_Y(y)$  denote the cumulative distribution function for random variable Y.

$$F_Y(y) = \begin{cases} 1 - e^{-\lambda_Y y} & y \geq 0 \\ 0 & \text{otherwise} \end{cases} \quad (5.9.6)$$

$$E(X) = \frac{1}{\lambda_X} \quad (5.9.7)$$

$$E(Y) = \frac{1}{\lambda_Y} \quad (5.9.8)$$

From 5.9.7 and 5.9.8,

$$\lambda_X = \lambda_Y = 1 \quad (5.9.9)$$

Let Z be a random variable such that Z =




Fig. 5.9.1: Parallel system

$\max(X, Y)$

$$P(Z \leq z) = P(\max(X, Y) \leq z) \quad (5.9.10)$$

$$= P(X \leq z, Y \leq z) \quad (5.9.11)$$

$$= P(X \leq z)P(Y \leq z) \quad (5.9.12)$$

$$= (F_X(z) - F_X(0))(F_Y(z) - F_Y(0)) \quad (5.9.13)$$

$$= 1 - e^{-(\lambda_X)z} - e^{-(\lambda_Y)z} + e^{-(\lambda_X + \lambda_Y)z} \quad (5.9.14)$$

$P(Z \leq z)$  denotes the probability that the system dies in the first  $z$  seconds.

$$\text{Expectation} = \int_0^\infty z d(P(Z \leq z)) \quad (5.9.15)$$

$$= \int_0^\infty z(\lambda_X e^{-(\lambda_X)z} + \lambda_Y e^{-(\lambda_Y)z} - (\lambda_X + \lambda_Y) e^{-(\lambda_X + \lambda_Y)z}) dz \quad (5.9.16)$$

$$= \frac{1}{\lambda_X} + \frac{1}{\lambda_Y} - \frac{1}{\lambda_X + \lambda_Y} \quad (5.9.17)$$

From 5.9.9,

$$\text{Expectation} = \frac{3}{2} \quad (5.9.18)$$

Therefore, option C correct.

## 6 BINOMIAL DISTRIBUTION

6.1. The probability that a part manufactured by a company will be defective is 0.05. If 15 such parts are selected randomly and inspected, the probability that atleast two parts will be defec-

tive is ...

**Solution:** The desired probability is

$$\Pr(X \geq 2) = 1 - \Pr(X < 2) \quad (6.1.1)$$

$$= 1 - \Pr(X = 0) - \Pr(X = 1) \quad (6.1.2)$$

$$= 1 - {}^{15}C_0 p^0 q^{15} - {}^{15}C_1 p^1 q^{14} \quad (6.1.3)$$

$$= 0.1709 \quad (6.1.4)$$

where

$$p = 0.05, q = 1 - p = 0.95 \quad (6.1.5)$$

and  $X$  is binomial with parameters  $(15, p)$ .

6.2. Let  $X$  be a binomial random variable with parameters  $(11, \frac{1}{3})$ . At which value(s) of  $k$  is  $\Pr(X = k)$  maximized?

a)  $k = 2$

b)  $k = 3$

c)  $k = 4$

d)  $k = 5$

**Solution:**

The binomial distribution is given by:

$$\Pr(X = k) = {}^nC_k \times p^k \times q^{n-k} \quad (6.2.1)$$

We are given  $n = 11$ ,  $p = \frac{1}{3}$  and hence  $q = \frac{2}{3}$

$$\Pr(X = k) = {}^{11}C_k \left(\frac{2}{3}\right)^{11-k} \left(\frac{1}{3}\right)^k \quad (6.2.2)$$

$$\frac{\Pr(X = k)}{\Pr(X = k + 1)} \geq 1 \quad (6.2.3)$$

$$\Rightarrow \frac{{}^{11}C_k \left(\frac{2}{3}\right)^{11-k} \left(\frac{1}{3}\right)^k}{{}^{11}C_{k+1} \left(\frac{2}{3}\right)^{10-k} \left(\frac{1}{3}\right)^{k+1}} \geq 1 \quad (6.2.4)$$

$$\text{or, } k \geq 3 \quad (6.2.5)$$

after some algebra. Similarly,

$$\frac{\Pr(X = k)}{\Pr(X = k - 1)} \geq 1 \quad (6.2.6)$$

$$k \leq 4 \quad (6.2.7)$$

From (6.2.5) and (6.2.7), it is obvious that  $\Pr(X = k)$  is maximized for  $k = 3, k = 4$ . Hence, options 2 and 3 are correct. See Fig. 6.2.1 for a graphical verification.

6.3. The probability that a ticketless traveler is caught during a trip is 0.1. If the traveler




Fig. 6.2.1: Binomial Distribution of X

makes 4 trips, the probability that he/she will be caught during at least one of the trips is:

- a)  $1 - (0.9)^4$
- b)  $(1 - 0.9)^4$
- c)  $1 - (1 - 0.9)^4$
- d)  $(0.9)^4$

**Solution:** Let  $X_i \in \{0, 1\}$  represent the  $i$ th trip where 1 denotes a ticketless traveller is caught. Given,

$$\Pr(X_i = 1) = p = 0.1 \quad (6.3.1)$$

Let,

$$X = \sum_{i=1}^n X_i \quad (6.3.2)$$

where  $n$  is the number of trips and  $X$  has a binomial distribution.

$$p_X(k) = \begin{cases} {}^nC_k p^k (1-p)^{n-k}, & 0 \leq k \leq n \\ 0, & \text{otherwise} \end{cases} \quad (6.3.3)$$

As he/she makes 4 trips in total, Using (6.3.1) and (6.3.3),

$$\Pr(X = 0) = p_X(0) \quad (6.3.4)$$

$$= {}^4C_0 p^0 (1-p)^4 \quad (6.3.5)$$

$$\Pr(X = 0) = (0.9)^4 \quad (6.3.6)$$

Then probability of being caught in atleast one

trip is, (Using (6.3.6))

$$\Pr(X \geq 1) = 1 - \Pr(X < 1) \quad (6.3.7)$$

$$= 1 - \Pr(X = 0) \quad (6.3.8)$$

$$= 1 - (0.9)^4 \quad (6.3.9)$$

6.4. Let  $X$  be a binomial random variable with parameters  $\left(11, \frac{1}{3}\right)$ . At which value(s) of  $k$  is  $\Pr(X = k)$  maximized?

- a)  $k=2$
- b)  $k=3$
- c)  $k=4$
- d)  $k=5$

**Solution:**  $X$  has a binomial distribution :

$$\Pr(X = k) = {}^nC_k (q)^{n-k} (p)^k \quad (6.4.1)$$

Where,

- $n=11$
- $p = \frac{1}{3}$
- $q = 1 - p = 1 - \frac{1}{3} = \frac{2}{3}$

$$\Pr(X = k) = {}^{11}C_k \left(\frac{2}{3}\right)^{11-k} \left(\frac{1}{3}\right)^k \quad (6.4.2)$$

For  $\Pr(X = k)$  to be maximized

$$\Pr(X = k) \geq \Pr(X = k + 1) \quad (6.4.3)$$

$$\frac{\Pr(X = k)}{\Pr(X = k + 1)} = \frac{{}^{11}C_k \left(\frac{2}{3}\right)^{11-k} \left(\frac{1}{3}\right)^k}{{}^{11}C_{k+1} \left(\frac{2}{3}\right)^{10-k} \left(\frac{1}{3}\right)^{k+1}} \geq 1 \quad (6.4.4)$$

$$\frac{2(k+1)}{11-k} \geq 1 \quad (6.4.5)$$

$$\Rightarrow k \geq 3 \quad (6.4.6)$$

$$\Pr(X = k) \geq \Pr(X = k-1) \quad (6.4.7)$$

$$\frac{\Pr(X = k)}{\Pr(X = k-1)} = \frac{{}^{11}C_k \left(\frac{2}{3}\right)^{11-k} \left(\frac{1}{3}\right)^k}{{}^{11}C_{k-1} \left(\frac{2}{3}\right)^{12-k} \left(\frac{1}{3}\right)^{k-1}} \geq 1 \quad (6.4.8)$$

$$\frac{12-k}{2k} \geq 1 \quad (6.4.9)$$

$$\Rightarrow k \leq 4 \quad (6.4.10)$$

From (6.4.6) , (6.4.10) and since k is an integer

$\Pr(X = k)$  is maximized for  $k=3, k=4$

Thus options 2) and 3) are correct

## 7 POISSON DISTRIBUTION

7.1. Let  $X$  be a Poisson random variable with p.m.f

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k = 0, 1, 2, \dots; \lambda > 0 \\ 0 & \text{otherwise} \end{cases} \quad (7.1.1)$$

If  $Y = X^2 + 3$ , then what is  $P(Y = y)$  equal to?

- (A)  $\frac{e^{-\lambda} \lambda^{\sqrt{y-3}}}{\sqrt{(y-3)!}}$ , for  $y = \{3, 4, 7, 12, \dots\}$   
 (B)  $\frac{e^{-\lambda} \lambda^{\sqrt{y-3}}}{\sqrt{(3-y)!}}$ , for  $y = \{3, 4, 7, 12, \dots\}$   
 (C)  $\frac{e^{-\lambda} \lambda^{\sqrt{3-y}}}{\sqrt{(3-y)!}}$ , for  $y = \{4, 7, 12, \dots\}$   
 (D)  $\frac{e^{-\lambda} \lambda^{\sqrt{3-y}}}{\sqrt{(3-y)!}}$ , for  $y = \{4, 7, 12, \dots\}$

**Solution:**

$$Y = X^2 + 3 \quad (7.1.2)$$

$$\Rightarrow X = \sqrt{Y-3} \quad (7.1.3)$$

Substituting  $k = \sqrt{y-3}$  in (7.1.1),

$$p_Y(y) = \begin{cases} \frac{e^{-\lambda} \lambda^{\sqrt{y-3}}}{\sqrt{(y-3)!}}, & y = 3, 4, 7, 12, \dots \\ 0 & \text{otherwise} \end{cases} \quad (7.1.4)$$

Hence, the correct option is (A).

7.2. For  $n \geq 1$ , let  $X_n$  be a Poisson random variable with mean  $n^2$ . Which of the following are equal

$$\text{to } \frac{1}{\sqrt{2\pi}} \int_2^\infty e^{-x^2/2} dx$$

- a)  $\lim_{n \rightarrow \infty} \Pr(X_n > (n+1)^2)$   
 b)  $\lim_{n \rightarrow \infty} \Pr(X_n \leq (n+1)^2)$   
 c)  $\lim_{n \rightarrow \infty} \Pr(X_n < (n-1)^2)$   
 d)  $\lim_{n \rightarrow \infty} \Pr(X_n < (n-2)^2)$

**Solution:**

**Lemma 7.1.** Let  $Y_i$  be a Poisson random variable with mean 1 for  $i \in (1, n^2)$  Then,

$$\sum_{i=1}^{n^2} Y_i = X_n \quad (7.2.1)$$

**Lemma 7.2.**

$$\lim_{n \rightarrow \infty} \frac{X_n - n^2}{n} = \mathcal{N}(0, 1) \quad (7.2.2)$$

*Proof.* Using the central limit theorem,

$$\lim_{n \rightarrow \infty} \frac{Y_1 + Y_2 + \dots + Y_{n^2} - n^2}{n} = \mathcal{N}(0, 1) \quad (7.2.3)$$

yielding (7.2.2).  $\square$

**Lemma 7.3.**

$$Q(2) = \frac{1}{\sqrt{2\pi}} \int_2^\infty e^{-x^2/2} dx \quad (7.2.4)$$




Fig. 7.1.1: Poisson stem plot for  $X$  ( $\lambda = 5$ )

**Lemma 7.4.**

$$\Pr(X_n > k) \quad (7.2.5)$$

$$= Q\left(\frac{k - n^2}{n}\right) \quad (7.2.6)$$

*Proof.* For  $X \sim \mathcal{N}(0, 1)$ , using Lemma 7.2,

$$\Pr(X_n > k) = \Pr\left(X > \frac{k - n^2}{n}\right) \quad (7.2.7)$$

yielding (7.2.6).  $\square$

**Lemma 7.5.**

$$Q(X) = 1 - Q(-x) \quad (7.2.8)$$

a) Using Lemma 7.4,

$$\lim_{n \rightarrow \infty} \Pr(X_n > (n+1)^2) = \lim_{n \rightarrow \infty} Q\left(\frac{(n+1)^2 - n^2}{n}\right) \quad (7.2.9)$$

$$= Q(2) \quad (7.2.10)$$

$\therefore$  **Option 1 is correct**

b) Using Lemma 7.4,

$$\lim_{n \rightarrow \infty} \Pr(X_n \leq (n+1)^2) \quad (7.2.11)$$

$$= 1 - \lim_{n \rightarrow \infty} \Pr(X_n > (n+1)^2)$$

$$= 1 - \lim_{n \rightarrow \infty} Q\left(\frac{(n+1)^2 - n^2}{n}\right) \quad (7.2.12)$$

$$= 1 - Q(2) \quad (7.2.13)$$

$$\neq Q(2) \quad (7.2.14)$$

$\therefore$ , from Lemma 7.3 **Option 2 is incorrect**

c) Using Lemma 7.4 and Lemma 7.5

$$\begin{aligned} \lim_{n \rightarrow \infty} \Pr(X_n < (n-1)^2) \\ = 1 - \lim_{n \rightarrow \infty} \Pr(X_n \geq (n-1)^2) \end{aligned} \quad (7.2.15)$$

$$= 1 - \lim_{n \rightarrow \infty} Q\left(\frac{(n-1)^2 - n^2}{n}\right) \quad (7.2.16)$$

$$= 1 - Q(-2) \quad (7.2.17)$$

$$= Q(2) \quad (7.2.18)$$

$\therefore$  **Option 3 is also correct**

d) Using Lemma 7.4 and Lemma 7.5,

$$\begin{aligned} \lim_{n \rightarrow \infty} \Pr(X_n < (n-2)^2) \\ = 1 - \lim_{n \rightarrow \infty} \Pr(X_n \geq (n-2)^2) \end{aligned} \quad (7.2.19)$$

$$= 1 - \lim_{n \rightarrow \infty} Q\left(\frac{(n-2)^2 - n^2}{n}\right) \quad (7.2.20)$$

$$= 1 - Q(-4) \quad (7.2.21)$$

$$= Q(4) < Q(2) \quad (7.2.22)$$

$\therefore$  **Option 4 is incorrect**

7.3. Suppose customers arrive in a shop according to a Poisson process with rate 4 per hour. The shop opens at 10 : 00 am. If it is given that the second customer arrives at 10 : 40 am, what is the probability that no customer arrived before 10 : 30 am?

- a)  $\frac{1}{4}$
- b)  $e^{-2}$
- c)  $\frac{1}{2}$
- d)  $e^{\frac{1}{2}}$

**Solution:** We need to find




Fig. 7.1.2: Stem plot for Y (Simulated and Theoretical) ( $\lambda = 5$ )

Random Variable	Time at which people arrive
$X_p$	$p = 10 : 00 - 10 : 30$
$X_q$	$q = 10 : 30 - 10 : 40$
$X_r$	$r = 10 : 00 - 10 : 40$
$Y$	$10 : 40$

TABLE 7.3.1: Random Variables

$$\Pr(X_p = 0 | Y = 2) \quad (7.3.1)$$

In the world where the  $2^{nd}$  person arrives at

10 : 40 am the (7.3.1) becomes:

$$= \frac{\Pr(X_p = 0, X_q = 1)}{\Pr(X_r = 1)} \quad (7.3.2)$$

$$= \frac{\Pr(X_p = 0) \times \Pr(X_q = 1)}{\Pr(X_r = 1)} \quad (7.3.3)$$

The Poisson function distribution for time interval  $t$  and rate  $\lambda$  for a random variable  $X$ :

$$f_X(x; t) = \frac{(\lambda t)^x \exp(-\lambda t)}{x!}$$

For the time interval  $p$ :

$$\lambda = 4, t = 0.5, x = 0 \quad (7.3.4)$$

$$\Pr(X_p = 0) = f_X\left(0; \frac{1}{2}\right) \quad (7.3.5)$$

$$= e^{-2} \quad (7.3.6)$$

$$(7.3.7)$$

For the time interval  $q$ :

$$\lambda = 4, t = \frac{1}{6}, x = 1 \quad (7.3.8)$$

$$\Pr(X_q = 1) = f_X\left(1; \frac{1}{6}\right) \quad (7.3.9)$$

$$= \frac{2}{3} e^{-\frac{2}{3}} \quad (7.3.10)$$

For the time interval  $r$ :

$$\lambda = 4, t = \frac{2}{3}, x = 1 \quad (7.3.11)$$

$$\Pr(X_r = 1) = f_X\left(1; \frac{2}{3}\right) \quad (7.3.12)$$

$$= \frac{8}{3} e^{-\frac{8}{3}} \quad (7.3.13)$$

Substituting (7.3.6) (7.3.10) (7.3.13) in (7.3.3):

$$\Pr(X_p = 0 | Y = 2) = \frac{1}{4} \quad (7.3.14)$$

7.4. Men arrive in a queue according to a Poisson process with rate  $\lambda_1$  and women arrive in the same queue according to another Poisson process with rate  $\lambda_2$ . The arrivals of men and women are independent. The probability that the first person to arrive in the queue is a man is:

$$a) \frac{\lambda_1}{\lambda_1 + \lambda_2}$$

$$b) \frac{\lambda_2}{\lambda_1 + \lambda_2}$$

$$c) \frac{\lambda_1}{\lambda_2}$$

$$d) \frac{\lambda_2}{\lambda_1}$$

**Solution:** Let  $X$  and  $Y$  be Poisson random variables, with the values  $X$  takes being the number of men joining the queue in an arbitrary time  $t$ , and the values  $Y$  takes being the number of women joining the queue in an arbitrary time  $t$ .

$$\Pr(X = i) = \frac{\lambda_1^i \cdot e^{-\lambda_1}}{i!} \quad (7.4.1)$$

$$\Pr(Y = i) = \frac{\lambda_2^i \cdot e^{-\lambda_2}}{i!} \quad (7.4.2)$$

For 2 independent Poisson distributions with means  $\lambda_1$  and  $\lambda_2$ , the simultaneous distribution can be represented by:

$$\Pr(X + Y = i) = \frac{(\lambda_1 + \lambda_2)^i \cdot e^{-(\lambda_1 + \lambda_2)}}{i!} \quad (7.4.3)$$

Now we take conditional probability that if only one person entered the queue within a certain time  $t$ , then the probability of them being a man and not a woman is given by:

$$\Pr(X = 1 | (X + Y) = 1) = \frac{\Pr((X = 1) + (Y = 0))}{\Pr(X + Y = 1)} \quad (7.4.4)$$

$$(7.4.5)$$

Since  $X$  and  $Y$  are independent,

$$\Pr(X = 1 | (X + Y) = 1) = \frac{\Pr(X = 1) \cdot \Pr(Y = 0)}{\Pr(X + Y = 1)} \quad (7.4.6)$$

$$= \frac{\frac{\lambda_1^1 \cdot e^{-\lambda_1}}{1!} \cdot \frac{\lambda_2^0 \cdot e^{-\lambda_2}}{0!}}{\frac{(\lambda_1 + \lambda_2)^1 \cdot e^{-(\lambda_1 + \lambda_2)}}{1!}} \quad (7.4.7)$$

$$= \frac{\lambda_1}{\lambda_1 + \lambda_2} \quad (7.4.8)$$

The probability that the first person to arrive in the queue is a man is option A, i.e.  $\frac{\lambda_1}{\lambda_1 + \lambda_2}$

## 8 GAUSSIAN DISTRIBUTION

8.1. Let  $U$  and  $V$  be two independent zero mean Gaussian random variables of variances  $\frac{1}{4}$  and  $\frac{1}{9}$  respectively. The probability  $\Pr(3V \geq 2U)$  is



...

**Solution:** From the given information,

$$U = \mathcal{N}\left(0, \frac{1}{4}\right) V = \mathcal{N}\left(0, \frac{1}{9}\right) \quad (8.1.1)$$

Let  $Y = 3V - 2U$ . Then,

$$E(Y) = 3E(V) - 2E(U) = 0 \quad (8.1.2)$$

$$\text{var}(Y) = 3^2\text{var}(V) + 2^2\text{var}(U) = 2 \quad (8.1.3)$$

$$\therefore Y = \mathcal{N}(0, 2) \quad (8.1.4)$$

Thus,

$$\Pr(3V \geq 2U) = \Pr(3V - 2U \geq 0) \quad (8.1.5)$$

$$= \Pr(Y \geq 0) = \frac{1}{2} \quad (8.1.6)$$

$\therefore Y$  is symmetric about the origin.

8.2.  $(X, Y)$  follows bivariate normal distribution  $N_2(0, 0, 1, 1, \rho)$ ,  $-1 < \rho < 1$ . Then,

- $X + Y$  and  $X - Y$  are uncorrelated only if  $\rho = 0$
- $X + Y$  and  $X - Y$  are uncorrelated only if  $\rho < 0$
- $X + Y$  and  $X - Y$  are uncorrelated only if  $\rho > 0$
- $X + Y$  and  $X - Y$  are uncorrelated for all values of  $\rho$

**Solution:** Given that

$$\mathbf{M} = \begin{pmatrix} X \\ Y \end{pmatrix} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \quad (8.2.1)$$

where

$$\boldsymbol{\mu} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (8.2.2)$$

$$\boldsymbol{\Sigma} = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix} \quad (8.2.3)$$

Also,

$$X + Y = \mathbf{A}^T \mathbf{M} \quad (8.2.4)$$

$$X - Y = \mathbf{B}^T \mathbf{M} \quad (8.2.5)$$

where

$$\mathbf{A} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad (8.2.6)$$

Thus,

$$\text{Cov}(X + Y, X - Y) = \mathbf{A}^T \boldsymbol{\Sigma} \mathbf{B} \quad (8.2.7)$$

$$= 0 \quad (8.2.8)$$

$\therefore X + Y$  and  $X - Y$  are uncorrelated irrespective of value of  $\rho$  where  $\rho \in (-1, 1)$ .

8.3. Let  $U$  and  $V$  be two independent zero mean Gaussian random variables of variances  $\frac{1}{4}$  and  $\frac{1}{9}$  respectively. The probability  $\Pr(3V \geq 2U)$  is

...

**Solution:** From the given information,

$$U = \mathcal{N}\left(0, \frac{1}{4}\right) V = \mathcal{N}\left(0, \frac{1}{9}\right) \quad (8.3.1)$$

Let  $Y = 3V - 2U$ . Then,

$$E(Y) = 3E(V) - 2E(U) = 0 \quad (8.3.2)$$

$$\text{var}(Y) = 3^2\text{var}(V) + 2^2\text{var}(U) = 2 \quad (8.3.3)$$

$$\therefore Y = \mathcal{N}(0, 2) \quad (8.3.4)$$

Thus,

$$\Pr(3V \geq 2U) = \Pr(3V - 2U \geq 0) \quad (8.3.5)$$

$$= \Pr(Y \geq 0) = \frac{1}{2} \quad (8.3.6)$$

$\therefore Y$  is symmetric about the origin.

8.4. Suppose  $\begin{pmatrix} X \\ Y \end{pmatrix}$  is a random vector such that the marginal distribution of  $X$  and the marginal distribution of  $Y$  are the same and each is normally distributed with mean 0 and variance 1. Then, which of the following conditions imply independence of  $X$  and  $Y$ ?

- $\text{Cov}(X, Y) = 0$
- $aX + bY$  is normally distributed with mean 0 and variance  $a^2 + b^2$  for all real  $a$  and  $b$
- $\Pr(X \leq 0, Y \leq 0) = \frac{1}{4}$
- $E[e^{itX + isY}] = E[e^{itX}]E[e^{isY}]$  for all real  $s$  and  $t$

**Solution:**

An important property of dirac delta function that will be used at multiple occasions in this solution is

$$\int_{-\infty}^{\infty} f(x)\delta(x - a)dx = f(a) \quad (8.4.1)$$

Given  $X \sim N(0, 1)$ ,  $Y \sim N(0, 1)$

a)

$$\text{Cov}(X, Y) = 0 \quad (8.4.2)$$

$$E[XY] - E[X]E[Y] = 0 \quad (8.4.3)$$

$$E[XY] = 0 \quad (8.4.4)$$

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dx dy = 0 \quad (8.4.5)$$

This doesn't imply independence. Counter example given below

Lets consider a case where  $X$  and  $Y$  are dependent based on the following relation,  $Y$  being independent of  $K$

$$X = KY \quad (8.4.6)$$

PMF for  $K$  is given as

$$p_K(k) = \begin{cases} \frac{1}{2} & k = 1 \\ \frac{1}{2} & k = -1 \\ 0 & \text{otherwise} \end{cases} \quad (8.4.7)$$

A simulation is given below,  $Y$  is gaussian, then  $X$  also follows gaussian Theoretically




Fig. 8.4.1:  $X$  and  $Y$ , if  $Y$  is normal

it can be proved in the following manner, Since  $K$  and  $Y$  are independent

$$f_X(x) = \Pr(K = 1) f_Y(x) + \Pr(K = -1) f_Y(-x) \quad (8.4.8)$$

$$= \frac{1}{2} (f_Y(x) + f_Y(-x)) \quad (8.4.9)$$

$$= f_Y(x) \quad (8.4.10)$$

Therefore,  $X$  follows identical but not in-

dependent distribution as  $Y$ , An alternative proof is given below as a proof for marginal probability

Now consider that  $X$  is normally distributed, we will establish  $Y$  is also normally distributed. The joint probability distribution is therefore

$$\begin{aligned} f_{XY}(x, y) &= f_{X|Y}(x|y) f_Y(y) \\ &= f_X(x) \frac{1}{2} (\delta(x+y) + \delta(x-y)) \end{aligned} \quad (8.4.11)$$

The marginal probability distribution function for  $X$  is given as

$$\int_{-\infty}^{\infty} f_X(x) \frac{1}{2} (\delta(x+y) + \delta(x-y)) dy \quad (8.4.12)$$

Using (8.4.1), we get

$$\int_{-\infty}^{\infty} f_X(x) \frac{1}{2} (\delta(x+y) + \delta(x-y)) dy = f_X(x) \quad (8.4.13)$$

We know that  $X \sim N(0, 1)$ ,  $f_X(x)$  represents gaussian probability distribution function.

Further, using symmetry of (8.4.6), we can establish that marginal distribution of  $Y$  is gaussian. Here is a proof anyways

$$f_Y(y) = \int_{-\infty}^{\infty} f_X(x) \frac{1}{2} (\delta(x+y) + \delta(x-y)) dx \quad (8.4.14)$$

Using (8.4.1), we get

$$f_Y(y) = \frac{1}{2} (f_X(y) + f_X(-y)) = f_X(y) \quad (8.4.15)$$

Since  $Y$  has identical probability distribution function,  $Y \sim N(0, 1)$

The covariance is given as

$$\text{Cov}(X, Y) = E[XY] - E[X]E[Y] = E[XY] \quad (8.4.16)$$

$$E[XY] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dy dx \quad (8.4.17)$$

$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_X(x) \frac{1}{2} (\delta(x+y) + \delta(x-y)) dy dx \quad (8.4.18)$$

$$= \int_{-\infty}^{\infty} x f_X(x) \int_{-\infty}^{\infty} y \frac{1}{2} (\delta(x+y) + \delta(x-y)) dy dx \quad (8.4.19)$$

Using (8.4.1)

$$E[XY] = \int_{-\infty}^{\infty} x f_X(x) \frac{1}{2} (x - x) dx = 0 \quad (8.4.20)$$

b) Defining the following matrices/vectors  
Given

vector/matrix	expression
$\mathbf{Z}$	$\begin{pmatrix} X & Y \end{pmatrix}^\top$
$\mathbf{C}$	$\begin{pmatrix} a & b \end{pmatrix}^\top$
$\boldsymbol{\mu}$	$\begin{pmatrix} 0 & 0 \end{pmatrix}^\top$
$\boldsymbol{\Sigma}$	$\begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$

TABLE 8.4.1: vectors/matrices and their expressions

$$\mathbf{C}^\top \mathbf{Z} \sim N(0, a^2 + b^2) \quad (8.4.21)$$

Since this is true for all  $a$  and  $b$ , it is equivalent to  $X$  and  $Y$  being jointly gaussian

$$\mathbf{Z} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \quad (8.4.22)$$

For correlated random variables  $X$  and  $Y$  in bivariate normal distribution, we have

$$\sigma_Z^2 = \sum_{i,j} \Sigma_{ij} \quad (8.4.23)$$

$$a^2 + b^2 = a^2 + b^2 + 2\rho ab \quad (8.4.24)$$

$$\therefore \rho = 0 \quad (8.4.25)$$

The joint distribution is given as

$$f_{\mathbf{Z}}(x, y) = \frac{\exp\left(-\frac{1}{2} (\mathbf{z} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{z} - \boldsymbol{\mu})\right)}{\sqrt{(2\pi)^2 |\boldsymbol{\Sigma}|}} \quad (8.4.26)$$

$$f_{\mathbf{Z}}(x, y) = \frac{\exp\left(-\frac{1}{2} \begin{pmatrix} x & y \end{pmatrix} I_2 \begin{pmatrix} x & y \end{pmatrix}^\top\right)}{\sqrt{(2\pi)^2}} \quad (8.4.27)$$

Where  $I_2$  is the identity matrix of order 2

$$f_{\mathbf{Z}}(x, y) = \frac{\exp\left(-\frac{1}{2} \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} x & y \end{pmatrix}^\top\right)}{\sqrt{(2\pi)^2}} \quad (8.4.28)$$

$$f_{\mathbf{Z}}(x, y) = \frac{\exp\left(-\frac{1}{2} (x^2 + y^2)\right)}{\sqrt{(2\pi)^2}} = f_X(x) f_Y(y) \quad (8.4.29)$$

$\therefore$  **Option(2) is correct**, A simulation for bivariate gaussian is given below




Fig. 8.4.2: bivariate gaussian while 0 mean vector and identity covariance matrix

c)

$$\Pr(X \leq 0, Y \leq 0) = \frac{1}{4} \quad (8.4.30)$$

This doesn't imply independence, it can be true even for dependent  $X$  and  $Y$ , the counter example is (8.4.11), the joint probability function is symmetric across all 4 quadrants

$$\therefore \Pr(X \leq 0, Y \leq 0) = \frac{1}{4} \quad (8.4.31)$$

Alternatively, here is proof

$$\Pr(X \leq 0) = F_X(0) = \frac{1}{2} \quad (8.4.32)$$

Using (8.4.6)

$$\Pr(Y \leq 0 | X \leq 0) = \frac{1}{2} \quad (8.4.33)$$

Using (8.4.32) and (8.4.33)

$$\Pr(X \leq 0, Y \leq 0) = \frac{1}{4} \quad (8.4.34)$$

d)

$$E[e^{itX+isY}] = E[e^{itX}]E[e^{isY}] \quad (8.4.35)$$

$$E[e^{itX+isY}] = \varphi_X(t)\varphi_Y(s) \quad (8.4.36)$$

The inverse is given as

$$f_{XY}(x, y) = \frac{1}{4\pi^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-itX-isY} E[e^{itX+isY}] ds dt \quad (8.4.37)$$

Using (8.4.36)

$$f_{XY}(x, y) = \frac{1}{4\pi^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-itX-isY} \varphi_X(t)\varphi_Y(s) ds dt \quad (8.4.38)$$

$$f_{XY}(x, y) = f_X(x)f_Y(y) \quad (8.4.39)$$

$\therefore$  **Option(4) is correct**

## 9 GEOMETRIC DISTRIBUTION

9.1. Suppose  $X$  has density

$$f(x|\theta) = \frac{1}{\theta} e^{-x/\theta}, x > 0 \quad (9.1.1)$$

Define

$$Y = k, \quad k \leq X < k+1, \quad k = 0, 1, 2, \dots \quad (9.1.2)$$

Then the distribution of  $Y$  is

- a) Normal
- b) Binomial
- c) Poisson
- d) Geometric

**Solution:**

$$\Pr(Y = k) = \Pr(k \leq X < k+1) \quad (9.1.3)$$

$$= \int_k^{k+1} f(x|\theta) dx \quad (9.1.4)$$

$$= \int_k^{k+1} \frac{1}{\theta} e^{-\frac{x}{\theta}} dx \quad (9.1.5)$$

$$= \left[ -e^{-\frac{x}{\theta}} \right]_k^{k+1} \quad (9.1.6)$$

$$= e^{-\frac{k}{\theta}} \left( 1 - e^{-\frac{1}{\theta}} \right) \quad (9.1.7)$$

$$\Rightarrow \Pr(Y = k) = (1-p)^k p, k = 0, 1, 2, \dots \quad (9.1.8)$$

where

$$p = 1 - e^{-\frac{1}{\theta}} \quad (9.1.9)$$

Therefore, the distribution of  $Y$  is 4) Geometric.

## 10 TWO DIMENSIONS

10.1. Let  $c \in \mathbb{R}$  be a constant. Let  $X, Y$  be random variables with joint probability density function

$$f(x, y) = \begin{cases} cxy & 0 < x < y < 1, \\ 0 & \text{otherwise} \end{cases} \quad (10.1.1)$$

Which of the following statements are correct ?

- a)  $c = \frac{1}{8}$
- b)  $c = 8$
- c)  $X$  and  $Y$  are independent
- d)  $\Pr(X = Y) = 0$

**Solution:**

- a) False
- b) By definition,

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx \quad (10.1.2)$$

$$= \int_0^y cxy dx \quad (10.1.3)$$

$$= cy \left( \frac{x^2}{2} \right) \Big|_0^y \quad (10.1.4)$$

$$= \frac{cy^3}{2} \quad (10.1.5)$$

$$\Rightarrow f_Y(y) = \begin{cases} \frac{cy^3}{2}, & 0 < y < 1 \\ 0 & \text{otherwise.} \end{cases} \quad (10.1.6)$$

$\therefore$  the area under the pdf is 1, from (10.1.6),

$$\Rightarrow \int_{-\infty}^{\infty} f_Y(y) dy = 1 \quad (10.1.7)$$

$$\Rightarrow \int_0^1 c \frac{y^3}{2} = 1 \quad (10.1.8)$$

$$\Rightarrow \frac{c}{8} = 1 \quad (10.1.9)$$

$$\text{or, } c = 8 \quad (10.1.10)$$

Also,

$$f_Y(y) = \begin{cases} 4y^3, & \text{if } 0 < y < 1 \\ 0, & \text{otherwise} \end{cases} \quad (10.1.11)$$

c)

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy \quad (10.1.12)$$

$$= \int_x^1 cxy dy \quad (10.1.13)$$

$$= cx \left( \frac{y^2}{2} \right) \Big|_x^1 \quad (10.1.14)$$

$$= cx \left( \frac{1 - x^2}{2} \right) \quad (10.1.15)$$

$$\Rightarrow f_X(x) = \begin{cases} 4x(1 - x^2), & 0 < x < 1 \\ 0, & \text{otherwise} \end{cases} \quad (10.1.16)$$

From (10.1.16) and (10.1.11)

$$f_X(x) \times f_Y(y) = \begin{cases} 16xy^3(1 - x^2), & \text{if } 0 < x, y < 1 \\ 0, & \text{otherwise} \end{cases} \quad (10.1.17)$$

$$\neq f(x, y) \quad (10.1.18)$$

Hence,  $X$  and  $Y$  are not independent.

d)

$$F_X(x) = \int_{-\infty}^x f_X(x) dx \quad (10.1.19)$$

$$= \int_0^x 4x(1 - x^2) dx \quad (10.1.20)$$

$$= \int_0^x 4x - 4x^3 dx \quad (10.1.21)$$

$$= 2x^2 - 4x^4 \text{ for } 0 < x < 1 \quad (10.1.22)$$

yielding

$$F_X(x) = \begin{cases} 0 & x \leq 0 \\ 2x^2 - 4x^4 & 0 < x < 1 \\ 1 & x \geq 1 \end{cases} \quad (10.1.23)$$

From (10.1.23),

$$\begin{aligned} \Pr(Y - \epsilon < X < Y + \epsilon) \\ &= F_X(Y + \epsilon) - F_X(Y - \epsilon) \\ &= 8\epsilon Y(1 - Y^2 - \epsilon^2) \end{aligned} \quad (10.1.24)$$

upon simplification. Letting

$$g(Y) = 8\epsilon Y(1 - Y^2 - \epsilon^2), \quad (10.1.25)$$

$$E[g(Y)] = \int_{-\infty}^{\infty} g(y) f_Y(y) dy \quad (10.1.26)$$

$$= \int_0^1 (4y^3)(8\epsilon y)(1 - y^2 - \epsilon^2) dy \quad (10.1.27)$$

$$\begin{aligned} \Rightarrow \Pr(Y - \epsilon < X < Y + \epsilon) \\ &= 32\epsilon \left( \frac{2 - 7\epsilon^2}{35} \right) \end{aligned} \quad (10.1.28)$$

Now substituting  $\epsilon = 0$  in the above,

$$\Pr(X = Y) = 0 \quad (10.1.29)$$

10.2. Let  $X$  and  $Y$  be random variables having the joining probability density function

$$f_{XY}(x, y) = \begin{cases} \frac{1}{\sqrt{2\pi y}} e^{\frac{-1}{2y}(x-y)^2} & x \in (-\infty, \infty), \\ & y \in (0, 1) \\ 0 & \text{otherwise} \end{cases} \quad (10.2.1)$$

The covariance between the random variables  $X$  and  $Y$  is

**Solution:**

10.3. Let a random variable  $X$  follow exponential distribution with mean 2. Define  $Y = [X - 2 | X > 2]$ . The value of  $\Pr(Y \geq t)$  is ...

**Solution:** From the given information,

$$\Pr(Y \geq t) = \frac{\Pr(X - 2 \geq t, X > 2)}{\Pr(X > 2)} \quad (10.3.1)$$

$$= \frac{\Pr(X \geq t + 2, X > 2)}{\Pr(X > 2)} \quad (10.3.2)$$

$\therefore X$  has an exponential distribution with pa-

parameter  $\lambda = \frac{1}{2}$ ,

$$F_X(x) = \begin{cases} 1 - e^{-\lambda x}, & \text{if } 0 < x < \infty \\ 0, & \text{otherwise} \end{cases} \quad (10.3.3)$$

and

$$\Pr(X > 2) = 1 - F_X(2) = e^{-2\lambda} \quad (10.3.4)$$

Also,

$$\Pr(X \geq t + 2, X > 2) = \begin{cases} \Pr(X \geq t + 2) & t \geq 0 \\ \Pr(X > 2) & t < 0 \end{cases} \quad (10.3.5)$$

Substituting (10.3.5) in (10.3.2), using (10.3.4) and simplifying,

$$\Pr(Y \geq t) = \begin{cases} e^{-\frac{t}{2}} & t \geq 0 \\ 1 & t < 0 \end{cases} \quad (10.3.6)$$

10.4. Let X and Y be two random variables with joint probability density function

$$f(x, y) = \begin{cases} \frac{1}{\pi} & 0 \leq x^2 + y^2 \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Which of the following statements are correct?

a) X and Y are independent.

b)  $\Pr(X > 0) = \frac{1}{2}$

c)  $E(Y) = 0$

d)  $\text{Cov}(X, Y) = 0$

**Solution:**

a) The marginal PDF of X is given by

$$f_X(x) = \int_{y=-\infty}^{y=\infty} f_{XY}(x, y) dy \quad (10.4.1)$$

$$= \int_{y=-\sqrt{1-x^2}}^{y=\sqrt{1-x^2}} \frac{1}{\pi} dy \quad (10.4.2)$$

$$= \frac{2\sqrt{1-x^2}}{\pi} \quad (10.4.3)$$

The marginal PDF of Y is given by

$$f_Y(y) = \int_{x=-\infty}^{x=\infty} f_{XY}(x, y) dx \quad (10.4.4)$$

$$= \int_{x=-\sqrt{1-y^2}}^{x=\sqrt{1-y^2}} \frac{1}{\pi} dx \quad (10.4.5)$$

$$= \frac{2\sqrt{1-y^2}}{\pi} \quad (10.4.6)$$

Now,

$$f_X(x) \times f_Y(y) = \frac{2\sqrt{1-x^2}}{\pi} \times \frac{2\sqrt{1-y^2}}{\pi} \quad (10.4.7)$$

$$= \frac{4(1-x^2)(1-y^2)}{\pi^2} \quad (10.4.8)$$

$$\neq \frac{1}{\pi} \quad (10.4.9)$$

$$\neq f_{XY}(x, y) \quad (10.4.10)$$

Therefore, X and Y are not independent.

b) Now,

$$\Pr(X > 0) = \int_{x=0}^{x=\infty} f_X(x) dx \quad (10.4.11)$$

$$= \int_{x=0}^{x=1} \frac{2\sqrt{1-x^2}}{\pi} dx \quad (10.4.12)$$

$$= \left( \frac{\arcsin(x) + x\sqrt{1-x^2}}{\pi} \right) \Big|_0^1 \quad (10.4.13)$$

$$= \frac{1}{2} \quad (10.4.14)$$

Therefore, option(2) is correct.

c) Now,

$$E[Y] = \int_{y=-\infty}^{y=\infty} y f_Y(y) dy \quad (10.4.15)$$

$$= \int_{y=-1}^{y=1} \frac{2y\sqrt{1-y^2}}{\pi} dy \quad (10.4.16)$$

$$= \left( \frac{-2(1-y^2)^{\frac{3}{2}}}{3\pi} \right)_{-1}^1 \quad (10.4.17)$$

$$= 0 \quad (10.4.18)$$

Therefore, option(3) is also correct.

d) Now,

$$E[XY] = \int_x \int_y xy f_{XY}(x, y) dy dx \quad (10.4.19)$$

$$= \int_{x=-1}^{x=1} \int_{y=-\sqrt{1-x^2}}^{y=\sqrt{1-x^2}} \frac{xy}{\pi} dy dx \quad (10.4.20)$$

$$= \frac{x}{\pi} \int_{x=-1}^{x=1} \left( \frac{y^2}{2} \right)_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} dx \quad (10.4.21)$$

$$= 0 \quad (10.4.22)$$

Now,

$$Cov(X, Y) = E[XY] - E[X]E[Y] \quad (10.4.23)$$

$$= 0 - E[X] \times 0 \quad (10.4.24)$$

$$= 0 \quad (10.4.25)$$

Therefore, option(4) is also correct.

10.5. Let  $X$  and  $Y$  be two random variables satisfying  $X \geq 0, Y \geq 0, E(X) = 3, Var(X) = 9, E(Y) = 2$  and  $Var(Y) = 4$ . Which of the following statements are correct?

A)  $0 \leq Cov(X, Y) \leq 4$

B)  $E(XY) \leq 3$

C)  $Var(X + Y) \leq 25$

D)  $E(X + Y)^2 \geq 25$

**Solution:**

$$E(X^2) = Var(X) + (E(X))^2 = 18 \quad (10.5.1)$$

Similarly,

$$E(Y^2) = Var(Y) + (E(Y))^2 = 8 \quad (10.5.2)$$

We can use the Covariance inequality for this question,

$$(Cov(X, Y))^2 \leq Var(X)Var(Y) \quad (10.5.3)$$

The proof of this inequality is as shown,

$$Var\left(\frac{X}{\sigma_X} \pm \frac{Y}{\sigma_Y}\right) = Var\left(\frac{X}{\sigma_X}\right) + Var\left(\frac{\pm Y}{\sigma_Y}\right) + 2Cov\left(\frac{X}{\sigma_X}, \frac{\pm Y}{\sigma_Y}\right) \quad (10.5.4)$$

$$= \frac{1}{\sigma_X^2} Var(X) + \frac{1}{\sigma_Y^2} Var(Y)$$

$$+ 2Cov\left(\frac{X}{\sigma_X}, \frac{\pm Y}{\sigma_Y}\right) \quad (10.5.5)$$

$$= 2 \pm 2 \frac{Cov(X, Y)}{\sigma_X \sigma_Y} \quad (10.5.6)$$

Since Variance is always positive,

$$Var\left(\frac{X}{\sigma_X} \pm \frac{Y}{\sigma_Y}\right) \geq 0 \quad (10.5.7)$$

$$2 \pm 2 \frac{Cov(X, Y)}{\sigma_X \sigma_Y} \geq 0 \quad (10.5.8)$$

$$1 \pm 1 \frac{Cov(X, Y)}{\sigma_X \sigma_Y} \geq 0 \quad (10.5.9)$$

$$|(Cov(X, Y))| \leq (\sigma_X)(\sigma_Y) \quad (10.5.10)$$

$$(Cov(X, Y))^2 \leq Var(X)Var(Y) \quad (10.5.11)$$

a) Substituting values of variance we get,

$$-6 \leq Cov(X, Y) \leq 6 \quad (10.5.12)$$

**Therefore, option A is incorrect.**

b) From equation (10.5.12),

$$Cov(X, Y) = E(XY) - E(X)E(Y) \quad (10.5.13)$$

$$-6 \leq E(XY) - E(X)E(Y) \leq 6 \quad (10.5.14)$$

$$0 \leq E(XY) \leq 12 \quad (10.5.15)$$

Also, if  $X$  and  $Y$  are independent,

$$E(XY) = E(X)E(Y) = 6 \quad (10.5.16)$$

**Therefore, Option B is incorrect.**

c) Now,

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y) \quad (10.5.17)$$

$$= 13 + 2\text{Cov}(X, Y) \quad (10.5.18)$$

From equation (10.5.12),

$$1 \leq \text{Var}(X + Y) \leq 25 \quad (10.5.19)$$

**Therefore, Option C is correct.**

d) Now,

$$E(X + Y)^2 = E(X^2) + E(Y^2) + 2E(XY) \quad (10.5.20)$$

$$E(X + Y)^2 = 26 + 2E(XY) \quad (10.5.21)$$

From equation (10.5.15),

$$26 \leq E(X + Y)^2 \leq 50 \quad (10.5.22)$$

**Therefore, Option D is correct.**

10.6. The joint probability density function of (X,Y) is

$$f(x, y) = \begin{cases} 6(1 - x) & \text{if } 0 < y < x, 0 < x < 1 \\ 0 & \text{otherwise} \end{cases} \quad (10.6.1)$$

Which among the following are correct?

a) X and Y are not independent

$$b) f_Y(y) = \begin{cases} 3(y - 1)^2 & \text{if } 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

c) X and Y are independent

$$d) f_Y(y) = \begin{cases} 3\left(y - \frac{1}{2}y^2\right) & \text{if } 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

**Solution:** Given joint probability density function of X and Y, marginal probability density functions are as follows:

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy \quad (10.6.2)$$

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx \quad (10.6.3)$$

Calculating  $f_X(x)$

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy \quad (10.6.4)$$

$$= \int_0^x 6(1 - x) dy \quad (10.6.5)$$

$$f_X(x) = \begin{cases} 6x(1 - x) & 0 < x < 1 \\ 0 & \text{otherwise} \end{cases} \quad (10.6.6)$$

Calculating  $f_Y(y)$

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx \quad (10.6.7)$$

$$= \int_y^1 6(1 - x) dx \quad (10.6.8)$$

$$= 6x - 3x^2 \Big|_y^1 \quad (10.6.9)$$

$$= 3 - 6y + 3y^2 \quad (10.6.10)$$

$$= 3(y - 1)^2 \quad (10.6.11)$$

$$f_Y(y) = \begin{cases} 3(y - 1)^2 & 0 < y < 1 \\ 0 & \text{otherwise} \end{cases} \quad (10.6.12)$$

To check whether X and Y are independent, we calculate  $f_X(x) \times f_Y(y)$ . From (10.6.6) and (10.6.12)

$$f_X(x) \times f_Y(y) = \begin{cases} 18x(1 - x)(y - 1)^2 & 0 < x < 1, 0 < y < 1 \\ 0 & \text{otherwise} \end{cases} \quad (10.6.13)$$

$$\neq f(x, y) \quad (10.6.14)$$

Since  $f(x, y)$  and  $f_X(x) \times f_Y(y)$  are different, random variables X and Y are not independent.

Options 1 and 2 are correct

10.7. Suppose that (X, Y) has a joint probability distribution with the marginal distribution of X being  $N(0, 1)$  and  $E(Y|X = x) = x^3$  for all  $x \in R$ . Then, which of the following statements are true?

a)  $\text{Corr}(X, Y) = 0$

b)  $\text{Corr}(X, Y) > 0$

c)  $\text{Corr}(X, Y) < 0$

d) X and Y are independent

**Solution:** The following result shall be useful



later. For  $n \in N$

$$\int_{-\infty}^{\infty} \frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx = \begin{cases} 0 & n \text{ is odd} \\ (n-1) \times \dots \times 3 \times 1 & n \text{ is even} \end{cases} \quad (10.7.1)$$

The proof for the above can be found at the end of the solution.

$$\text{Corr}(X, Y) = \frac{\sigma_{XY}^2}{\sigma_X \sigma_Y} \quad (10.7.2)$$

We know  $X \sim N(0, 1)$ . Thus,

$$f_X(x) = \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} \quad (10.7.3)$$

$$E(X) = 0 \quad (10.7.4)$$

$$\sigma_X^2 = 1 \quad (10.7.5)$$

$$\sigma_Y^2 = E(Y^2) - E(Y)^2 \quad (10.7.6)$$

$$E(Y) = \int_{-\infty}^{\infty} E(Y|X = x) f_X(x) dx \quad (10.7.7)$$

$$= \int_{-\infty}^{\infty} \frac{x^3 e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (10.7.8)$$

$$= 0 \quad (10.7.9)$$

$$E(Y^2) = \int_{-\infty}^{\infty} E(Y^2|X = x) f_X(x) dx \quad (10.7.10)$$

$$= \int_{-\infty}^{\infty} \frac{x^6 e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (10.7.11)$$

$$= 15 \quad (10.7.12)$$

Substituting in (10.7.6)

$$\sigma_Y^2 = 15 \quad (10.7.13)$$

$$\sigma_{XY}^2 = E(XY) - E(X)E(Y) \quad (10.7.14)$$

$$E(XY) = \int_{-\infty}^{\infty} E(XY|X = x) f_X(x) dx \quad (10.7.15)$$

$$= \int_{-\infty}^{\infty} E(xY|X = x) f_X(x) dx \quad (10.7.16)$$

$$= \int_{-\infty}^{\infty} x E(Y|X = x) f_X(x) dx \quad (10.7.17)$$

$$= \int_{-\infty}^{\infty} \frac{x^4 e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (10.7.18)$$

$$= 3 \quad (10.7.19)$$

Substituting in (10.7.14)

$$\sigma_{XY}^2 = 3 \quad (10.7.20)$$

Substituting in (10.7.2)

$$\text{Corr}(X, Y) = \frac{3}{\sqrt{15}} > 0 \quad (10.7.21)$$

Since  $\text{Corr}(X, Y) \neq 0$ , X and Y are dependent. Thus option 2 is the only correct option. **Proof**

**for the integral:** If n is odd,  $\frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}}$  is an odd function, thus

$$\int_{-\infty}^{\infty} \frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx = 0 \quad (10.7.22)$$

If n is even,

$$\int_{-\infty}^{\infty} \frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx = \int_{-\infty}^{\infty} (x^{n-1}) \left( \frac{x e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} \right) dx \quad (10.7.23)$$

Using integration by parts,

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx &= \left( x^{n-1} \int \frac{x e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \right) \Big|_{-\infty}^{\infty} \\ &- (n-1) \int_{-\infty}^{\infty} x^{n-2} \left( \int \frac{x e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \right) dx \end{aligned} \quad (10.7.24)$$

$$= \left( x^{n-1} \left( -\frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} \right) \right) \Big|_{-\infty}^{\infty} - (n-1) \int_{-\infty}^{\infty} x^{n-2} \left( -\frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} \right) dx \quad (10.7.25)$$

c)  $\frac{7}{9}$   
d)  $\frac{1}{3}$

**Solution:** Let  $U \in \{1, 2, 3\}$  and  $V \in \{1, 2, 3\}$

$$= (n-1) \int_{-\infty}^{\infty} \frac{x^{n-2} e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (10.7.26)$$

TABLE 10.8.1: Probability of selecting values for  $U$

$$= (n-1)(n-3) \int_{-\infty}^{\infty} \frac{x^{n-4} e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (10.7.27)$$

$k$	1	2	3
$\Pr(U = k)$	1/3	1/3	1/3

$$= (n-1) \times \dots \times 3 \times 1 \int_{-\infty}^{\infty} \frac{x^0 e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (10.7.28)$$

TABLE 10.8.2: Probability of selecting values for  $V$

$$= (n-1) \times \dots \times 3 \times 1 \quad (10.7.29)$$

$k$	1	2	3
$\Pr(V = k)$	1/3	1/3	1/3

Alternative proof for the integral:

If  $n$  is odd,  $\frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}}$  is an odd function, thus

$$\int_{-\infty}^{\infty} \frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx = 0 \quad (10.7.30)$$

If  $n$  is even, let  $n = 2k$ . We differentiate the following identity  $k$  times w.r.t.  $\alpha$ .

$$\int_{-\infty}^{\infty} e^{-\alpha x^2} dx = \sqrt{\left(\frac{\pi}{\alpha}\right)} \quad (10.7.31)$$

On differentiating  $k$  times, we get

$$\int_{-\infty}^{\infty} x^{2k} e^{-\alpha x^2} dx = \frac{1 \times 3 \times \dots \times (2k-1)}{2^k} \sqrt{\left(\frac{\pi}{\alpha^{2k+1}}\right)} \quad (10.7.32)$$

On substituting  $\alpha = \frac{1}{2}$ , we get

$$\int_{-\infty}^{\infty} x^n e^{-\frac{x^2}{2}} dx = 1 \times 3 \times \dots \times (n-1) \sqrt{2\pi} \quad (10.7.33)$$

Thus

$$\int_{-\infty}^{\infty} \frac{x^n e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx = (n-1) \times \dots \times 3 \times 1 \quad (10.7.34)$$

10.8. Consider the quadratic equation  $x^2 + 2Ux + V = 0$  where  $U$  and  $V$  are chosen independently and randomly from  $\{1, 2, 3\}$  with equal probabilities. Then probability that the equation has both roots real

- a)  $\frac{2}{3}$   
b)  $\frac{1}{2}$

For  $x^2 + 2Ux + V = 0$  to have real roots,

$$b^2 - 4ac \geq 0 \quad (10.8.1)$$

$$(2U)^2 - 4(1)(V) \geq 0 \quad (10.8.2)$$

$$U^2 \geq V \quad (10.8.3)$$

$$\Pr(U^2 \geq V) = 1 - \Pr(U^2 < V) \quad (10.8.4)$$

The possible pairs  $(U, V)$  for  $\Pr(U^2 < V)$ ,

TABLE 10.8.3: Table for  $\Pr(U^2 < V)$

$(U, V)$ for $U^2 < V$	Probability
(1, 2)	$\Pr(U = 1)\Pr(V = 2) = 1/9$
(1, 3)	$\Pr(U = 1)\Pr(V = 3) = \frac{1}{9}$
Total	$\Pr(U^2 < V) = \frac{2}{9}$

$$\Pr(U^2 \geq V) = 1 - \frac{2}{9} \quad (10.8.5)$$

$$\Pr(U^2 \geq V) = \frac{7}{9} \quad (10.8.6)$$

Hence, Option 3 is correct.

## 11 INTEGRAL TRANSFORMS

11.1.  $X$  and  $Y$  are independent random variables each having the density

$$f(t) = \frac{1}{\pi} \frac{1}{1+t^2} \quad -\infty < t < +\infty \quad (11.1.1)$$

Then the density function of  $\frac{X+Y}{3}$  for  $-\infty < t < +\infty$  is

a)  $\frac{6}{\pi} \frac{1}{4+9t^2}$

b)  $\frac{6}{\pi} \frac{1}{9+4t^2}$

c)  $\frac{3}{\pi} \frac{1}{1+9t^2}$

d)  $\frac{3}{\pi} \frac{1}{9+t^2}$

**Solution:** Let us consider the random variables X and Y. The Characteristic function of the probability density  $f(t)$  is

$$g(w) = \int_{-\infty}^{\infty} f(t)e^{iwt} dt \quad (11.1.2)$$

$$= \int_{-\infty}^{\infty} \frac{1}{\pi} \frac{1}{1+t^2} e^{iwt} dt \quad (11.1.3)$$

$$= e^{-|w|}, -\infty < w < \infty \quad (11.1.4)$$

The product of the Characteristic function of probability density of X and Y is

$$h(w) = g_1(w) \times g_2(w) = e^{-2|w|} \quad (11.1.5)$$

To get the probability density of X+Y, we find the inverse characteristic function of h(w). But since there is a one to one correspondence between a function and its fourier transform and  $h(w) = g(2w)$

$$F_{X+Y}(t) = \frac{1}{2} f\left(\frac{t}{2}\right) \quad (11.1.6)$$

$$= \frac{1}{2\pi} \frac{4}{4+t^2}, -\infty < t < \infty \quad (11.1.7)$$

We know that if a random variable M has a probability density  $f_M(x)$ , then the probability density of random variable kM is

$$f_{kM}(x) = \frac{1}{|k|} f_M\left(\frac{x}{|k|}\right) \quad (11.1.8)$$

Probability density of  $Z = \frac{X+Y}{3}$  given  $F_{X+Y}(t)$

is

$$F_Z(t) = 3 \times f_{X+Y}(3t) \quad (11.1.9)$$

$$= \frac{6}{\pi} \frac{1}{4+9t^2} \quad (11.1.10)$$

11.2. Suppose X and Y are independent and identically distributed random variables and let  $Z = X + Y$ . Then the distribution of Z is in the same family as that of X and Y if X is **Solution:**

- |            |                |
|------------|----------------|
| 1) Normal  | 2) Exponential |
| 3) Uniform | 4) Binomial    |

1) Let X and Y be independent and identically distributed normal random variables. Then the characteristic function of X and Y is given by

$$\Phi_X(\omega) = e^{j\eta\omega - \sigma^2\omega^2/2} \quad (11.2.1)$$

The characteristic function of Z is given by

$$\Phi_Z(\omega) = \Phi_X^2(\omega) \quad (11.2.2)$$

$$= e^{2j\eta\omega - \sigma^2\omega^2} \quad (11.2.3)$$

Thus Z is a normal random variable with parameters  $2\eta$  and  $2\sigma^2$ . Thus option (1) is correct.

2) Let X and Y be independent and identically distributed exponential random variables. Then the characteristic function of X and Y is given by

$$\Phi_X(\omega) = \frac{\lambda}{1-j\omega} \quad (11.2.4)$$

The characteristic function of Z is given by

$$\Phi_Z(\omega) = \Phi_X^2(\omega) \quad (11.2.5)$$

$$= \frac{\lambda^2}{(1-j\omega)^2} \quad (11.2.6)$$

Thus Z is not an exponential random variable. Therefore option (2) is wrong.

3) Let X and Y be independent and identically distributed uniform random variables such that  $X, Y \sim U(a,b)$ . Then the characteristic function of X and Y is given by

$$\Phi_X(\omega) = \frac{e^{jb\omega} - e^{ja\omega}}{j\omega(b-a)} \quad (11.2.7)$$

The characteristic function of  $Z$  is given by

$$\Phi_Z(\omega) = \Phi_X^2(\omega) \quad (11.2.8)$$

$$= -\frac{(e^{jb\omega} - e^{ja\omega})^2}{\omega^2(b-a)^2} \quad (11.2.9)$$

Thus  $Z$  is not a uniform random variable.

Thus option (3) is wrong.

- 4) Let  $X$  and  $Y$  be independent and identically distributed binomial random variables. Then the characteristic function of  $X$  and  $Y$  is given by

$$\Phi_X(\omega) = (pe^{j\omega} + q)^n \quad (11.2.10)$$

The characteristic function of  $Z$  is given by

$$\Phi_Z(\omega) = \Phi_X^2(\omega) \quad (11.2.11)$$

$$= (pe^{j\omega} + q)^{2n} \quad (11.2.12)$$

Thus  $Z$  is a binomial random variable with parameter  $2n$ . Thus option (4) is correct.

The following figures show the experimental distributions for  $Z$  in each case. The simulation length was kept one million.




Fig. 11.2.1:  $Z$  when  $X$  is standard normal

- 11.3.  $N, A_1, A_2, \dots$  are independent real valued random variables such that

$$\Pr(N = k) = (1 - p)p^k, k = 0, 1, 2, 3, \dots \quad (11.3.1)$$

where  $0 < p < 1$  and  $\{A_i : i = 1, 2, \dots\}$  is a sequence of independent and identically




Fig. 11.2.2:  $Z$  when  $X$  is exponential with  $\lambda = 1$




Fig. 11.2.3:  $Z$  when  $X \sim U(0,1)$




Fig. 11.2.4:  $Z$  when  $X \sim B(100,0.5)$

distributed bounded random variables. Let

$$X(w) = \begin{cases} 0 & \text{if } N(w) = 0 \\ \sum_{j=1}^k A_j & \text{if } N(w) = k, k = 1, 2, 3 \dots \end{cases} \quad (11.3.2)$$

Which of the following are necessarily correct?

- a)  $X$  is a bounded random variable.  
b) Moment generating function  $m_X$  of  $X$  is

$$m_X(t) = \frac{1-p}{1-pm_A(t)}, t \in \mathbb{R}, \quad (11.3.3)$$

where  $m_A$  is moment generating function of  $A_1$ .

- c) Characteristic function  $\varphi_X$  of  $X$  is

$$\varphi_X(t) = \frac{1-p}{1-p\varphi_A(t)}, t \in \mathbb{R}, \quad (11.3.4)$$

where  $\varphi_A$  is the characteristic function of  $A_1$ .

- d)  $X$  is symmetric about 0.

## 12 MARKOV CHAIN

### 12.1. Step 1. Flip a coin twice.

**Step 2.** If the outcomes are (TAILS, HEADS) then output  $Y$  and stop.

**Step 3.** If the outcomes are either (HEADS, HEADS) or (HEADS, TAILS), then output  $N$  and stop.

**Step 4.** If the outcomes are (TAILS, TAILS), then go to Step 1.

The probability that the output of the experiment is  $Y$  is (upto two decimal places).....

**Solution:** The given problem can be represented using Table 12.1.1 and the Markov chain in Fig. 12.1.1. The state transition matrix for the

State	Description
1	$\{T, T\}$
2	$Y = \{T, H\}$
3	$N = \{\{H, H\}, \{H, T\}\}$

TABLE 12.1.1: States and their notations

Markov chain can be expressed as

$$P = \begin{matrix} & \begin{matrix} 2 & 3 & 1 \end{matrix} \\ \begin{matrix} 2 \\ 3 \\ 1 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0.25 & 0.5 & 0.25 \end{bmatrix} \end{matrix} \quad (12.1.1)$$




Fig. 12.1.1

Clearly, the state 1 is transient, while 2, 3 are absorbing. Comparing (12.1.1) with the standard form of the state transition matrix

$$P = \begin{matrix} & \begin{matrix} A & N \end{matrix} \\ \begin{matrix} A \\ N \end{matrix} & \begin{bmatrix} I & O \\ R & Q \end{bmatrix} \end{matrix} \quad (12.1.2)$$

where, From (12.1.1) and (12.1.2),

TABLE 12.1.2: Notations and their meanings

Notation	Meaning
$A$	All absorbing states
$N$	All non-absorbing states
$I$	Identity matrix
$O$	Zero matrix
$R, Q$	Other submatrices

$$R = \begin{pmatrix} 0.25 & 0.5 \end{pmatrix}, Q = \begin{pmatrix} 0.25 \end{pmatrix} \quad (12.1.3)$$

The limiting matrix for absorbing Markov chain is

$$\bar{P} = \begin{pmatrix} I & O \\ FR & O \end{pmatrix} \quad (12.1.4)$$

where

$$F = (I - Q)^{-1} = (1 - 0.25)^{-1} = \frac{4}{3} \quad (12.1.5)$$

is called the fundamental matrix of  $P$ . Upon substituting from (12.1.3) in (12.1.5),

$$F = (1 - 0.25)^{-1} = \frac{4}{3} \quad (12.1.6)$$

and

$$FR = \begin{pmatrix} \frac{1}{3} & \frac{2}{3} \end{pmatrix} \quad (12.1.7)$$

which, upon substituting in (12.1.4) yields

$$\bar{P} = \begin{matrix} & 2 & 3 & 1 \\ \begin{matrix} 2 \\ 3 \\ 1 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{3} & \frac{2}{3} & 0 \end{bmatrix} \end{matrix} \quad (12.1.8)$$

$$\therefore \bar{p}_{12} = \frac{1}{3} \quad (12.1.9)$$


12.2. Consider a simple symmetric random walk on integers ,Where from every state  $i$  you to move to  $i-1$  and  $i+1$  with probability half each. Then which of the following are correct?

- The random walk is aperiodic
- The random walk is irreducible
- The random walk is null recurrent
- The random walk is positive recurrent

**Solution:** This is a Markov Chain ,Where the state space consists of the integers ( $i = 0, \pm 1, \pm 2, \pm 3, \dots$ ) and transition probability is given as

$$P_{i,i+1} = p = \frac{1}{2} \quad (12.2.1)$$

$$P_{i,i-1} = q = \frac{1}{2} \quad (12.2.2)$$



Let  $P_{i,j}^n$  denotes the probability of being in state  $j$  after  $n$ th transition starting from state  $i$ .

- We know that for state  $j$  in Markov chain to be **aperiodic** ,Then their exist  $k$  such that  $P_{j,j}^n > 0$  for all  $n \geq k$ . but for to return to same state  $j$  after  $n$  transitions ,Number of forward steps should be equal to Backward

steps , i.e for odd  $n$  in  $(2m + 1)$ form

$$P_{j,j}^{2m+1} = 0 \quad (12.2.3)$$

when  $n$  is even in  $2m$  form

$$P_{j,j}^{2m} = \binom{2m}{m} p^m q^m \quad (12.2.4)$$

$$= \frac{(2m)!}{m!.m!} p^m q^m \quad (12.2.5)$$

,As for odd  $n$   $P_{j,j}^n = 0$  , $P_{j,j}^n > 0$  for all  $n \geq k$  is not possible .which implies all states are

**Periodic**

Option (1) is **incorrect**.

- In a Markov Chain for state  $j$  to be recurrent then it should satisfy following condition

$$\lim_{t \rightarrow \infty} \sum_{n=1}^t P_{j,j}^n = \infty \quad (12.2.6)$$

using Stirling approximation in equation (12.2.5)

$$P_{j,j}^{2m} = \frac{((2m)^{2m+\frac{1}{2}}) \cdot e^{-2m} \cdot (2\pi)^{\frac{1}{2}}}{m^{m+\frac{1}{2}} \cdot e^{-m} \cdot m^{m+\frac{1}{2}} \cdot e^{-m} \cdot 2\pi} \cdot p^m q^m \quad (12.2.7)$$

$$= \frac{(4pq)^{2m}}{(m\pi)^{\frac{1}{2}}} \quad (12.2.8)$$

In this question  $p = \frac{1}{2} = q$ , then using (12.2.3) and (12.2.8)

$$\lim_{t \rightarrow \infty} \sum_{n=1}^t P_{j,j}^n = \sum_{n=2k, k=1}^{\infty} \frac{1}{(\frac{n}{2}\pi)^{\frac{1}{2}}} \quad (12.2.9)$$

Since  $\frac{1}{n^{\frac{1}{2}}}$  is divergent,

$$\lim_{t \rightarrow \infty} \sum_{n=1}^t P_{j,j}^n = \infty \quad (12.2.10)$$

Therefore state  $j$  is recurrent ,as what we calculated is independent of  $j$  ,all states are **recurrent** . The first-passage-time probability,  $f_{i,j}(n)$ , of a Markov chain is the probability, given as

$$f_{i,j}(n) = \Pr(X_n = j, X_{n-1} \neq j, X_{n-2} \neq j, \dots, X_1 \neq j | X_0 = i) \quad (12.2.11)$$

The first-passage time  $T_{j,j}$  from a state  $j$  back to itself is of particular importance. It has the

PMF  $f_{j,j}(n)$  and Distribution function  $F_{j,j}(n)$

$$F_{j,j}(n) = \sum_{k=0}^n f_{j,j}(k) \quad (12.2.12)$$

We Know that all states are recurrent .Now i will find whether they are null recurrent or positive recurrent . For positive recurrent

$$\overline{T}_{j,j} < \infty \quad (12.2.13)$$

For null recurrent

$$\overline{T}_{j,j} = \infty \quad (12.2.14)$$

Where  $\overline{T}_{j,j}$  is mean time to enter state j starting from j. Now calculating  $\overline{T}_{j,j}$  using below formula,

$$\overline{T}_{j,j} = 1 + \sum_{k=0}^n (1 - F_{j,j}(k)) \quad (12.2.15)$$

Using (12.2.15) and (12.2.12),We get

$$\overline{T}_{j,j} = \infty \quad (12.2.16)$$

Therefore all states are null recurrent. Option(3) is **correct**

- c) Since all states are recurrent,they communicate with each other ,therefore Markov chain is irreducible , option (2) is **correct**  
d) As all states are null recurrent , option (4) is **incorrect**

Therefore correct options are **2,3**

12.3. Consider a Markov Chain with state space  $\{0, 1, 2\}$  and transition matrix

$$P = \begin{pmatrix} 0 & 1 & 2 \\ 0 & \frac{1}{4} & \frac{5}{8} & \frac{1}{8} \\ 1 & \frac{1}{4} & 0 & \frac{3}{4} \\ 2 & \frac{1}{2} & \frac{3}{8} & \frac{1}{8} \end{pmatrix} \quad (12.3.1)$$

Then which of the following are true?

- a)  $\lim_{n \rightarrow \infty} p_{12}^{(n)} = 0$   
b)  $\lim_{n \rightarrow \infty} p_{12}^{(n)} = \lim_{n \rightarrow \infty} p_{21}^{(n)}$   
c)  $\lim_{n \rightarrow \infty} p_{22}^{(n)} = \frac{1}{8}$   
d)  $\lim_{n \rightarrow \infty} p_{21}^{(n)} = \frac{1}{3}$

12.4. Consider a Markov chain with state space  $1, 2, \dots, 100$ . Suppose states  $2i$  and  $2j$  communicate with each other and states  $2i-1$  and  $2j-1$  communicate with each other for every  $i, j = 1, 2, \dots, 50$ . Further suppose that  $p_{3,3}^{(2)} \neq 0, p_{4,4}^{(3)} \neq 0$

and  $p_{2,5}^{(7)} \neq 0$ . Then

- a) The Markov chain is irreducible.  
b) The Markov chain is aperiodic.  
c) State 8 is recurrent.  
d) State 9 is recurrent.


**Solution:**

12.5. A standard fair die is rolled until some face other than 5 or 6 turns up. Let  $X$  denote the face value of the last roll. Let  $A = \{X \text{ is even}\}$  and  $B = \{X \text{ is at most } 2\}$  Then,

a)  $\Pr(A \cap B) = 0$       c)  $\Pr(A \cap B) = \frac{1}{4}$

b)  $\Pr(A \cap B) = \frac{1}{6}$       d)  $\Pr(A \cap B) = \frac{1}{3}$

**Solution:** Let us assume the following table.



Let us represent the markov chain diagram in a

TABLE 12.5.1

state 1	state 2	state 3	state 4
$X = 5 \text{ or } X = 6$	$X = 2$	$X = 1 \text{ or } X = 3 \text{ or } X = 4$	end

matrix. Let  $P_{ij}$  represent the element of a matrix which is in  $i^{th}$  row and  $j^{th}$  column. The value of  $P_{ij}$  is equal to probability of transition from state  $i$  to state  $j$

$$P = \begin{bmatrix} \frac{1}{3} & \frac{1}{6} & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (12.5.1)$$

We need the probability that  $X = 2$ . Hence required probability is

$$P_{12} + (P_{12})^2 + \cdots + \infty \quad (12.5.2)$$

where  $P_{12}^n$  represents the 1st row, 2nd column element in the  $P^n$

$$P^2 = \begin{bmatrix} \frac{1}{3} & \frac{1}{6} & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \times \begin{bmatrix} \frac{1}{3} & \frac{1}{6} & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (12.5.3)$$

$$= \begin{bmatrix} \frac{1}{9} & \frac{1}{18} & \frac{1}{6} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (12.5.4)$$

$$P^3 = (P^2)(P^1) \quad (12.5.5)$$

$$= \begin{bmatrix} \frac{1}{9} & \frac{1}{18} & \frac{1}{6} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \times \begin{bmatrix} \frac{1}{3} & \frac{1}{6} & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (12.5.6)$$

$$= \begin{bmatrix} \frac{1}{27} & \frac{1}{54} & \frac{1}{18} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (12.5.7)$$

From above we can notice that each time  $P_{12}$  reduces by  $\frac{1}{3}$ . Hence from (12.5.2),

$$\sum_{i=0}^{\infty} \left(\frac{1}{3}\right)^i \frac{1}{6} \quad (12.5.8)$$

From Geometric progression we can write, required probability  $= \frac{1}{4} \therefore$  **option C is correct**

12.6. Consider a Markov chain with five states  $\{1, 2, 3, 4, 5\}$  and transition matrix

$$P = \begin{pmatrix} \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 \\ 0 & \frac{1}{7} & 0 & 0 & \frac{6}{7} \\ \frac{1}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} \\ \frac{1}{3} & 0 & 0 & \frac{2}{3} & 0 \\ 0 & \frac{5}{8} & 0 & 0 & \frac{3}{8} \end{pmatrix} \quad (12.6.1)$$

Which of the following are true?

- a) 3 and 1 are in the same communicating class
- b) 1 and 4 are in the same communicating class
- c) 4 and 2 are in the same communicating class
- d) 2 and 5 are in the same communicating class

**Solution:** See Tables 12.6.1 and 12.6.2



Accessibility of states in Markov's chain	We say that state $j$ is accessible from state $i$ , written as $i \rightarrow j$ , if $p_{ij}^{(n)} > 0$ for some $n$ . Every state is accessible from itself since $p_{ii}^{(0)} = 1$
Communication between states	Two states $i$ and $j$ are said to communicate, written as $i \leftrightarrow j$ , if they are accessible from each other. In other words,  $i \leftrightarrow j \text{ means } i \rightarrow j \text{ and } j \rightarrow i.$
Communicating class	For each Markov chain, there exists a unique decomposition of the state space $S$ into a sequence of disjoint subsets $C_1, C_2, \dots$ ,  $S = \bigcup_{i=1}^{\infty} C_i$  in which each subset has the property that all states within it communicate. Each such subset is called a communication class of the Markov chain.

TABLE 12.6.1: Definition and Result used

Drawing Transition diagram	
Checking whether the states 3 and 1 are in the same communicating class	<p>Here,        State 1 is accessible from the state 3.        But, State 3 is not accessible from the state 1        i.e. <math>3 \rightarrow 1, 1 \nrightarrow 3</math>  <math>\Rightarrow \boxed{3 \leftrightarrow 1}</math></p> <p>Therefore, 3 and 1 are not in the same communicating class.</p>
Checking whether the states 1 and 4 are in the same communicating class	<p>Here,        State 1 is accessible from the state 4.        Also, State 4 is accessible from the state 1        i.e. <math>3 \rightarrow 1, 1 \rightarrow 3</math>  <math>\Rightarrow \boxed{3 \leftrightarrow 1}</math></p> <p>Therefore, 1 and 4 are in the same communicating class.</p>
Checking whether the states 4 and 2 are in the same communicating class	<p>Here,        State 2 is not accessible from the state 4.        Also, State 4 is not accessible from the state 2        i.e. <math>4 \nrightarrow 2, 2 \nrightarrow 4</math></p>

	$\Rightarrow \boxed{4 \leftrightarrow 2}$ <p>Therefore, 4 and 2 are not in the same communicating class.</p>
Checking whether the states 2 and 5 are in the same communicating class	<p>Here,          State 2 is accessible from the state 5.          Also, State 5 is accessible from the state 2          i.e. <math>5 \rightarrow 2, 2 \rightarrow 5</math>  <math display="block">\Rightarrow \boxed{2 \leftrightarrow 5}</math></p> <p>Therefore, 2 and 5 are in the same communicating class.</p>
Conclusion	<p>Communication classes are:</p> $\boxed{S = \{1, 4\} \cup \{3\} \cup \{2, 5\}}$ <p>Option 2) and 4) are true.</p>

TABLE 12.6.2: Solution

12.7.  $A$  and  $B$  play a game of tossing a fair coin.  $A$  starts the game by tossing the coin once and  $B$  then tosses the coin twice, followed by  $A$  tossing the coin once and  $B$  tossing the coin twice and this continues until a head turns up. Whoever gets the first head wins the game. Then,

- a)  $P(B \text{ Wins}) > P(A \text{ Wins})$
- b)  $P(B \text{ Wins}) = 2P(A \text{ Wins})$
- c)  $P(A \text{ Wins}) > P(B \text{ Wins})$
- d)  $P(A \text{ Wins}) = 1 - P(B \text{ Wins})$

**Solution:** Given, a fair coin is tossed till heads turns up.

$$p = \frac{1}{2}, q = \frac{1}{2} \quad (104.1)$$

Let's define a Markov chain  $\{X_n, n = 0, 1, 2, \dots\}$ , where  $X_n \in S = \{1, 2, 3, 4, 5\}$ , such that The state transition matrix for the Markov

TABLE 12.7.1: States and their notations

Notation	State
$S = 1$	$A$ 's turn
$S = 2$	$B$ 's first turn
$S = 3$	$B$ 's second turn
$S = 4$	$A$ wins
$S = 5$	$B$ wins

chain is

$$P = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} & \begin{bmatrix} 0 & 0.5 & 0 & 0.5 & 0 \\ 0 & 0 & 0.5 & 0 & 0.5 \\ 0.5 & 0 & 0 & 0 & 0.5 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \end{matrix} \quad (104.2)$$

Clearly, the states 1, 2, 3 are transient, while 4, 5 are absorbing. The standard form of a state transition matrix is

$$P = \begin{matrix} & \begin{matrix} A & N \end{matrix} \\ \begin{matrix} A \\ N \end{matrix} & \begin{bmatrix} I & O \\ R & Q \end{bmatrix} \end{matrix} \quad (104.3)$$

where, Converting (104.2) to standard form, we

TABLE 12.7.2: Notations and their meanings

Notation	Meaning
$A$	All absorbing states
$N$	All non-absorbing states
$I$	Identity matrix
$O$	Zero matrix
$R, Q$	Other submatrices

get

$$P = \begin{matrix} & \begin{matrix} 4 & 5 & 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 4 \\ 5 \\ 1 \\ 2 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0.5 & 0 & 0 & 0.5 & 0 \\ 0 & 0.5 & 0 & 0 & 0.5 \\ 0 & 0.5 & 0.5 & 0 & 0 \end{bmatrix} \end{matrix} \quad (104.4)$$

From (104.4),

$$R = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \\ 0 & 0.5 \end{bmatrix}, Q = \begin{bmatrix} 0 & 0.5 & 0 \\ 0 & 0 & 0.5 \\ 0.5 & 0 & 0 \end{bmatrix} \quad (104.5)$$

The limiting matrix for absorbing Markov chain is

$$\bar{P} = \begin{bmatrix} I & O \\ FR & O \end{bmatrix} \quad (104.6)$$

where,

$$F = (I - Q)^{-1} \quad (104.7)$$

is called the fundamental matrix of  $P$ .

On solving, we get

$$\bar{P} = \begin{matrix} & \begin{matrix} 4 & 5 & 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 4 \\ 5 \\ 1 \\ 2 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0.5714 & 0.4285 & 0 & 0 & 0 \\ 0.1428 & 0.8571 & 0 & 0 & 0 \\ 0.2857 & 0.7142 & 0 & 0 & 0 \end{bmatrix} \end{matrix} \quad (104.8)$$

A element  $\bar{p}_{ij}$  of  $\bar{P}$  denotes the absorption probability in state  $j$ , starting from state  $i$ . Then,

a)  $Pr(A \text{ wins}) = \bar{p}_{14} \approx 0.5714$

b)  $Pr(B \text{ wins}) = \bar{p}_{15} \approx 0.4285$

$$\therefore \bar{p}_{14} > \bar{p}_{15} \quad (104.9)$$

Also, in  $\bar{P}$ , all the terms in every row should sum to 1.

$$\Rightarrow \bar{p}_{14} + \bar{p}_{15} + 0 + 0 + 0 = 1 \quad (104.10)$$

$$\therefore \bar{p}_{14} = 1 - \bar{p}_{15} \quad (104.11)$$

Therefore, options 3), 4) are correct.

12.8. Consider a Markov chain with state space  $1, 2, \dots, 100$ . Suppose states  $2i$  and  $2j$  communicate with each other and states  $2i-1$  and  $2j-1$  communicate with each other for every  $i, j = 1, 2, \dots, 50$ . Further suppose that  $p_{3,3}^{(2)} < 0, p_{4,4}^{(3)} < 0$  and  $p_{2,5}^{(7)} < 0$ . Then


- The Markov chain is irreducible.
- The Markov chain is aperiodic.
- State 8 is recurrent.
- State 9 is recurrent.

**Solution:**

### 13 INEQUALITIES

13.1. Let  $X'_i$ 's be independent random variables such that  $X'_i$ 's are symmetric about 0

**Markov chain diagram**



and  $var(X_i) = 2i - 1$ , for  $i \geq 1$ . then,  
 $\lim_{n \rightarrow \infty} Pr(X_1 + X_2 + \dots + X_n > n \log n)$

- does not exist.
- equals  $\frac{1}{2}$ .
- equals 1.
- equals 0.

**Solution:** Let  $X = X_1 + X_2 + \dots + X_n$ , as  $X'_i$ 's are symmetric about 0. The mean of  $X$  is given by,

$$E[X] = 0 \quad (13.1.1)$$

the variance of  $X$  is given by,

$$\text{var}[X] = \sum_{i=1}^n (2i-1) \quad (13.1.2)$$

$$= \frac{2n(n+1)}{2} - n \quad (13.1.3)$$

$$= n^2 \quad (13.1.4)$$

the standard deviation,

$$\sigma_X = n \quad (13.1.5)$$

Applying Chebyshev's Inequality for the random variable  $X$ , for any  $k > 0$

$$\Pr(|X - E[X]| > k\sigma_X) \leq \frac{1}{k^2} \quad (13.1.6)$$

let  $k = \log n$ , using (13.1.1) and (13.1.5) in (13.1.6),

$$\Pr(|X| > n \log n) \leq \frac{1}{(\log n)^2} \quad (13.1.7)$$

$$\Pr(X > n \log n) + \Pr(X < -n \log n) \leq \frac{1}{(\log n)^2} \quad (13.1.8)$$

As,  $X$  is symmetric about 0,

$$\Pr(X > n \log n) = \Pr(X < -n \log n) \quad (13.1.9)$$

using (13.1.9) in (13.1.8),

$$2 \Pr(X > n \log n) \leq \frac{1}{(\log n)^2} \quad (13.1.10)$$

$$\Pr(X > n \log n) \leq \frac{1}{2(\log n)^2} \quad (13.1.11)$$

as any probability is greater than 0,

$$0 < \Pr(X > n \log n) \leq \frac{1}{2(\log n)^2} \quad (13.1.12)$$

applying sandwich principle to (13.1.12),

$$\lim_{n \rightarrow \infty} 0 < \lim_{n \rightarrow \infty} \Pr(X > n \log n) \leq \lim_{n \rightarrow \infty} \frac{1}{2(\log n)^2} \quad (13.1.13)$$

$$\lim_{n \rightarrow \infty} \Pr(X_1 + X_2 + \dots + X_n > n \log n) = 0 \quad (13.1.14)$$

Hence the option.4 is correct.

13.2. Let  $X_1, X_2, \dots$  be independent random variables each following exponential distribution with mean 1. Then which of the following state-

ments are correct?

- a)  $\Pr(X_n > \log n \text{ for infinitely many } n \geq 1) = 1$
- b)  $\Pr(X_n > 2 \text{ for infinitely many } n \geq 1) = 1$
- c)  $\Pr(X_n > \frac{1}{2} \text{ for infinitely many } n \geq 1) = 0$
- d)  $\Pr(X_n > \log n, X_{n+1} > \log(n+1) \text{ for infinitely many } n \geq 1) = 0$

**Solution:** PDF of  $X_i$  is

$$f_{X_i}(x) = \begin{cases} \lambda_i e^{-\lambda_i x}, & x \geq 0 \\ 0, & x < 0 \end{cases}$$

Mean of  $X_i$  is expressed as

$$\begin{aligned} E(X_i) &= \int_{-\infty}^{\infty} x f_{X_i}(x) dx \\ &= \int_{-\infty}^0 0 dx + \int_0^{\infty} x \lambda_i e^{-\lambda_i x} \\ &= \frac{1}{\lambda_i} \end{aligned} \quad (13.2.1)$$

From (13.2.1) and  $E(X_i) = 1$ , we have  $\lambda_i = 1 \forall i \geq 1$  Now, for some constant  $c \geq 0$

$$\begin{aligned} \Pr(X_n > c) &= \int_c^{\infty} f_{X_n}(x) dx \\ &= \int_c^{\infty} e^{-x} dx \\ &= e^{-c} \end{aligned} \quad (13.2.2)$$

**Borel-Cantelli Lemma:**

Let  $E_1, E_2, \dots$  be a sequence of events in some probability space. The Borel-Cantelli lemma states that, if the sum of the probabilities of the events  $E_n$  is finite

$$\sum_{n=1}^{\infty} \Pr(E_n) < \infty \quad (13.2.3)$$

then the probability that infinitely many of them occur is 0

$$\Pr\left(\limsup_{n \rightarrow \infty} E_n\right) = 0 \quad (13.2.4)$$

**Second Borel-Cantelli Lemma:**

If the events  $E_n$  are independent and the sum of the probabilities of the  $E_n$  diverges to infinity, then the probability that infinitely many of them occur is 1. If for independent events

$E_1, E_2, \dots$

$$\sum_{n=1}^{\infty} \Pr(E_n) = \infty \quad (13.2.5)$$

Then

$$\Pr\left(\limsup_{n \rightarrow \infty} E_n\right) = 1 \quad (13.2.6)$$

- a) **Option 1:** We can say the events  $X_n > \log n$  are independent  $\forall n \geq 1$  as  $X_n$  are independent random variable.

From (13.2.2)

$$\begin{aligned} \sum_{n=1}^{\infty} \Pr(X_n > \log n) &= \sum_{n=1}^{\infty} e^{-\log n} \\ &= \sum_{n=1}^{\infty} \frac{1}{n} \\ &= \infty \text{ (Cauchy's Criterion)} \end{aligned}$$

Now, from second Borel-Cantelli lemma

$$\begin{aligned} &\Pr(X_n > \log n \text{ for infinitely many } n \geq 1) \\ &= \Pr\left(\limsup_{n \rightarrow \infty} X_n > \log n\right) \\ &= 1 \end{aligned}$$

$\therefore$  Option 1 is correct.

- b) **Option 2:** We can say the events  $X_n > 2$  are independent  $\forall n \geq 1$  as  $X_n$  are independent random variable.

From (13.2.2)

$$\begin{aligned} \sum_{n=1}^{\infty} \Pr(X_n > 2) &= \sum_{n=1}^{\infty} e^{-2} \\ &= \infty \end{aligned}$$

Now, from second Borel-Cantelli lemma

$$\begin{aligned} &\Pr(X_n > 2 \text{ for infinitely many } n \geq 1) \\ &= \Pr\left(\limsup_{n \rightarrow \infty} X_n > 2\right) \\ &= 1 \end{aligned}$$

$\therefore$  Option 2 is correct.

- c) **Option 3:** We can say the events  $X_n > \frac{1}{2}$  are independent  $\forall n \geq 1$  as  $X_n$  are independent random variable.

From (13.2.2)

$$\begin{aligned} \sum_{n=1}^{\infty} \Pr\left(X_n > \frac{1}{2}\right) &= \sum_{n=1}^{\infty} e^{-\frac{1}{2}} \\ &= \infty \end{aligned}$$

Now, from second Borel-Cantelli lemma

$$\begin{aligned} &\Pr\left(X_n > \frac{1}{2} \text{ for infinitely many } n \geq 1\right) \\ &= \Pr\left(\limsup_{n \rightarrow \infty} X_n > \frac{1}{2}\right) \\ &= 1 \end{aligned}$$

$\therefore$  Option 3 is incorrect.

- d) **Option 4:** We can say the events  $X_n > \log n$  are independent  $\forall n \geq 1$  as  $X_n$  are independent random variable.

Let the event  $X_n > \log n, X_{n+1} > \log(n+1)$  be represented by  $E_n$

From (13.2.2)

$$\begin{aligned} &\sum_{n=1}^{\infty} \Pr(E_n) \\ &= \sum_{n=1}^{\infty} \Pr(X_n > \log n) \Pr(X_{n+1} > \log(n+1)) \\ &= \sum_{n=1}^{\infty} e^{-\log n} e^{-\log(n+1)} \\ &= \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \\ &= \sum_{n=1}^{\infty} \frac{1}{n} - \frac{1}{n+1} \\ &= 1 \end{aligned} \quad (13.2.7)$$

Now, from Borel-Cantelli lemma

$$\begin{aligned} &\Pr(E_n \text{ for infinitely many } n \geq 1) \\ &= \Pr\left(\limsup_{n \rightarrow \infty} (X_n > \log n, X_{n+1} > \log(n+1))\right) \\ &= 0 \end{aligned}$$

$\therefore$  Option 4 is correct.

**Solution:** Options 1, 2, 4

## 14 CONVERGENCE

14.1. Let  $X_1, X_2, \dots$  be independent and identically distributed random variables each following a

uniform distribution on  $(0,1)$ . Denote

$$T_n = \max \{X_1, X_2, \dots, X_n\}. \quad (14.1.1)$$

Then, which of the following statements are true?

- a)  $T_n$  converges to 1 in probability.
- b)  $n(1 - T_n)$  converges in distribution.
- c)  $n^2(1 - T_n)$  converges in distribution.
- d)  $\sqrt{n}(1 - T_n)$  converges to 0 in probability.

**Solution:**

**Definition 3.** Random Sampling : A collection of random variables  $X_1, X_2, \dots, X_n$  is said to be a random sample of size  $n$  if they are independent and identically distributed, i.e,

- a)  $X_1, X_2, \dots, X_n$  are independent random variables
- b) They have the same distribution (Let us denote it by  $F_X(x)$ ), i.e,

$$F_X(x) = F_{X_i}(x), i = 1, 2, \dots, n \forall x \in \mathbb{R} \quad (14.1.2)$$

**Definition 4.** Order Statistics : Given a random sample  $X_1, X_2, \dots, X_n$ , the sequence  $X_{(1)}, X_{(2)}, \dots, X_{(n)}$  is called the order statistics of it. Here,

$$X_{(1)} = \min (X_1, X_2, \dots, X_n) \quad (14.1.3)$$

$$X_{(2)} = \text{the } 2^{\text{nd}} \text{ smallest of } X_1, X_2, \dots, X_n \quad (14.1.4)$$

$$\vdots \quad (14.1.5)$$

$$X_{(n)} = \max (X_1, X_2, \dots, X_n) \quad (14.1.6)$$

**Lemma 14.1.** Distribution of the maximum :

$$f_{T_n}(x) = \begin{cases} nx^{n-1}, & 0 < x < 1 \\ 0, & \text{otherwise} \end{cases} \quad (14.1.7)$$

$$F_{T_n}(x) = \begin{cases} x^n, & 0 < x < 1 \\ 1, & x \geq 1 \\ 0, & \text{otherwise} \end{cases} \quad (14.1.8)$$

**Proof:**

$$F_{X_{(n)}}(x) = \Pr (X_{(n)} \leq x) \quad (14.1.9)$$

$$= \Pr (X_1 \leq x, X_2 \leq x, \dots, X_n \leq x) \quad (14.1.10)$$

$$= \Pr (X_1 \leq x) \Pr (X_2 \leq x) \dots \Pr (X_n \leq x) \quad (14.1.11)$$

$$= [\Pr (X_1 \leq x)]^n \text{ (i.i.d)} \quad (14.1.12)$$

$$= [F_X(x)]^n \quad (14.1.13)$$

and

$$f_{X_{(n)}}(x) = \frac{d}{dx} (F_{X_{(n)}}(x)) = \frac{d}{dx} ([F_X(x)]^n) \quad (14.1.14)$$

$$= n ([F_X(x)]^{n-1}) \frac{d}{dx} (F_X(x)) \quad (14.1.15)$$

$$= n [F_X(x)]^{n-1} f_X(x) \quad (14.1.16)$$

$\therefore$

$$f_{X_i}(x) = \begin{cases} 1, & 0 < x < 1 \\ 0, & \text{otherwise} \end{cases}, \quad (14.1.17)$$

$$F_{X_i}(x) = \begin{cases} x, & 0 < x < 1 \\ 1, & x \geq 1 \\ 0, & \text{otherwise,} \end{cases} \quad (14.1.18)$$

$\forall i \in \mathbb{N}$ . Substituting the above in (14.1.16) and (14.1.13) yields (14.1.7) and (14.1.8) respectively. Then, as  $T_n = \max\{X_1, X_2, \dots, X_n\} = X_{(n)}$ ,

**Lemma 14.2.** If  $Y = aX + b$  and  $a < 0$ , then

$$F_Y(y) = 1 - F_X\left(\frac{y-b}{a}\right) \quad (14.1.19)$$

**Definition 5.** Convergence in Probability : A sequence of random variables  $X_1, X_2, X_3, \dots$  converges in probability to a random variable  $X$ , shown by  $X_n \xrightarrow{p} X$ , if

$$\lim_{n \rightarrow \infty} \Pr (|X_n - X| \geq \epsilon) = 0, \forall \epsilon > 0 \quad (14.1.20)$$

**Definition 6.** Convergence in Distribution : A sequence of random variables  $X_1, X_2, X_3, \dots$  converges in distribution to a random variable  $X$ , shown by  $X_n \xrightarrow{d} X$ , if

$$\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x) \quad (14.1.21)$$

for all  $x$  at which  $F_X(x)$  is continuous.



a)

$$\begin{aligned}\lim_{n \rightarrow \infty} \Pr(|T_n - 1| \geq \epsilon) &= \lim_{n \rightarrow \infty} \Pr(1 - T_n \geq \epsilon) \\ &= \lim_{n \rightarrow \infty} \Pr(T_n \leq 1 - \epsilon) = \lim_{n \rightarrow \infty} F_{T_n}(1 - \epsilon)\end{aligned}\quad (14.1.22)$$

$$\therefore F_{T_n}(1 - \epsilon) = \begin{cases} (1 - \epsilon)^n, & 0 < \epsilon < 1 \\ 0, & \epsilon \geq 1 \end{cases}\quad (14.1.23)$$

and

$$\therefore \lim_{n \rightarrow \infty} (1 - \epsilon)^n = 0 \text{ for } 0 < \epsilon < 1 \quad (14.1.24)$$

$$(14.1.25)$$

from (14.1.24), (14.1.23) and (14.1.22),

$$\lim_{n \rightarrow \infty} \Pr(|T_n - 1| \geq \epsilon) = 0, \forall \epsilon > 0 \quad (14.1.26)$$

 $\therefore T_n$  converges to 1 in probability.b) Substituting  $a = -n, b = n$  in (14.1.19),

$$F_{n(1-T_n)}(x) = 1 - F_{T_n}\left(1 - \frac{x}{n}\right) \quad (14.1.27)$$

$$(14.1.28)$$

where

$$F_{T_n}\left(1 - \frac{x}{n}\right) = \begin{cases} \left(1 - \frac{x}{n}\right)^n, & 0 < x < n \\ 1, & x \leq 0 \\ 0, & x \geq n \end{cases}\quad (14.1.29)$$

$$\text{from (14.1.8)} \quad (14.1.30)$$

$$\therefore \lim_{n \rightarrow \infty} \left(1 - \frac{y}{n}\right)^n = e^{-y}, \quad (14.1.31)$$

$$(14.1.32)$$

$$\therefore \lim_{n \rightarrow \infty} F_{T_n}\left(1 - \frac{x}{n}\right) = \begin{cases} e^{-x}, & x > 0 \\ 1, & x \leq 0 \end{cases}\quad (14.1.33)$$

$$\Rightarrow \lim_{n \rightarrow \infty} F_{n(1-T_n)}(x) = 1 - \lim_{n \rightarrow \infty} F_{T_n}\left(1 - \frac{x}{n}\right) \quad (14.1.34)$$

which can be expressed as

$$\therefore \lim_{n \rightarrow \infty} F_{n(1-T_n)}(x) = \begin{cases} 1 - e^{-x}, & x > 0 \\ 0, & x \leq 0 \end{cases}\quad (14.1.35)$$

 $\therefore n(1 - T_n)$  converges in distribution to the random variable  $X \sim \text{Exponential}(1)$ .c) Substituting  $a = -n^2, b = n^2$  in (14.1.19),

$$F_{n^2(1-T_n)}(x) = 1 - F_{T_n}\left(1 - \frac{x}{n^2}\right) \quad (14.1.36)$$

$$F_{T_n}\left(1 - \frac{x}{n^2}\right) = \begin{cases} \left(1 - \frac{x}{n^2}\right)^n, & 0 < x < n^2 \\ 1, & x \leq 0 \\ 0, & x \geq n^2 \end{cases}\quad (14.1.37)$$

$$= \begin{cases} 1, & x > 0 \\ 1, & x \leq 0 \end{cases}\quad (14.1.38)$$

$$\therefore \lim_{n \rightarrow \infty} \left(1 - \frac{x}{n^2}\right)^n = 1 \quad (14.1.39)$$

yielding

$$\lim_{n \rightarrow \infty} F_{n^2(1-T_n)}(x) = \begin{cases} 0, & x > 0 \\ 0, & x \leq 0 \end{cases}\quad (14.1.40)$$

which is not a valid CDF. Hence,  $n^2(1 - T_n)$  does not converge in distribution.

d)

$$\begin{aligned}\lim_{n \rightarrow \infty} \Pr(|\sqrt{n}(1 - T_n) - 0| \geq \epsilon) &= \lim_{n \rightarrow \infty} \Pr\left(1 - T_n \geq \frac{\epsilon}{\sqrt{n}}\right) \\ &= \lim_{n \rightarrow \infty} \Pr\left(T_n \leq 1 - \frac{\epsilon}{\sqrt{n}}\right) \\ &= \lim_{n \rightarrow \infty} F_{T_n}\left(1 - \frac{\epsilon}{\sqrt{n}}\right) \\ &= \begin{cases} \left(1 - \frac{\epsilon}{\sqrt{n}}\right)^n, & 0 < \epsilon < \sqrt{n} \\ 0, & \epsilon \geq \sqrt{n} \end{cases}\end{aligned}\quad (14.1.41)$$

$$\begin{aligned} \therefore \lim_{n \rightarrow \infty} \left(1 - \frac{\epsilon}{\sqrt{n}}\right)^n &= 0 \text{ for } 0 < \epsilon < \sqrt{n}, \\ \lim_{n \rightarrow \infty} \Pr(|\sqrt{n}(1 - T_n) - 0| \geq \epsilon) &= 0, \forall \epsilon > 0 \end{aligned} \quad (14.1.42)$$

$\therefore \sqrt{n}(1 - T_n)$  converges to 0 in probability.

Hence, options 1), 2), 4) are correct.

14.2. Let  $\{X_i\}_{i \geq 1}$  be a sequence of i.i.d. random variables with  $E(X_i) = 0$  and  $V(X_i) = 1$ . Which of the following are true?

- a)  $\frac{1}{n} \sum_{i=1}^n X_i^2 \rightarrow 0$  in probability
- b)  $\frac{1}{n^{3/4}} \sum_{i=1}^n X_i \rightarrow 0$  in probability
- c)  $\frac{1}{n^{1/2}} \sum_{i=1}^n X_i \rightarrow 0$  in probability
- d)  $\frac{1}{n} \sum_{i=1}^n X_i^2 \rightarrow 1$  in probability

**Solution:**

**Lemma 14.3.** Strong Law of Large Numbers: Let  $X_1, X_2, \dots$  be a sequence of i.i.d. random variables, each having finite mean  $E(X_i)$ . Then for any  $\epsilon > 0$ ,

$$\lim_{n \rightarrow \infty} \Pr\left(\left|\frac{1}{n} \sum_{i=1}^n X_i - E(X_i)\right| \geq \epsilon\right) = 0 \quad (14.2.1)$$

Or,  $\frac{1}{n} \sum_{i=1}^n X_i$  converges in probability to  $E(X_i)$ .

14.3. Let  $\{X_n\}$  be a sequence of independent random variables where the distribution of  $X_n$  is normal with mean  $\mu$  and variance  $n$  for  $n = 1, 2, \dots$ . Define

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i \quad (14.3.1)$$

$$S_n = \frac{\sum_{i=1}^n \frac{1}{i} X_i}{\sum_{i=1}^n \frac{1}{i}} \quad (14.3.2)$$

Which of the following are true?

- a)  $E(\bar{X}_n) = E(S_n)$  for sufficiently large  $n$
- b)  $\text{Var}(S_n) < \text{Var}(\bar{X}_n)$  for sufficiently large  $n$
- c)  $\bar{X}_n$  is consistent for  $\mu$
- d)  $\bar{X}_n$  is sufficient for  $\mu$

**Solution:** As  $X_i$  for  $i = 1, 2, \dots, n$  are independent random variables we can use this property to state

$$\text{Var}\left(\sum_{i=1}^n g(X_i)\right) = \sum_{i=1}^n \text{Var}(g(X_i)) \quad (14.3.3)$$

**Definition 7.** Random Sample: The random variables  $X_1, X_2, X_3, \dots, X_n$  are said to be random sample if

- a) the  $X_i$ 's are independent
- b)  $F_{X_1}(x) = F_{X_2}(x) = \dots = F_{X_n}(x) = F_X(x)$
- c)  $EX_i = EX = \mu < \infty$
- d)  $0 < \text{Var}(X_i) = \text{Var}(X) = \sigma^2 < \infty$

Let  $n = 2$  and hence  $X_1$  and  $X_2$  are sequence of independent random variables and

$$\text{Var}(X_1) = 1 \quad (14.3.4)$$

$$\text{Var}(X_2) = 2 \quad (14.3.5)$$

$$\text{Var}(X_1) \neq \text{Var}(X_2) \quad (14.3.6)$$

The equation (14.3.6) doesn't follow point(14.3d) in definition(7) and hence the random variables are not a random sample.

a) Expectation of  $\bar{X}_n$  and  $S_n$

$$E(\bar{X}_n) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) \quad (14.3.7)$$

$$= \frac{1}{n} \sum_{i=1}^n E(X_i) \quad (14.3.8)$$

$$= \frac{1}{n} \sum_{i=1}^n \mu \quad (14.3.9)$$

$$= \mu \quad (14.3.10)$$

$$E(S_n) = E\left(\frac{\sum_{i=1}^n \frac{1}{i} X_i}{\sum_{i=1}^n \frac{1}{i}}\right) \quad (14.3.11)$$

$$= \frac{1}{\left(\sum_{i=1}^n \frac{1}{i}\right)} \sum_{i=1}^n E\left(\frac{1}{i} X_i\right) \quad (14.3.12)$$

$$= \frac{1}{\left(\sum_{i=1}^n \frac{1}{i}\right)} \sum_{i=1}^n \frac{\mu}{i} \quad (14.3.13)$$

$$= \mu \quad (14.3.14)$$

From (14.3.10) and (14.3.14) we get option(14.3a) is correct.

b) Variance of  $\bar{X}_n$  and  $S_n$  using (14.3.3)

$$\text{Var}(\bar{X}_n) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) \quad (14.3.15)$$

$$= \frac{1}{n^2} \left( \sum_{i=1}^n \text{Var}(X_i) \right) \quad (14.3.16)$$

$$= \frac{1}{n^2} \left( \sum_{i=1}^n i \right) \quad (14.3.17)$$

$$= \frac{1}{2} + \frac{1}{2n} \quad (14.3.18)$$

$$\text{Var}(S_n) = \text{Var}\left(\frac{\sum_{i=1}^n \frac{1}{i} X_i}{\sum_{i=1}^n \frac{1}{i}}\right) \quad (14.3.19)$$

$$= \frac{1}{\left(\sum_{i=1}^n \frac{1}{i}\right)^2} \sum_{i=1}^n \frac{1}{i^2} \text{Var}(X_i) \quad (14.3.20)$$

$$= \frac{1}{\left(\sum_{i=1}^n \frac{1}{i}\right)^2} \sum_{i=1}^n \frac{1}{i^2} i \quad (14.3.21)$$

$$= \frac{1}{\sum_{i=1}^n \frac{1}{i}} \quad (14.3.22)$$

As  $n$  is sufficiently large

$$\text{Var}(\bar{X}_n) = \frac{1}{2} \quad (14.3.23)$$

$$\text{Var}(S_n) = 0 \quad (14.3.24)$$

$$\text{Var}(S_n) < \text{Var}(\bar{X}_n) \quad (14.3.25)$$

from (14.3.25) we get option(14.3b) as correct.

c)

**Definition 8. Point Estimator :** Let  $\theta$  be an unknown fixed(non-random) parameter be estimated. To estimate  $\theta$  we define a point estimator  $\hat{\Theta}$  that is a function of the random sample  $X_1, X_2, X_3, \dots, X_n$  i.e.,

$$\hat{\Theta} = h(X_1, X_2, \dots, X_n) \quad (14.3.26)$$

**Definition 9. Consistent Estimator :** Let  $\hat{\Theta}_1, \hat{\Theta}_2, \dots, \hat{\Theta}_n, \dots$  be a sequence of point estimators of  $\theta$ . We say that  $\hat{\Theta}_n$  is a consistent estimator of  $\theta$ , if

$$\lim_{n \rightarrow \infty} P(|\hat{\Theta}_n - \theta| \geq \epsilon) = 0, \text{ for all } \epsilon > 0. \quad (14.3.27)$$

From (14.3.6) as given data is not a random sample we don't define point estimator and

hence option(14.3c) is incorrect.

d)

**Definition 10. Statistic :** A statistic is a function  $T = r(X_1, X_2, \dots, X_n)$  of the random sample  $X_1, X_2, \dots, X_n$ .

**Definition 11. Sufficient Statistics :** A statistic  $t = T(X)$  is sufficient for  $\theta$  if the conditional probability distribution of data  $X$ , given the statistic  $t = T(X)$ , doesn't depend on the parameter  $\theta$ .

Equation (14.3.6) suggests that given data is not a random sample we don't define statistic and hence option(14.3d) is incorrect.

Hence option(14.3a) and option(14.3b) are correct.

14.4. Let  $X_1, X_2, \dots$  be i.i.d.  $N(0, 1)$  random variables. Let

$$S_n = X_1^2 + X_2^2 + \dots + X_n^2 \forall n \geq 1. \quad (14.4.1)$$

Which of the following statements are correct?

a)

$$\frac{S_n - n}{\sqrt{2}} \sim N(0, 1) \quad \forall n \geq 1 \quad (14.4.2)$$

b)

$$\forall \epsilon > 0, \Pr\left(\left|\frac{S_n}{n} - 2\right| > \epsilon\right) \rightarrow 0, n \rightarrow \infty \quad (14.4.3)$$

c)  $\frac{S_n}{n} \rightarrow 1$  with probability 1

d)

$$\Pr(S_n \leq n + \sqrt{nx}) \rightarrow \Pr(Y \leq x) \quad \forall x \in \mathbb{R}, Y \sim N(0, 2) \quad (14.4.4)$$

14.5. Let  $X_1, X_2, \dots$  be i.i.d.  $N(0, 1)$  random variables. Let  $S_n = X_1^2 + X_2^2 + \dots + X_n^2, \forall n \geq 1$ . Which of the following statements are correct?

a)  $\frac{S_n - n}{\sqrt{2}} \sim N(0, 1)$  for all  $n \geq 1$

b) For all  $\epsilon > 0, \Pr\left(\left|\frac{S_n}{n} - 2\right| > \epsilon\right) \rightarrow 0$  as  $n \rightarrow \infty$

c)  $\frac{S_n}{n} \rightarrow 1$  with probability 1

d)  $\Pr(S_n \leq n + \sqrt{nx}) \rightarrow \Pr(Y \leq x) \forall x \in \mathbb{R}$ , where  $Y \sim N(0, 2)$

**Solution:**

**Definition 12** (Almost sure convergence). A sequence of random variables  $\{X_n\}_{n \in \mathbb{N}}$  is said to converge almost surely or with probability 1 (denoted by a.s or w.p 1) to  $X$  if

$$\Pr(\omega | X_n(\omega) \rightarrow X(\omega)) = 1 \quad (14.5.1)$$

**Definition 13** (Convergence in probability). A sequence of random variables  $\{X_n\}_{n \in \mathbb{N}}$  is said to converge in probability (denoted by i.p) to  $X$  if

$$\lim_{n \rightarrow \infty} \Pr(|X_n - X| > \epsilon) = 0, \forall \epsilon > 0 \quad (14.5.2)$$

**Theorem 14.1** (Weak law of large numbers). Let  $X_1, X_2, \dots$  be i.i.d random variables with same expectation( $\mu$ ) and finite variance( $\sigma^2$ ). Let  $S_n = X_1 + X_2 + \dots + X_n$ , Then as  $n \rightarrow \infty$

$$\frac{S_n}{n} \xrightarrow{i.p} \mu, \quad (14.5.3)$$

in probability

**Theorem 14.2** (Strong law of large numbers). Let  $X_1, X_2, \dots$  be i.i.d random variables with same expectation( $\mu$ ) and finite variance( $\sigma^2$ ). Let  $S_n = X_1 + X_2 + \dots + X_n$ , Then as  $n \rightarrow \infty$

$$\frac{S_n}{n} \xrightarrow{a.s} \mu, \quad (14.5.4)$$

almost surely.

**Theorem 14.3** (Central limit theorem). The Central limit theorem states that the distribution of the sample approximates a normal distribution as the sample size becomes larger, given that all the samples are equal in size, regardless of the distribution of the individual samples.

Given  $X_1, X_2, \dots$  follow normal distribution with mean 0 and variance 1.

$$f_{X_i}(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}, i \in \{1, 2, \dots\} \quad (14.5.5)$$

As  $X_1, X_2, \dots$  are i.i.d random variables therefore  $X_1^2, X_2^2, \dots$  are also identical and independent. We can write

$$E(X^2) = \text{Var}(X) \quad (14.5.6)$$

a)

$$E\left(\frac{S_n - n}{\sqrt{2}}\right) = E\left(\frac{\sum_i (X_i^2 - 1)}{\sqrt{2}}\right) \quad (14.5.7)$$

$$= \frac{\sum_i E(X_i^2 - 1)}{\sqrt{2}} \quad (14.5.8)$$

From (14.5.6) we can write

$$E\left(\frac{S_n - n}{\sqrt{2}}\right) = 0 \quad (14.5.9)$$

$$\text{Var}\left(\frac{S_n - n}{\sqrt{2}}\right) = \text{Var}\left(\frac{\sum_i (X_i^2 - 1)}{\sqrt{2}}\right) \quad (14.5.10)$$

$$= \frac{\sum_i \text{Var}(X_i^2 - 1)}{\sqrt{2}} \quad (14.5.11)$$

$$\text{Var}(X_i^2 - 1) = \int_{-\infty}^{\infty} (X_i^2 - 1)^2 f_{X_i}(x) dx \quad (14.5.12)$$

$$= \int_{-\infty}^{\infty} (X_i^4 + 1 - 2X_i^2) f_{X_i}(x) dx \quad (14.5.13)$$

$$= 2 \quad (14.5.14)$$

$$\text{Var}\left(\frac{S_n - n}{\sqrt{2}}\right) = n \sqrt{2} \quad (14.5.15)$$

Hence from theorem 14.2 as  $n \rightarrow \infty$

$$\left(\frac{S_n - n}{\sqrt{2}}\right) \sim N(0, n \sqrt{2}) \quad (14.5.16)$$

Hence **Option A is false.**

b) Given

$$S_n = X_1^2 + X_2^2 + \dots + X_n^2, \forall n \geq 1 \quad (14.5.17)$$

Hence from theorem 14.1 we can write

$$\frac{S_n}{n} \xrightarrow{i.p} \text{Var}(X) \quad (14.5.18)$$

$$\Rightarrow \frac{S_n}{n} \xrightarrow{i.p} 1 \quad (14.5.19)$$

in probability. From definition 13 we can write,

$$\Rightarrow \Pr\left(\left|\frac{S_n}{n} - 1\right| > \epsilon\right) \rightarrow 0, \forall \epsilon > 0 \quad (14.5.20)$$

Hence **Option B is false .**

c) Given

$$S_n = X_1^2 + X_2^2 + \dots + X_n^2, \forall n \geq 1 \quad (14.5.21)$$

Hence from theorem 14.1 we can write

$$\frac{S_n}{n} \xrightarrow{i.p} \text{Var}(X) \quad (14.5.22)$$

$$\implies \frac{S_n}{n} \xrightarrow{a.s} 1 \quad (14.5.23)$$

almost surely. From definition 12 we can write,

$$\frac{S_n}{n} \xrightarrow{w.p.1} 1 \quad (14.5.24)$$

with probability 1. Hence **Option C is true.**

d) Consider,

$$E\left(\frac{S_n - n}{\sqrt{n}}\right) = 0 \quad (14.5.25)$$

using (14.5.6) and (14.5.8).

$$\text{Var}\left(\frac{S_n - n}{\sqrt{n}}\right) = \frac{2n}{\sqrt{n}} \quad (14.5.26)$$

$$= 2\sqrt{n}. \quad (14.5.27)$$

using (14.5.14). From theorem 14.3 we can write,

$$\left(\frac{S_n - n}{\sqrt{n}}\right) \sim N(0, 2\sqrt{n}) \quad (14.5.28)$$

$$\Pr\left(\frac{S_n - n}{\sqrt{n}} \leq x\right) = \Pr(S_n \leq n + \sqrt{n}x) \quad (14.5.29)$$

Hence using (14.5.28), **Option D is false.**

14.6. Let  $X_1, X_2, \dots, X_n$  be independent and identically distributed, each having a uniform distribution on  $(0, 1)$ . Let  $S_n = \sum_{i=1}^n X_i$  for  $n \geq 1$ . Then, which of the following statements are true?

- A)  $\frac{S_n}{n \log n} \rightarrow 0$  as  $n \rightarrow \infty$  with probability 1.
- B)  $\Pr\left(\left(S_n > \frac{2n}{3}\right) \text{ occurs for infinitely many } n\right) = 1$
- C)  $\frac{S_n}{\log n} \rightarrow 0$  as  $n \rightarrow \infty$  with probability 1.
- D)  $\Pr\left(\left(S_n > \frac{n}{3}\right) \text{ occurs for infinitely many } n\right) = 1$

**Solution:**

a) Given

$$S_n = \sum_{i=1}^n X_i, n \geq 1 \quad (14.6.1)$$

Symbol	expression/definition
$S_n$	$\sum_{i=1}^n X_i$
$\mu_n$	$\frac{1}{n} \sum_{i=1}^n X_i$
$X$	Independent continuous random variable identical to $X_1, X_2, \dots, X_n$

TABLE 14.6.1: Variables and their definitions

Dividing by  $n$  on both sides

$$\frac{S_n}{n} = \frac{1}{n} \sum_{i=1}^n X_i = \mu_n \quad (14.6.2)$$

It can be said that  $X_1, X_2, \dots, X_n$  are the trials of  $X$ . By definition

$$E[X] = \lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n X_i}{n} = \lim_{n \rightarrow \infty} \frac{S_n}{n} \quad (14.6.3)$$

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = E[X] = \frac{1}{2} \quad (14.6.4)$$

$$\therefore \lim_{n \rightarrow \infty} \frac{S_n}{n \log n} = 0 \quad (14.6.5)$$

b) Using weak law, (14.6.4), and table (14.6.1)

$$\lim_{n \rightarrow \infty} \Pr(|\mu_n - E[X]| > \epsilon) = 0, \forall \epsilon > 0 \quad (14.6.6)$$

$$\lim_{n \rightarrow \infty} \Pr\left(S_n = \frac{n}{2}\right) = 1 \quad (14.6.7)$$

It can be easily implied from (14.6.7) that option B is false.

c) It is easy to observe from (14.6.4) that option C is false.

d) Using (14.6.7), we get

$$\Pr\left(\left(S_n > \frac{n}{3}\right) \text{ occurs for infinitely many } n\right) = 1 \quad (14.6.8)$$

14.7. Let  $U_1, U_2, \dots, U_n$  be independent and identically distributed random variables each having a uniform distribution on  $(0, 1)$ . Then,  $\lim_{n \rightarrow +\infty} \Pr(U_1 + U_2, \dots, U_n \leq \frac{3}{4}n)$

- a) does not exist
- b) exists and equals 0
- c) exists and equals 1
- d) exists and equals  $\frac{3}{4}$

**Solution:** We use Weak law for large num-

bers to solve this problem. Let the collection of identically distributed random variables  $U_1, U_2, \dots, U_n$  have a finite mean  $\mu$  and finite variance  $\sigma^2$ .

$$\mu = E(U_i) \quad \text{for } i \in (1, 2, 3, \dots, n) \quad (14.7.1)$$

Since the distribution is uniform on (0,1),  $\mu = 0.5$ . Let  $M_n$  be the sample mean

$$M_n = \frac{U_1 + U_2 + U_3 + \dots + U_n}{n} \quad (14.7.2)$$

Expected value of  $M_n$  (using (14.7.2) and (14.7.1)) is

$$E(M_n) = \frac{E(U_1 + U_2 + U_3 + \dots + U_n)}{E(n)} \quad (14.7.3)$$

$$= \frac{E(U_1) + E(U_2) + \dots + E(U_n)}{n} \quad (14.7.4)$$

$$= \frac{n \times \mu}{n} \quad (14.7.5)$$

$$= \mu \quad (14.7.6)$$

Variance of  $M$

$$Var(M_n) = \frac{Var(U_1 + U_2 + U_3 + \dots + U_n)}{n^2} \quad (14.7.7)$$

$$= \frac{Var(U_1) + Var(U_2) + \dots + Var(U_n)}{n^2} \quad (14.7.8)$$

$$= \frac{n \times \sigma^2}{n^2} \quad (14.7.9)$$

$$= \frac{\sigma^2}{n} \quad (14.7.10)$$

From Chebyshev inequality, for any  $\epsilon > 0$

$$\Pr(|M_n - \mu| \geq \epsilon) \leq \frac{Var(M_n)}{\epsilon^2} \quad (14.7.11)$$

From (14.7.1) and (14.7.10)

$$\begin{aligned} \Pr\left(\left|\frac{U_1 + U_2 + \dots + U_n}{n} - \mu\right| \geq \epsilon\right) &\leq \frac{\sigma^2}{n \times \epsilon^2} \\ \lim_{n \rightarrow \infty} \Pr\left(\left|\frac{U_1 + U_2 + \dots + U_n}{n} - \mu\right| \geq \epsilon\right) & \leq \lim_{n \rightarrow \infty} \frac{\sigma^2}{n \times \epsilon^2} \leq 0 \quad \text{for fixed } \epsilon > 0 \end{aligned} \quad (14.7.12)$$

But since Probabilities are always non-negative,

$$\lim_{n \rightarrow \infty} \Pr\left(\left|\frac{U_1 + U_2 + \dots + U_n}{n} - \mu\right| \geq \epsilon\right) \rightarrow 0 \quad (14.7.13)$$

This is known as the weak law of large numbers

The inverse of (14.7.13) is also true

$$\lim_{n \rightarrow \infty} \Pr\left(\left|\frac{U_1 + U_2 + \dots + U_n}{n} - \mu\right| \leq \epsilon\right) \rightarrow 1 \quad (14.7.14)$$

$$\left|\frac{U_1 + U_2 + \dots + U_n}{n} - \mu\right| \leq \epsilon \quad \text{as } n \rightarrow \infty \quad (14.7.15)$$

From  $\epsilon, n$  definition of limits, it is clear that

$$\frac{U_1 + U_2 + \dots + U_n}{n} \rightarrow \mu \quad (14.7.16)$$

$$U_1 + U_2 + \dots + U_n \rightarrow n \times \mu \quad \text{as } n \rightarrow \infty \quad (14.7.17)$$

Since  $\mu = \frac{1}{2}$ ,

$$\lim_{n \rightarrow +\infty} U_1 + U_2 + \dots + U_n = \frac{1}{2}n < \frac{3}{4}n \quad (14.7.18)$$

So

$$\lim_{n \rightarrow +\infty} \Pr\left(U_1 + U_2 + \dots + U_n \leq \frac{3}{4}n\right) = 1 \quad (14.7.19)$$

14.8. Let  $X_1, X_2, \dots$  be independent and identically distributed random variables each following a uniform distribution on (0,1). Denote  $T_n = \max\{X_1, X_2, \dots, X_n\}$ . Then, which of the following statements are true?

- a)  $T_n$  converges to 1 in probability.
- b)  $n(1 - T_n)$  converges in distribution.
- c)  $n^2(1 - T_n)$  converges in distribution.
- d)  $\sqrt{n}(1 - T_n)$  converges to 0 in probability.

**Solution:** The PDF, CDF of each  $X_1, X_2, X_3, \dots$  is

$$f_{X_i}(x) = \begin{cases} 1, & 0 < x < 1 \\ 0, & \text{otherwise} \end{cases} \quad (72.1)$$

$$F_{X_i}(x) = \begin{cases} x, & 0 < x < 1 \\ 1, & x \geq 1 \\ 0, & \text{otherwise} \end{cases} \quad (72.2)$$

$\forall i \in \mathbb{N}$ . Then, as  $T_n = \max\{X_1, X_2, \dots, X_n\}$ ,

$$f_{T_n}(x) = \begin{cases} nx^{n-1}, & 0 < x < 1 \\ 0, & \text{otherwise} \end{cases} \quad (72.3)$$

$$F_{T_n}(x) = \begin{cases} x^n, & 0 < x < 1 \\ 1, & x \geq 1 \\ 0, & \text{otherwise} \end{cases} \quad (72.4)$$

NOTE : If  $Y = aX + b$  and  $a < 0$ , then

$$F_Y(y) = 1 - F_X\left(\frac{y-b}{a}\right) \quad (72.5)$$

a) OPTION-1:

Convergence in Probability :

A sequence of random variables  $X_1, X_2, X_3, \dots$  converges in probability to a random variable  $X$ , shown by  $X_n \xrightarrow{p} X$ , if

$$\lim_{n \rightarrow \infty} \Pr(|X_n - X| \geq \epsilon) = 0, \forall \epsilon > 0 \quad (72.6)$$

To evaluate :  $\lim_{n \rightarrow \infty} \Pr(|T_n - 1| \geq \epsilon), \forall \epsilon > 0$

$$\lim_{n \rightarrow \infty} \Pr(|T_n - 1| \geq \epsilon) = \lim_{n \rightarrow \infty} \Pr(1 - T_n \geq \epsilon) \quad (72.7)$$

$$= \lim_{n \rightarrow \infty} \Pr(T_n \leq 1 - \epsilon) = \lim_{n \rightarrow \infty} F_{T_n}(1 - \epsilon) \quad (72.8)$$

$$F_{T_n}(1 - \epsilon) = \begin{cases} (1 - \epsilon)^n, & 0 < \epsilon < 1 \\ 0, & \epsilon \geq 1 \end{cases} \quad (72.9)$$

$$\therefore \lim_{n \rightarrow \infty} (1 - \epsilon)^n = 0 \text{ for } 0 < \epsilon < 1 \quad (72.10)$$

$$\therefore \lim_{n \rightarrow \infty} \Pr(|T_n - 1| \geq \epsilon) = 0, \forall \epsilon > 0 \quad (72.11)$$

$\therefore T_n$  converges to 1 in probability.

b) OPTION-2:

Convergence in Distribution :

A sequence of random variables  $X_1, X_2, X_3, \dots$  converges in distribution to a random variable  $X$ , shown by  $X_n \xrightarrow{d} X$ , if

$$\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x) \quad (72.12)$$

for all  $x$  at which  $F_X(x)$  is continuous.

To evaluate :  $\lim_{n \rightarrow \infty} F_{n(1-T_n)}(x)$

Substituting  $a = -n, b = n$  in (72.5),

$$F_{n(1-T_n)}(x) = 1 - F_{T_n}\left(1 - \frac{x}{n}\right) \quad (72.13)$$

$$F_{T_n}\left(1 - \frac{x}{n}\right) = \begin{cases} \left(1 - \frac{x}{n}\right)^n, & 0 < x < n \\ 1, & x \leq 0 \\ 0, & x \geq n \end{cases} \quad (72.14)$$

$$\therefore \lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x} \quad (72.15)$$

$$\therefore \lim_{n \rightarrow \infty} F_{T_n}\left(1 - \frac{x}{n}\right) = \begin{cases} e^{-x}, & 0 < x < n \\ 1, & x \leq 0 \\ 0, & x \geq n \end{cases} \quad (72.16)$$

$$\therefore F_{n(1-T_n)}(x) = \begin{cases} 1 - e^{-x}, & 0 < x < n \\ 0, & x \leq 0 \\ 1, & x \geq n \end{cases} \quad (72.17)$$

$\therefore n(1 - T_n)$  converges in distribution to a




Fig. 14.8.1: CDF

random variable with CDF in (72.17).

c) OPTION-3:

Convergence in Distribution :

A sequence of random variables  $X_1, X_2, X_3, \dots$  converges in distribution to a random variable  $X$ , shown by  $X_n \xrightarrow{d} X$ ,

if

$$\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x) \quad (72.18)$$

for all  $x$  at which  $F_X(x)$  is continuous.

To evaluate :  $\lim_{n \rightarrow \infty} F_{n^2(1-T_n)}(x)$

Substituting  $a = -n^2, b = n^2$  in (72.5),

$$F_{n^2(1-T_n)}(x) = 1 - F_{T_n}\left(1 - \frac{x}{n^2}\right) \quad (72.19)$$

$$F_{T_n}\left(1 - \frac{x}{n^2}\right) = \begin{cases} \left(1 - \frac{x}{n^2}\right)^n, & 0 < x < n^2 \\ 1, & x \leq 0 \\ 0, & x \geq n^2 \end{cases} \quad (72.20)$$

$$\therefore \lim_{n \rightarrow \infty} \left(1 - \frac{y}{n^2}\right)^n \text{ is not defined} \quad (72.21)$$

$\therefore n^2(1 - T_n)$  does not converge in distribution.

d) OPTION-4:

Convergence in Probability :

A sequence of random variables  $X_1, X_2, X_3, \dots$  converges in probability to a random variable  $X$ , shown by  $X_n \xrightarrow{p} X$ , if

$$\lim_{n \rightarrow \infty} \Pr(|X_n - X| \geq \epsilon) = 0, \forall \epsilon > 0 \quad (72.22)$$

To evaluate :

$$\lim_{n \rightarrow \infty} \Pr(|\sqrt{n}(1 - T_n) - 0| \geq \epsilon), \forall \epsilon > 0$$

$$= \lim_{n \rightarrow \infty} \Pr\left(1 - T_n \geq \frac{\epsilon}{\sqrt{n}}\right) \quad (72.23)$$

$$= \lim_{n \rightarrow \infty} \Pr\left(T_n \leq 1 - \frac{\epsilon}{\sqrt{n}}\right) \quad (72.24)$$

$$= \lim_{n \rightarrow \infty} F_{T_n}\left(1 - \frac{\epsilon}{\sqrt{n}}\right) \quad (72.25)$$

$$F_{T_n}\left(1 - \frac{\epsilon}{\sqrt{n}}\right) = \begin{cases} \left(1 - \frac{\epsilon}{\sqrt{n}}\right)^n, & 0 < \epsilon < \sqrt{n} \\ 0, & \epsilon \geq \sqrt{n} \end{cases} \quad (72.26)$$

$$\therefore \lim_{n \rightarrow \infty} \left(1 - \frac{\epsilon}{\sqrt{n}}\right)^n = 0 \text{ for } 0 < \epsilon < \sqrt{n} \quad (72.27)$$

$$\therefore \lim_{n \rightarrow \infty} \Pr(|\sqrt{n}(1 - T_n) - 0| \geq \epsilon) = 0, \forall \epsilon > 0 \quad (72.28)$$

$\therefore \sqrt{n}(1 - T_n)$  converges to 0 in probability.

Hence, options 1), 2), 4) are correct.

## 15 STATISTICS

15.1. Let  $Y_1$  denote the first order statistic in a random sample of size  $n$  from a distribution that has the pdf

$$f(x) = \begin{cases} e^{-(x-\theta)} & \text{when } \theta < x < \infty \\ 0 & \text{otherwise} \end{cases} \quad (15.1.1)$$

Obtain the distribution of  $Z_n = n(Y_1 - \theta)$ .

**Solution:** From the given information

$$Y_1 = \min\{X_1, X_2, \dots, X_n\} \quad (15.1.2)$$

and

$$F_{Z_n}(z) = \Pr(n(Y_1 - \theta) \leq z) \quad (15.1.3)$$

$$= \Pr\left(Y_1 \leq \frac{z}{n} + \theta\right) \quad (15.1.4)$$

$$= 1 - \Pr\left(Y_1 > \frac{z}{n} + \theta\right) \quad (15.1.5)$$

Let

$$\left(\frac{z}{n} + \theta\right) = z' \quad (15.1.6)$$

Then

$$F_{Z_n}(z) = 1 - \prod_{i=1}^n \Pr(X_i > z') \quad (15.1.7)$$

$$= 1 - (1 - F(z'))^n \quad (15.1.8)$$

$$\Rightarrow F_{Z_n}(z) = 1 - \left(1 - F\left(\frac{z}{n} + \theta\right)\right)^n \quad (15.1.9)$$

where

$$F(x) = \int_{-\infty}^x f(t) dt \quad (15.1.10)$$

$$= \begin{cases} 1 - e^{-(x-\theta)} & \text{when } \theta < x < \infty \\ 0 & \text{otherwise} \end{cases} \quad (15.1.11)$$



Substituting from (15.1.11) in (15.1.9),

$$F_{Z_n}(z) = \begin{cases} 1 - e^{-n(\frac{z}{n} + \theta - \theta)} & \theta < \frac{z}{n} + \theta < \infty \\ 0 & \text{otherwise} \end{cases} \quad (15.1.12)$$

$$= \begin{cases} 1 - e^{-z} & \text{when } 0 < z < \infty \\ 0 & \text{otherwise} \end{cases} \quad (15.1.13)$$

and

$$f_{Z_n}(z) = \frac{d}{dz} F_{Z_n}(z) \quad (15.1.14)$$

$$= \begin{cases} e^{-z} & 0 < z < \infty \\ 0 & \text{otherwise} \end{cases} \quad (15.1.15)$$

The plots for the cdf in (15.1.13) and the pdf in (15.1.15) are shown in Fig. 15.1.1 and Fig. 15.1.2 respectively:




Fig. 15.1.1: cdf of  $Z_n$




Fig. 15.1.2: pdf of  $Z_n$

15.2. Let  $X_1, X_2, \dots, X_n$  be a random sample of size  $n$  ( $\geq 2$ ) from a distribution having the probability density function

$$f(x; \theta) = \begin{cases} \frac{1}{\theta} \exp\left(-\frac{x}{\theta}\right) & x > 0, \\ 0, & \text{otherwise,} \end{cases} \quad (15.2.1)$$

where  $\theta \in (0, \infty)$ . Let  $X_{(1)} = \min\{X_1, X_2, \dots, X_n\}$  and  $T = \sum_{i=1}^n X_i$ . Then  $E(X_{(1)}|T)$  equals

- (A)  $\frac{T}{n^2}$
- (B)  $\frac{T}{n}$
- (C)  $\frac{(n+1)T}{2n}$
- (D)  $\frac{(n+1)^2 T}{4n^2}$

**Solution:**

**Lemma 15.1.** Lehmann–Scheffé theorem : If  $T$  is a complete sufficient statistic for  $\theta$  and

$$E(g(T)) = \tau(\theta) \quad (15.2.2)$$

then  $g(T)$  is the uniformly minimum-variance unbiased estimator (UMVUE) of  $\tau(\theta)$ .

We know that

$$T = \sum_{i=1}^n X_i \quad (15.2.3)$$

is a complete and sufficient statistic. By the law of total expectation,

$$E(E(X_{(1)}|T)) = E(X_{(1)}) \quad (15.2.4)$$

By Lehmann–Scheffé theorem, with

$$\theta = X_{(1)}, \quad (15.2.5)$$

$$\tau(x) = E(x), \quad (15.2.6)$$

$$g(T) = E(X_{(1)}|T). \quad (15.2.7)$$

it follows from (15.2.4) that  $E(X_{(1)}|T)$  is the

UMVUE of  $E(X_{(1)})$ .

$$\Pr(X_{(1)} > x) = \Pr(X_1 > x) \dots \Pr(X_n > x) \quad (15.2.8)$$

$$= (1 - F_{X_1}(x)) \dots (1 - F_{X_n}(x)) \quad (15.2.9)$$

$$= (1 - F_{X_1}(x))^n \quad (15.2.10)$$

$$= \exp\left(-\frac{nx}{\theta}\right) \quad (15.2.11)$$

$$F_{X_{(1)}}(x) = 1 - \exp\left(-\frac{nx}{\theta}\right) \quad (15.2.12)$$

$$f_{X_{(1)}}(x) = \frac{n}{\theta} \exp\left(-\frac{nx}{\theta}\right) \quad (15.2.13)$$

Therefore,  $X_{(1)}$  follows an exponential distribution with mean  $\frac{\theta}{n}$ .

$$E(X_{(1)}) = \frac{\theta}{n} \quad (15.2.14)$$

Note that,

$$E\left(\frac{T}{n^2}\right) = E\left(\frac{\sum_{i=1}^n X_i}{n^2}\right) \quad (15.2.15)$$

$$= \frac{E(\sum_{i=1}^n X_i)}{n^2} \quad (15.2.16)$$

$$= \sum_{i=1}^n \frac{E(X_i)}{n^2} \quad (15.2.17)$$

$$= \sum_{i=1}^n \frac{\theta}{n^2} \quad (15.2.18)$$

$$= \frac{\theta}{n} \quad (15.2.19)$$

$$= E(X_{(1)}) \quad (15.2.20)$$

Therefore, by Lehmann–Scheffé theorem, with

$$\theta = X_{(1)}, \quad (15.2.21)$$

$$\tau(x) = E(x), \quad (15.2.22)$$

$$g(T) = \frac{T}{n^2}, \quad (15.2.23)$$

it follows that  $\frac{T}{n^2}$  is UMVUE of  $E(X_{(1)})$ .

Since there exists a unique UMVUE for  $E(X_{(1)})$ , it follows that

$$E(X_{(1)}|T) = \frac{T}{n^2} \quad (15.2.24)$$

Hence, option A is correct.

15.3. Suppose  $n$  units are drawn from a population




Fig. 15.2.1: Theory vs Simulated plot of  $E(X_{(1)}|T)$

of  $N$  units sequentially as follows. A random sample

$$U_1, U_2, \dots, U_N \text{ of size } N, \text{ drawn from } U(0, 1) \quad (15.3.1)$$

The  $k$ -th population unit is selected if

$$U_k < \frac{n - n_k}{N - k + 1}, k = 1, 2, \dots, N. \text{ where, } n_1 = 0, n_k = \quad (15.3.2)$$

number of units selected out of first  $k-1$  units for each  $k = 2, 3, \dots, N$ . Then,

a) The probability of inclusion of the second unit in the sample

$$\text{is } \frac{n}{N} \quad (15.3.3)$$

b) The probability of inclusion of the first and the second unit in the sample

$$\text{is } \frac{n(n-1)}{N(N-1)} \quad (15.3.4)$$

c) The probability of not including the first and including the second unit in the sample

$$\text{is } \frac{n(N-n)}{N(N-1)} \quad (15.3.5)$$

d) The probability of including the first and not including the second unit in the sample

$$\text{is } \frac{n(n-1)}{N(N-1)} \quad (15.3.6)$$

**Solution:**

Defining random variable  $X \in \{0, 1, 2, \dots, N\}$   
(15.3.7)

Where,  $X = i$  when  $i$ th unit is included.  
(15.3.8)

The first unit in the sample is included if

$$U_1 < \frac{n - n_1}{N - 1 + 1} \quad (15.3.9)$$

Here,  $n_1 = 0$  is given in the qn. (15.3.10)

$$\therefore \Pr(X = 1) = \frac{n}{N} \quad (15.3.11)$$

a) For  $k=2$ ,

$n_2 = 1$  when, first unit is included.  
(15.3.12)

$$U_2 < \frac{n - n_2}{N - 2 + 1} \left( = \frac{n - 1}{N - 1} \right) \quad (15.3.13)$$

$$\therefore \Pr(X = 2 | X = 1) = \frac{n - 1}{N - 1} \quad (15.3.14)$$

$\Pr(X = 1, X = 2)$

$$= \Pr(X = 2 | X = 1) \times \Pr(X = 1) \quad (15.3.15)$$

$$\therefore \Pr(X = 1, X = 2) = \frac{n(n - 1)}{N(N - 1)} \quad (15.3.16)$$

$n_2 = 0$  when, first unit is not included.  
(15.3.17)

$$U_2 < \frac{n - n_2}{N - 2 + 1} \left( = \frac{n}{N - 1} \right) \quad (15.3.18)$$

$$\therefore \Pr(X = 2 | X \neq 1) = \frac{n}{N - 1} \quad (15.3.19)$$

$\Pr(X \neq 1, X = 2)$

$$= \Pr(X = 2 | X \neq 1) \times \Pr(X \neq 1) \quad (15.3.20)$$

$$\therefore \Pr(X \neq 1, X = 2) = \left(1 - \frac{n}{N}\right) \times \frac{n}{N - 1} \quad (15.3.21)$$

$$\therefore \Pr(X \neq 1, X = 2) = \frac{n(N - n)}{N(N - 1)} \quad (15.3.22)$$

From (15.3.16) and (15.3.22)

$$\Pr(X = 2) = \frac{n(n - 1)}{N(N - 1)} + \frac{n(N - n)}{N(N - 1)} = \frac{n}{N} \quad (15.3.23)$$

Hence, option 1 is correct.

b) From (15.3.16)

$$\Pr(X = 1, X = 2) = \frac{n(n - 1)}{N(N - 1)} \quad (15.3.24)$$

Hence, option 2 is correct.

c) From (15.3.22)

$$\Pr(X \neq 1, X = 2) = \frac{n(N - n)}{N(N - 1)} \quad (15.3.25)$$

Hence, option 3 is correct.

d)

$$\Pr(X = 1, X \neq 2) = \frac{n}{N} \times \left(1 - \frac{n}{N}\right) = \frac{n(N - n)}{N^2} \quad (15.3.26)$$

Hence, option 4 is incorrect.

Therefore, Options 1, 2, 3 are correct

15.4. Let  $X_1, X_2, X_3, \dots, X_n$  be independent random variables follow a common continuous distribution  $F$ , which is symmetric about 0. For  $i=1, 2, 3, \dots, n$ , define

$$S_i = \begin{cases} 1 & \text{if } X_i > 0 \\ -1 & \text{if } X_i < 0 \\ 0 & \text{if } X_i = 0 \end{cases} \quad \text{and} \quad (1.1)$$

$R_i$  = rank of  $|X_i|$  in the set  $\{|X_1|, |X_2|, \dots, |X_n|\}$ . Which of the following statements are correct?

- a)  $S_1, S_2, \dots, S_n$  are independent and identically distributed.
- b)  $R_1, R_2, \dots, R_n$  are independent and identically distributed.
- c)  $S = (S_1, S_2, \dots, S_n)$  and  $R = (R_1, R_2, \dots, R_n)$  are independent.

**Solution:**

A sequence  $\{X_i\}$  is an Independent and identical if and only if  $F_{X_n}(x) = F_{X_k}(x) \forall n, k, x$  and any subset of terms of the sequence is a set of mutually independent random variables. Where  $F$  is the probability density function.

- a) As the probability distribution function of  $\{X_i\}$  is symmetric about origin we can say

that

$$F_{X_i}(-x) = F_{X_i}(x) \forall x \in R \quad (2.1)$$

and the mean of the distribution( $\mu$ )

$$\mu = 0 \quad (2.2)$$

The sequence  $S_i$  depend on  $X_i$  as mention in 1.1, as each  $S_i$  depend only on  $X_i$  we can say that sequence  $S_i$  is independent.

$$\Pr(S_1 = 1, S_2 = 1, \dots, S_n = 1) = \prod_{i=1}^n \Pr(S_i = 1) \quad (2.3)$$

Any subset of terms of sequence  $\{S_i\}$  is a set of mutually independent random variables and its distribution is identical.

$$F_{S_n}(s) = F_{S_k}(s) \quad \forall s, k, n \quad (2.4)$$

So, the sequence  $\{S_i\}$  is independent and identical.

- b) **Ranking** refers to the data transformation in which the numerical or ordinary values are replaced by the rank of numerical value when compared to a list of other values. Usually we follow increasing order for ranking.

Ranking of a sequence depend on every elements of the sequence. Let  $\{R_i\}$  be the output sequence of the ranking function of  $\{|X_i|\}$ .

$$R_k = \text{rank of } |X_k| \text{ in the set } \{|X_1|, |X_2|, \dots, |X_n|\} \quad (2.5)$$

As  $R_k$  depend not only on  $|X_k|$  but on the rest of the elements of the set  $\{|X_1|, |X_2|, \dots, |X_n|\}$ . So the sequence  $R_i$  is not independent. Hence  $R_i$  is not an independent and identical distribution.

- c) As the  $i^{th}$  element of sequence R depends only on set  $\{|X_1|, |X_2|, \dots, |X_n|\}$ , we can say that sequence S and R are independent.

Answer: A, C

- 15.5. A simple random variable of size n will be drawn from a class of 125 students, and the mean mathematics score of the sample will be computed, If the standard error of the sample mean for "with replacement sampling" is twice as much as the standard error of the sample mean for "without replacement sampling", the value of n is ?

- a) 32

- b) 63

- c) 79

- d) 94

**Solution:** Let N be the population size so, N=120. The given sample size is n. **Notations**

: y : student under consideration.  $y_i$  : Maths marks of  $i^{th}$  student in the sample. Y : student of class.  $Y_i$  : Maths marks of  $i^{th}$  student in the class.  $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$  : Average of sample

class.  $\bar{Y} = \frac{1}{N} \sum_{i=1}^N Y_i$  : Average of whole class.

$S^2 = \frac{1}{N-1} \sum_{i=1}^N (Y_i - \bar{Y})^2$  : S=Std dev of the class.  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (Y_i - \bar{Y})^2$  : Variance of the class. Standard error of sample mean  $SE_{mean} = \frac{s}{\sqrt{n}}$ .

Where

$s$  = standard deviation of sample mean.

$n$  = sample class size.

**Variance of the  $\bar{y}$**

$$V(\bar{y}) = E(\bar{y} - \bar{Y})^2 \quad (15.5.1)$$

$$= E \left[ \frac{1}{n} \sum_{i=1}^n (y_i - \bar{Y}) \right]^2 \quad (15.5.2)$$

$$= E \left[ \frac{1}{n^2} \sum_{i=1}^n (y_i - \bar{Y})^2 + \frac{1}{n^2} \sum_{1 \leq i \neq j \leq n} (y_i - \bar{Y})(y_j - \bar{Y}) \right] \quad (15.5.3)$$

$$= \frac{1}{n^2} \sum_{i=1}^n E(y_i - \bar{Y})^2 + \frac{1}{n^2} \sum_{1 \leq i \neq j \leq n} E(y_i - \bar{Y})(y_j - \bar{Y}) \quad (15.5.4)$$

$$\text{Let } K = \sum_{1 \leq i \neq j \leq n} E(y_i - \bar{Y})(y_j - \bar{Y}) \quad (15.5.5)$$

$$= \frac{1}{n^2} \sum_{i=1}^n \sigma^2 + \frac{K}{n^2} \quad (15.5.6)$$

$$= \frac{1}{n^2} n \sigma^2 + \frac{K}{n^2} \quad (15.5.7)$$

$$= \frac{N-1}{Nn} S^2 + \frac{K}{n^2} \quad (15.5.8)$$

Finding the value of K in case of Simple random sampling with repetition (SR-SWR) and Simple random sampling without repetition (SRSWOR) allows us to calculate the variance of mean. **K value in case of SR-**

**SWOR**

$$K = \sum_{1 \leq i \neq j \leq n} E(y_i - \bar{Y})(y_j - \bar{Y})$$

Consider

$$E(y_i - \bar{Y})(y_j - \bar{Y}) = \frac{1}{N(N-1)} \sum_{1 \leq k \neq l \leq n} E(y_k - \bar{Y})(y_l - \bar{Y})$$

Since

$$\left[ \sum_{k=1}^N (y_k - \bar{Y}) \right]^2 = \sum_{i=1}^N (y_i - \bar{Y})^2 + \sum_{1 \leq k \neq l \leq n} E(y_k - \bar{Y})(y_l - \bar{Y})$$

$$\Rightarrow 0 = (N-1)S^2 + \sum_{1 \leq k \neq l \leq n} E(y_k - \bar{Y})(y_l - \bar{Y})$$

$$\Rightarrow E(y_i - \bar{Y})(y_j - \bar{Y}) = \frac{1}{N(N-1)}(N-1)(-S^2)$$

$$\Rightarrow K = n(n-1) \frac{(-S^2)}{N}$$

Putting this value in (15.5.8) gives us

$$V(\bar{y})_{WOR} = \frac{N-1}{Nn} S^2 + \frac{n-1(-S^2)}{Nn} \quad (15.5.9)$$

$$= \frac{N-n}{Nn} S^2 \quad (15.5.10)$$

**K value in case of SRSWR**

$$K = \sum_{1 \leq i \neq j \leq n} E(y_i - \bar{Y})(y_j - \bar{Y})$$

Since we are selecting the samples with replacements choosing  $i^{th}$  and  $j^{th}$  sample is independent of each other. So,

$$K = \sum_{1 \leq i \neq j \leq n} E(y_i - \bar{Y})E(y_j - \bar{Y})$$

$$= 0$$

(Since deviation about mean is 0)

Putting  $K=0$  in (15.5.8) we get

$$V(\bar{y})_{WR} = \frac{N-1}{Nn} S^2 \quad (15.5.11)$$

From equation (15.5.10) standard error of mean

of sample class without repetition

$$SE_{WOR} = \frac{s}{\sqrt{n}} \quad (15.5.12)$$

$$= \sqrt{\frac{V(\bar{y})_{WOR}}{n}} \quad (15.5.13)$$

$$= \sqrt{\frac{N-n}{Nn^2}} S \quad (15.5.14)$$

From equation (15.5.11) standard error of mean of sample class with repetition

$$SE_{WR} = \sqrt{\frac{V(\bar{y})_{WR}}{n}} \quad (15.5.15)$$

$$= \sqrt{\frac{N-1}{Nn^2}} S \quad (15.5.16)$$

Given to find the value of n if  $2 \times SE_{WOR} = SE_{WR}$ . From (15.5.14) and (15.5.16) we can write

$$2 \sqrt{\frac{N-n}{Nn^2}} S = \sqrt{\frac{N-1}{Nn^2}} S \quad (15.5.17)$$

$$\Rightarrow 4(N-n) = N-1 \quad (15.5.18)$$

$$\Rightarrow 4N + 1 - N = 4n \quad (15.5.19)$$

$$\Rightarrow 4n = 3(125) + 1 \quad (15.5.20)$$

$$\Rightarrow n = 94 \quad (15.5.21)$$

Therefore the sample size for the given condition to be met is  $n=94$ . **(Option D)**

15.6. Let  $X_1$  and  $X_2$  be i.i.d. with probability mass function  $f_\theta(x) = \theta^x (1-\theta)^{1-x}$ ;  $x = 0, 1$  where  $\theta \in (0, 1)$ . Which of the following statements are true?

- a)  $X_1 + 2X_2$  is a sufficient statistic
- b)  $X_1 - X_2$  is a sufficient statistic
- c)  $X_1^2 + X_2^2$  is a sufficient statistic
- d)  $X_1^2 + X_2$  is a sufficient statistic

**Solution:** Given that,  $X_1$  and  $X_2$  are i.i.d. with probability mass function

$$f(x) = \begin{cases} (1-\theta) & x = 0 \\ \theta & x = 1 \end{cases} \quad (15.6.1)$$

A statistic  $t = T(X)$  is sufficient for a parameter  $\theta$  if the conditional probability distribution of the data, given the statistic  $t = T(X)$  does not depend on the parameter  $\theta$ . i.e.,

$$P_\theta(X_1 = x_1, X_2 = x_2 | T = t) \quad (15.6.2)$$

is independent of  $\theta$  for all  $x_1, x_2$  and  $t$

a) Let  $T = X_1 + 2X_2$

Consider a case where  $x_1 = 0, x_2 = 0$  and  $t = 0$

$$\Pr(T = 0) = \Pr(X_1 + 2X_2 = 0) \quad (15.6.3)$$

$$= \Pr(X_1 = 0, X_2 = 0) \quad (15.6.4)$$

As  $X_1$  and  $X_2$  are independent

$$\begin{aligned} \Pr(T = 0) &= \Pr(X_1 = 0) \Pr(X_2 = 0) \\ &= (1 - \theta)^2 \quad (15.6.5) \end{aligned}$$

The conditional probability,

$$\begin{aligned} \Pr(X_1 = 0, X_2 = 0 | T = 0) \\ = \frac{\Pr((X_1 = 0, X_2 = 0) \cap (T = 0))}{\Pr(T = 0)} \quad (15.6.6) \end{aligned}$$

From (15.6.4),  $(X_1 = 0, X_2 = 0) \subseteq (T = 0)$

$$= \frac{\Pr(X_1 = 0, X_2 = 0)}{\Pr(T = 0)} = \frac{(1 - \theta)^2}{(1 - \theta)^2} = 1 \quad (15.6.7)$$

Similarly, conditional probabilities for other values of  $x_1, x_2$  and  $t$  are given in table 15.6.1

$x_1$	$x_2$	$t$ $t = X_1 + 2X_2$	Conditional probability $P_\theta(X_1 = x_1, X_2 = x_2   T = t)$
0	0	0 otherwise	1 0
1	0	1 otherwise	1 0
0	1	2 otherwise	1 0
1	1	3 otherwise	1 0

TABLE 15.6.1: Conditional Probabilities

From table 15.6.1, all the conditional probabilities are independent of  $\theta$

$\therefore X_1 + 2X_2$  is a sufficient statistic.

b) Let  $T = X_1 - X_2$

Consider a case where  $x_1 = 0, x_2 = 0$  and  $t = 0$

$$\begin{aligned} \Pr(T = 0) &= \Pr(X_1 - X_2 = 0) \\ &= \Pr(X_1 = 0, X_2 = 0) + \Pr(X_1 = 1, X_2 = 1) \quad (15.6.8) \end{aligned}$$

As  $X_1$  and  $X_2$  are independent

$$\begin{aligned} &= \Pr(X_1 = 0) \Pr(X_2 = 0) \\ &+ \Pr(X_1 = 1) \Pr(X_2 = 1) = (1 - \theta)^2 + \theta^2 \quad (15.6.9) \end{aligned}$$

The conditional probability,

$$\begin{aligned} \Pr(X_1 = 0, X_2 = 0 | T = 0) \\ = \frac{\Pr((X_1 = 0, X_2 = 0) \cap (T = 0))}{\Pr(T = 0)} \quad (15.6.10) \end{aligned}$$

From (15.6.8),  $(X_1 = 0, X_2 = 0) \subseteq (T = 0)$

$$= \frac{\Pr(X_1 = 0, X_2 = 0)}{\Pr(T = 0)} = \frac{(1 - \theta)^2}{(1 - \theta)^2 + \theta^2} \quad (15.6.11)$$

depends on  $\theta$ .

$\therefore X_1 - X_2$  is not a sufficient statistic.

c) Let  $T = X_1^2 + X_2^2$

Consider a case where  $x_1 = 1, x_2 = 0$  and  $t = 1$

$$\begin{aligned} \Pr(T = 1) &= \Pr(X_1^2 + X_2^2 = 1) \\ &= \Pr(X_1 = 1, X_2 = 0) + \Pr(X_1 = 0, X_2 = 1) \\ &= \theta(1 - \theta) + (1 - \theta)\theta = 2\theta(1 - \theta) \quad (15.6.12) \end{aligned}$$

The conditional probability,

$$\begin{aligned} \Pr(X_1 = 1, X_2 = 0 | T = 1) \\ = \frac{\Pr((X_1 = 1, X_2 = 0) \cap (T = 1))}{\Pr(T = 1)} \quad (15.6.13) \end{aligned}$$

From (15.6.12),  $(X_1 = 1, X_2 = 0) \subseteq (T = 1)$

$$= \frac{\Pr(X_1 = 1, X_2 = 0)}{\Pr(T = 1)} = \frac{\theta(1 - \theta)}{2\theta(1 - \theta)} = \frac{1}{2} \quad (15.6.14)$$

Similarly, conditional probabilities for other values of  $x_1, x_2$  and  $t$  are given in table 15.6.2

From table 15.6.2, all the conditional probabilities are independent of  $\theta$

$\therefore X_1^2 + X_2^2$  is a sufficient statistic.

d) Let  $T = X_1^2 + X_2$

Consider a case where  $x_1 = 1, x_2 = 0$  and

$x_1$	$x_2$	$t$ $t = X_1^2 + X_2^2$	Conditional probability $P_\theta(X_1 = x_1, X_2 = x_2   T = t)$
0	0	0 otherwise	1 0
1	0	1 otherwise	$\frac{1}{2}$ 0
0	1	1 otherwise	$\frac{1}{2}$ 0
1	1	2 otherwise	1 0

TABLE 15.6.2: Conditional Probabilities

$t = 1$

$$\begin{aligned}
 \Pr(T = 1) &= \Pr(X_1^2 + X_2^2 = 1) \\
 &= \Pr(X_1 = 1, X_2 = 0) + \Pr(X_1 = 0, X_2 = 1) \\
 &= \theta(1 - \theta) + (1 - \theta)\theta = 2\theta(1 - \theta)
 \end{aligned} \tag{15.6.15}$$

The conditional probability,

$$\begin{aligned}
 \Pr(X_1 = 1, X_2 = 0 | T = 1) \\
 &= \frac{\Pr((X_1 = 1, X_2 = 0) \cap (T = 1))}{\Pr(T = 1)}
 \end{aligned} \tag{15.6.16}$$

From (15.6.15),  $(X_1 = 1, X_2 = 0) \subseteq (T = 1)$

$$= \frac{\Pr(X_1 = 1, X_2 = 0)}{\Pr(T = 1)} = \frac{\theta(1 - \theta)}{2\theta(1 - \theta)} = \frac{1}{2} \tag{15.6.17}$$

Similarly, conditional probabilities for other values of  $x_1, x_2$  and  $t$  are given in table 15.6.3

$x_1$	$x_2$	$t$ $t = X_1^2 + X_2^2$	Conditional probability $P_\theta(X_1 = x_1, X_2 = x_2   T = t)$
0	0	0 otherwise	1 0
1	0	1 otherwise	$\frac{1}{2}$ 0
0	1	1 otherwise	$\frac{1}{2}$ 0
1	1	2 otherwise	1 0

TABLE 15.6.3: Conditional Probabilities

From table 15.6.3, all the conditional prob-

abilities are independent of  $\theta$

$\therefore X_1^2 + X_2^2$  is a sufficient statistic.

Answer : Options 1,3,4

15.7. Let  $X_1$  and  $X_2$  be a random sample of size two from a distribution with probability density function

$$f_\theta(x) = \theta \left( \frac{1}{\sqrt{2\pi}} \right) e^{-\frac{1}{2}x^2} + (1 - \theta) \left( \frac{1}{2} \right) e^{-|x|},$$

$-\infty < x < \infty$ ,

where  $\theta \in \left\{0, \frac{1}{2}, 1\right\}$ . If the observed values of  $X_1$  and  $X_2$  are 0 and 2, respectively, then the maximum likelihood estimate of  $\theta$  is

- a) 0
- b)  $\frac{1}{2}$
- c) 1
- d) not unique

**Solution:** Given  $X_1 = 0, X_2 = 2, n = 2$  and

$$f_\theta(x) = \theta \left( \frac{1}{\sqrt{2\pi}} \right) e^{-\frac{1}{2}x^2} + (1 - \theta) \left( \frac{1}{2} \right) e^{-|x|} \tag{15.7.1}$$

Then log of likelihood function is given by

$$l(\theta) = \sum_{i=1}^{i=n} \log f_\theta(x_i) \tag{15.7.2}$$

$$= \log f_\theta(x_1) + \log f_\theta(x_2) \tag{15.7.3}$$

$$\begin{aligned}
 &= \log \left[ \theta \left( \frac{1}{\sqrt{2\pi}} \right) e^{-\frac{1}{2}0^2} + (1 - \theta) \left( \frac{1}{2} \right) e^{-|0|} \right] \\
 &\quad + \log \left[ \theta \left( \frac{1}{\sqrt{2\pi}} \right) e^{-\frac{1}{2}2^2} + (1 - \theta) \left( \frac{1}{2} \right) e^{-|2|} \right]
 \end{aligned} \tag{15.7.4}$$

$$\begin{aligned}
 &= \log \left( \theta \left( \frac{1}{\sqrt{2\pi}} \right) + (1 - \theta) \left( \frac{1}{2} \right) \right) \\
 &\quad + \log \left( \theta \left( \frac{1}{\sqrt{2\pi}} \right) e^{-2} + (1 - \theta) \left( \frac{1}{2} \right) e^{-2} \right)
 \end{aligned} \tag{15.7.5}$$

$$= 2 \log \left( \theta \left( \frac{1}{\sqrt{2\pi}} \right) + (1 - \theta) \left( \frac{1}{2} \right) \right) - 2 \tag{15.7.6}$$

Since likelihood  $L(\theta) = e^{l(\theta)}$ .

Likelihood function  $L(\theta)$  at  $\theta = 0, \frac{1}{2}, 1$  is

given by

a) At  $\theta = 0$   $L(\theta = 0) = \frac{1}{4}e^{-2} = 0.0338$

b) At  $\theta = 1$   $L(\theta = 1) = \frac{1}{2\pi}e^{-2} = 0.0215$

c) At  $\theta = \frac{1}{2}$   $L(\theta = \frac{1}{2}) = \left(\frac{1}{2\sqrt{2\pi}} + \frac{1}{4}\right)^2 e^{-2} = 0.0273$

Hence the maximum likelihood estimate of  $\theta$  is at  $\theta = 0$

15.8. Let  $\{X_n, n \geq 1\}$  be i.i.d. uniform  $(-1, 2)$  random variables. Which of the following statements are true?

a)  $\frac{1}{n} \sum_{i=1}^n X_i \rightarrow 0$  almost surely

b)  $\left\{ \frac{1}{2n} \sum_{i=1}^n X_{2i} - \frac{1}{2n} \sum_{i=1}^n X_{2i-1} \right\} \rightarrow 0$  almost surely

c)  $\sup \{X_1, X_2, \dots\} = 2$  almost surely

d)  $\inf \{X_1, X_2, \dots\} = -1$  almost surely

**Solution:** We using convergence in almost surely and Strong law of large number (SLLN)

a) *Almost sure convergence* : Let  $X_1, X_2, \dots$  be an infinite sequence of random variables. We shall say that the sequence  $\{X_i\}$  converges with probability 1 (or converges almost surely (a.s.)) to a random variable  $Y$ , if

$$\Pr\left(\lim_{n \rightarrow \infty} X_n = Y\right) = 1 \quad (15.8.1)$$

$$\text{and we write, } X_n \xrightarrow{\text{a.s.}} Y \quad (15.8.2)$$

b) *SLLN* : Let  $X_n$  be i.i.d with  $\mathbf{E}[|X_1|] < \infty$ . Then, as  $n \rightarrow \infty$ , we have

$$\frac{S_n}{n} \xrightarrow{\text{a.s.}} \mathbf{E}[X_1] \implies \frac{S_n}{n} \xrightarrow{\text{P}} \mathbf{E}[X_1] \quad (15.8.3)$$

$$\text{, where } S_n = X_1 + \dots + X_n \quad (15.8.4)$$

also,

$$X_i \xrightarrow{\text{a.s.}} X \implies g(X_i) \xrightarrow{\text{a.s.}} g(X) \quad (15.8.5)$$

a)

$$\frac{1}{n} (X_1 + \dots + X_n) \rightarrow E(X) \in (-1, 2) \quad (15.8.6)$$

$$\text{as } n \rightarrow \infty,$$

$$(15.8.7)$$

according to strong law of large numbers (SLLN).

So, option (A) is incorrect.

b) using this 15.8.5, we solve as

$$\left\{ \frac{1}{2n} \sum_{i=1}^n X_{2i} - \frac{1}{2n} \sum_{i=1}^n X_{2i-1} \right\} \xrightarrow{\text{a.s.}} \left\{ \frac{nX}{2n} - \frac{nX}{2n} \right\} \quad (15.8.8)$$

$$= 0 \quad (15.8.9)$$

option (B) is correct.

c) Similarly, Let  $M = \sup(S)$ . Then,

$$x \leq M, \quad \forall x \in S \quad (15.8.10)$$

$$\forall \epsilon > 0, \quad (M - \epsilon, M] \cap S \neq \emptyset \quad (15.8.11)$$

where,  $S$  be a nonempty subset of  $\mathbb{R}$  with an upper bound. Using  $X_i \xrightarrow{\text{a.s.}} X$  this, we conclude that

$$\sup \{X_1, X_2, \dots\} = 2 \text{ almost surely} \quad (15.8.12)$$

d) Let  $m = \inf(S)$ . Then

$$x \geq m, \quad \forall x \in S \quad (15.8.13)$$

$$\forall \epsilon > 0, \quad [m, m + \epsilon] \cap S \neq \emptyset \quad (15.8.14)$$

where,  $S$  be a nonempty subset of  $\mathbb{R}$  with a lower bound. Again using  $X_i \xrightarrow{\text{a.s.}} X$  this, we conclude that

$$\inf \{X_1, X_2, \dots\} = -1 \text{ almost surely} \quad (15.8.15)$$

Hence (B), (C) and (D) are correct options.

15.9.  $X_1, X_2, \dots$  are independent identically distributed random variables having common density  $f$ . Assume  $f(x) = f(-x)$  for all  $x \in \mathbb{R}$ . Which of the following statements is correct?

a)  $\frac{1}{n} (X_1 + \dots + X_n) \rightarrow 0$  in probability

b)  $\frac{1}{n} (X_1 + \dots + X_n) \rightarrow 0$  almost surely

c)  $\Pr\left(\frac{1}{\sqrt{n}} (X_1 + \dots + X_n) < 0\right) \rightarrow \frac{1}{2}$

d)  $\sum_{i=1}^n X_i$  has the same distribution as  $\sum_{i=1}^n (-1)^i X_i$

**Solution:** We using

a) (1) *Convergence in probability* : Let  $X_1, X_2, \dots$  be an infinite sequence of random variables, and let  $Y$  be another random variable. Then the sequence



$\{X_n\}$  converges in probability to  $Y$ , if

$$\forall \epsilon > 0, \lim_{n \rightarrow \infty} \Pr(|X_n - Y| \geq \epsilon) = 0, \quad (15.9.1)$$

and we write

$$X_n \xrightarrow{P} Y. \quad (15.9.2)$$

- (2) *Convergence in almost surely* : Let  $X_1, X_2, \dots$  be an infinite sequence of random variables. We shall say that the sequence  $\{X_i\}$  converges with probability 1 (or converges almost surely (a.s.)) to a random variable  $Y$ , if

$$\Pr\left(\lim_{n \rightarrow \infty} X_n = Y\right) = 1 \quad (15.9.3)$$

and we write

$$X_n \xrightarrow{a.s.} Y \quad (15.9.4)$$

- (3) *Strong law of large number (SLLN)* : Let  $X_1, X_2, \dots$  be an infinite sequence of random variables, If  $\mathbf{E}[|X_1|] < \infty$ . Then, as  $n \rightarrow \infty$ , we have

$$\frac{S_n}{n} \xrightarrow{a.s.} \mathbf{E}[X_1] \implies \frac{S_n}{n} \xrightarrow{P} \mathbf{E}[X_1], \quad (15.9.5)$$

$$\text{where, } S_n = X_1 + \dots + X_n \quad (15.9.6)$$

using SLLN, (B) are incorrect option.

- b) *Relation between in probability and almost surely* : Let  $Z, Z_1, Z_2, \dots$  be random variables. Suppose  $Z_n \rightarrow Z$  with probability 1. Then, we say

$$Z_n \xrightarrow{a.s.} Z \implies Z_n \xrightarrow{P} Z. \quad (15.9.7)$$

(15.9.5), also in probability also hold this equation. Hence (A) is incorrect option.

- c) *Central Limit Theorem* : Let  $X_1, X_2, \dots$  be i.i.d. with finite mean  $\mu$  and finite variance  $\sigma^2$ . Let  $Z \sim N(0, 1)$ . Set  $S_n = X_1 + \dots + X_n$ , and

$$Z_n = \frac{S_n - n\mu}{\sqrt{n\sigma^2}} \quad (15.9.8)$$

Then as  $n \rightarrow \infty$ , the sequence  $\{Z_n\}$  converges in distribution to the  $Z$ , i.e.,  $Z_n \xrightarrow{D} Z$ .

Consider,

$$Y = \frac{X_1 + X_2 + \dots + X_n}{\sqrt{n}} \quad (15.9.9)$$

So,

$$E(Y) = E\left(\frac{X_1 + X_2 + \dots + X_n}{\sqrt{n}}\right) = 0 \quad (15.9.10)$$

$$V(Y) = V\left(\frac{X_1 + X_2 + \dots + X_n}{\sqrt{n}}\right) = \frac{1}{n} 2n = 2 \quad (15.9.11)$$

$$Y \sim N[0, 2] \quad (15.9.12)$$

we know that,

$$f(x) = f(-x) \implies \text{Symmetry about Zero,} \quad (15.9.13)$$

So,

$$\Pr(Y < 0) = \frac{1}{2} \quad (15.9.14)$$

$$\Pr\left(\frac{1}{\sqrt{n}}(X_1 + \dots + X_n) < 0\right) = \frac{1}{2} \quad (15.9.15)$$

Hence, (C) is incorrect option.

- d) *Characteristic function* : For a scalar random variable  $X$  the characteristic function is defined as the expected value of  $e^{itx}$ , where  $i$  is the imaginary unit, and  $t \in \mathbf{R}$  is the argument of the characteristic function:

$$\begin{cases} \varphi_X : \mathbf{R} \rightarrow \mathbf{C} \\ \varphi_X(t) = \mathbf{E}[e^{itX}] = \int_{\mathbf{R}} e^{itx} dF_X(x) \\ = \int_{\mathbf{R}} e^{itx} f_X(x) dx = \int_0^1 e^{itQ_X(p)} dp \end{cases} \quad (15.9.16)$$

Here  $F_X$  is the cumulative distribution function of  $X$ , Consider,  $\phi_x(t)$  is characteristic function of  $X_i, i = 1, \dots, n$ .

$$f(x) = f(-x) \implies \phi_x(t) = \phi_{-x}(t) \quad (15.9.17)$$

Therefore,

$$\phi_{\sum_{i=1}^n X_i}(t) = \phi_{X_1 + \dots + X_n}(t) = \phi_{X_1}(t) \cdots \phi_{X_n}(t) \quad (15.9.18)$$

$$= [\phi_x(t)]^n \quad (15.9.19)$$

similarly,

$$\phi_{\sum_{i=1}^n (-1)^i X_i}(t) = \phi_{-X_1} + \phi_{X_2} + \dots + \phi_{(-1)^n X_n}(t) \quad (15.9.20)$$

$$= \phi_{-X_1}(t) \cdot \phi_{X_2}(t) \cdots \phi_{(-1)^n X_n}(t) \quad (15.9.21)$$

$$= [\phi_X(t)]^n \quad (15.9.22)$$

$$\phi_{\sum_{i=1}^n X_i}(t) = \phi_{\sum_{i=1}^n (-1)^i X_i}(t) \quad (15.9.23)$$

$\therefore \sum_{i=1}^n X_i$  has same distribution as  $\sum_{i=1}^n (-1)^i X_i$ .

Hence, only (D) is correct option.

15.10.  $X_1, X_2, \dots, X_n$  are independent and identically distributed as  $N(\mu, \sigma^2)$ ,  $-\infty < \mu < \infty$ ,  $\sigma^2 > 0$ . Then

a)  $\sum_1^n \frac{(X_i - \bar{X})^2}{n-1}$  is the Minimum Variance Unbiased Estimate of  $\sigma^2$

b)  $\sqrt{\sum_1^n \frac{(X_i - \bar{X})^2}{n-1}}$  is the Minimum Variance Unbiased Estimate of  $\sigma$

c)  $\sum_1^n \frac{(X_i - \bar{X})^2}{n}$  is the Maximum Likelihood Estimate of  $\sigma^2$

d)  $\sqrt{\sum_1^n \frac{(X_i - \bar{X})^2}{n}}$  is the Maximum Likelihood Estimate of  $\sigma$

**Solution:** The pdf for each random variable is same as they are all identical and independent Normal Distributions with same  $\mu$  and  $\sigma^2$ .

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{(x - \mu)^2}{2\sigma^2} \quad (15.10.1)$$

Let us take our maximum likelihood function for given random variable  $X_i$

$$L(\mu; \sigma | X_i) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{(X_i - \mu)^2}{2\sigma^2} \quad (15.10.2)$$

Since all the random variables are i.i.d

$$L(\mu; \sigma | X_1, X_2, \dots, X_n) = \prod_{i=1}^n L(\mu; \sigma | X_i) \quad (15.10.3)$$

Let us denote:

$$L_m : L(\mu; \sigma | X_1, X_2, \dots, X_n) \quad (15.10.4)$$

Substituting (15.10.2) for each Random Vari-

able in (15.10.3)

$$L_m = \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{(X_i - \mu)^2}{2\sigma^2} \quad (15.10.5)$$

Taking natural log on both sides and simplifying

$$\ln L_m = \frac{-n}{2} \ln 2\pi - n \ln \sigma - \sum_{i=1}^n \frac{(X_i - \mu)^2}{2\sigma^2} \quad (15.10.6)$$

In order to find Maximum Likelihood we need to maximise  $\mu$  and  $\sigma$  w.r.t. all Random variables. Taking partial derivative w.r.t  $\mu$  and taking  $\sigma$  as constant

$$\frac{\partial \ln L_m}{\partial \mu} = \sum_{i=1}^n \frac{(X_i - \mu)}{\sigma^2} \quad (15.10.7)$$

The value for  $\mu$  at which  $L_m$  achieves maximum value is same in  $\ln L_m$

$$\therefore \frac{\partial \ln L_m}{\partial \mu} = 0 \quad (15.10.8)$$

$$\therefore \sum_{i=1}^n \frac{(X_i - \mu)}{\sigma^2} = 0 \quad (15.10.9)$$

On simplifying the expression we get:

$$n\mu = \sum_{i=1}^n X_i \quad (15.10.10)$$

$$\mu = \frac{1}{n} \sum_{i=1}^n X_i \quad (15.10.11)$$

Let us denote the value achieved in (15.10.11) as  $\bar{X}$ . Taking partial derivative w.r.t  $\sigma$  and taking  $\mu$  as constant

$$\frac{\partial \ln L_m}{\partial \sigma} = \frac{-n}{\sigma} + \sum_{i=1}^n \frac{(X_i - \mu)^2}{\sigma^3} \quad (15.10.12)$$

The value for  $\sigma$  at which  $L_m$  achieves maximum value is same in  $\ln L_m$

$$\frac{\partial \ln L_m}{\partial \sigma} = 0 \quad (15.10.13)$$

$$\frac{-n}{\sigma} + \sum_{i=1}^n \frac{(X_i - \mu)^2}{\sigma^3} = 0 \quad (15.10.14)$$

Upon simplifying the expression

$$\frac{n}{\sigma} = \sum_{i=1}^n \frac{(X_i - \mu)^2}{\sigma^3} \quad (15.10.15)$$

$$\sigma^2 = \sum_{i=1}^n \frac{(X_i - \mu)^2}{n} \quad (15.10.16)$$

Substituting (15.10.11) in (15.10.16)

$$\sigma^2 = \sum_{i=1}^n \frac{(X_i - \bar{X})^2}{n} \quad (15.10.17)$$

$$\sigma = \sqrt{\sum_{i=1}^n \frac{(X_i - \bar{X})^2}{n}} \quad (15.10.18)$$

Hence **Option 3** and **Option 4** are correct

15.11. Suppose  $X_1$  and  $X_2$  are independent and identically distributed random variables each following an exponential distribution with mean  $\theta$ , i.e., the common pdf is given by  $f_\theta(x) = \frac{1}{\theta}e^{-\frac{x}{\theta}}, 0 < x < \infty, 0 < \theta < \infty$ . Then which of the following is true? Conditional distribution of  $X_2$  given  $X_1 + X_2 = t$  is

- exponential with mean  $\frac{t}{2}$  and hence  $X_1 + X_2$  is sufficient for  $\theta$
- exponential with mean  $\frac{t\theta}{2}$  and hence  $X_1 + X_2$  is not sufficient for  $\theta$
- uniform(0,  $t$ ) and hence  $X_1 + X_2$  is sufficient for  $\theta$
- uniform(0,  $t\theta$ ) and hence  $X_1 + X_2$  is not sufficient for  $\theta$

**Solution:** Let  $f_{X_1, X_2}(x_1, x_2)$  denote the joint probability distribution of random variables  $X_1$  and  $X_2$ . Let  $Z$  be another random variable such that  $Z = X_1 + X_2$ . Let  $\Phi_{X_1}(\omega)$  and  $\Phi_Z(\omega)$  be the characteristic functions of the probability density functions  $f_{X_1}(x)$  and  $f_Z(x)$  respectively. The conditional probability density function of  $X_2$  can be defined by:

$$f_{X_2|(X_1+X_2=t)}(x_2) = \begin{cases} \frac{f_{X_1, X_2}(x_1, x_2)}{f_{(X_1+X_2)}(t)} & \text{if } x_1 + x_2 = t \\ 0 & \text{otherwise} \end{cases} \quad (15.11.1)$$

$$x_1 + x_2 = t \quad (15.11.2)$$

$$0 < x_1, x_2 < \infty \quad (15.11.3)$$

$$x_1 = t - x_2 \quad (15.11.4)$$

$$t - x_2 > 0 \quad (15.11.5)$$

$$x_2 < t \quad (15.11.6)$$

From equations (15.11.3) and (15.11.6), we can conclude that  $x_2 \in (0, t)$  if  $x_1 + x_2 = t$ . Also, given in the question,

$$0 < \theta < \infty \quad (15.11.7)$$

$$f_{X_1}(x_1) = \frac{1}{\theta}e^{-\frac{x_1}{\theta}}, 0 < x_1 < \infty \quad (15.11.8)$$

$$f_{X_2}(x_2) = \frac{1}{\theta}e^{-\frac{x_2}{\theta}}, 0 < x_2 < \infty \quad (15.11.9)$$

Since  $X_1$  and  $X_2$  are independent,

$$f_{X_1, X_2}(x_1, x_2) = f_{X_1}(x_1) \times f_{X_2}(x_2) \quad (15.11.10)$$

$$= \frac{1}{\theta}e^{-\frac{x_1}{\theta}} \times \frac{1}{\theta}e^{-\frac{x_2}{\theta}} \quad (15.11.11)$$

$$= \frac{1}{\theta^2}e^{-\frac{(x_1+x_2)}{\theta}} \quad (15.11.12)$$

$$\Phi_{X_1}(\omega) = \frac{1}{\theta} \int_0^\infty e^{i\omega x} e^{-\frac{x}{\theta}} dx \quad (15.11.13)$$

$$= \frac{1}{\theta} \times \frac{1}{i\omega - \frac{1}{\theta}} \left( e^{x(i\omega - \frac{1}{\theta})} \right) \Big|_0^\infty \quad (15.11.14)$$

$$= \frac{1}{1 - i\omega\theta} - \frac{\lim_{x \rightarrow \infty} (e^{x(i\omega - \frac{1}{\theta})})}{1 - i\omega\theta} \quad (15.11.15)$$

$$= \frac{1}{1 - i\omega\theta} - 0 = \frac{1}{1 - i\omega\theta} \quad (15.11.16)$$

$$\Phi_Z(\omega) = \left( \frac{1}{1 - i\omega\theta} \right)^2 \quad (15.11.17)$$

$$f_Z(x) = \frac{1}{2\pi} \int_{-\infty}^\infty \frac{e^{-i\omega x}}{\left( \frac{1}{1 - i\omega\theta} \right)^2} d\omega \quad (15.11.18)$$

The equation (15.11.18) is the characteristic function expression of a gamma random variable with  $k=2$ . Thus,

$$f_Z(x) = \frac{x^{k-1} e^{-\frac{x}{\theta}}}{\Gamma(k)\theta^k} \quad (15.11.19)$$

$$= \frac{x^{2-1} e^{-\frac{x}{\theta}}}{\Gamma(2)\theta^2} \quad (15.11.20)$$

$$= \frac{xe^{-\frac{x}{\theta}}}{\theta^2} \quad (15.11.21)$$

$$f_{X_2|(X_1+X_2=t)}(x_2) = \begin{cases} \frac{f_{X_1,X_2}(x_1,x_2)}{f_Z(t)} & x_2 \in (0,t) \\ 0 & \text{otherwise} \end{cases} \quad (15.11.22)$$

Let  $x_2 \in (0, t)$ .

$$f_{X_2|(X_1+X_2=t)}(x_2) = \frac{f_{X_1,X_2}(x_1,x_2)}{f_Z(t)} \quad (15.11.23)$$

$$= \frac{\frac{1}{\theta^2} e^{-\frac{(x_1+x_2)}{\theta}}}{\frac{1}{\theta^2} e^{-\frac{t}{\theta}}} \quad (15.11.24)$$

$$= \frac{e^{-\frac{(t)}{\theta}}}{e^{-\frac{t}{\theta}}} \quad (15.11.25)$$

$$= \frac{1}{t} \quad \forall x_2 \in (0, t) \quad (15.11.26)$$

The obtained pdf is uniform(0, t). Any distribution is sufficient for underlying parameter  $\theta$  if the conditional probability distribution of the data does not depend on the parameter  $\theta$ . And since the conditional distribution of  $X_2$  does not depend on  $\theta$  for any value of  $t$ ,  $X_1 + X_2$  is sufficient for  $\theta$ . Verifying the pdf,

$$\text{total probability} = \int_0^t f_{X_2|(X_1+X_2=t)}(x_2) dx_2 \quad (15.11.27)$$

$$= \int_0^t \frac{1}{t} dx_2 \quad (15.11.28)$$

$$= 1 \quad (15.11.29)$$

Hence, the correct answer is option (15.11c)

15.12. Let  $X_1, X_2, X_3, \dots, X_n$  be i.i.d observations from a distribution with continuous probability density function  $f$  which is symmetric around  $\theta$  i.e

$$f(x - \theta) = f(\theta - x) \quad (15.12.1)$$

for all real  $x$ . Consider the test  $H_0 : \theta = 0$  vs  $H_1 : \theta > 0$  and the sign test statistic

$$S_n = \sum_{i=1}^n \text{sign}(X_i) \quad (15.12.2)$$

where

$$\text{sign}(x) = \begin{cases} 1 & x > 0 \\ 0 & x = 0 \\ -1 & x < 0 \end{cases} \quad (15.12.3)$$

Let  $z_\alpha$  be the upper  $100(1-\alpha)^{\text{th}}$  percentile of the standard normal distribution where  $0 < \alpha < 1$ .

Which of the following is/are correct?

a) If  $\theta = 0$  then  $\lim_{n \rightarrow \infty} P\{S_n > \sqrt{n}z_\alpha\} = 1$

b) If  $\theta = 0$  then  $\lim_{n \rightarrow \infty} P\{S_n > \sqrt{n}z_\alpha\} = \alpha$

c) If  $\theta > 0$  then  $\lim_{n \rightarrow \infty} P\{S_n > \sqrt{n}z_\alpha\} = 1$

d) If  $\theta > 0$  then  $\lim_{n \rightarrow \infty} P\{S_n > \sqrt{n}z_\alpha\} = \alpha$

**Solution:**  $H_0 : \theta = 0$  Assume hypothesis  $H_0 : \theta = 0$  is true.

a) Given  $X$  is symmetric around zero.

$$f_X(x) = f_X(-x) \quad (15.12.4)$$

$$\int_0^\infty f_X(x) dx = \int_0^\infty f_X(-x) dx \quad (15.12.5)$$

i) Solving LHS of (15.12.5)

$$\int_0^\infty f_X(x) dx = \Pr(X \geq 0) \quad (15.12.6)$$

ii) Solving RHS of (15.12.5)

$$\int_0^\infty f_X(-x) dx \quad (15.12.7)$$

Changing  $-x \rightarrow x$  we get

$$\int_0^\infty f_X(-x) dx = \int_{-\infty}^0 f_X(x) dx \quad (15.12.8)$$

$$= \Pr(X \leq 0) \quad (15.12.9)$$

but

$$\int_{-\infty}^0 f_X(x) dx + \int_0^\infty f_X(x) dx = 1 \quad (15.12.10)$$

from (15.12.5), (15.12.8) and (15.12.10)

$$\int_{-\infty}^0 f_X(x) dx = \int_0^\infty f_X(x) dx = \frac{1}{2} \quad (15.12.11)$$

$$\implies \Pr(X \leq 0) = \Pr(X \geq 0) = \frac{1}{2} \quad (15.12.12)$$

b) Let  $Y$  be a random variable such that

$$Y = \text{sign}(X) \quad (15.12.13)$$

$$Y = \begin{cases} 1 & X > 0 \\ -1 & X < 0 \end{cases} \quad (15.12.14)$$

From (15.12.12) and (15.12.14) we have

$$\Pr(Y = -1) = \Pr(Y = 1) = \frac{1}{2} \quad (15.12.15)$$

So  $Y = \text{sign}(X)$  is also symmetric around zero.

$$\Rightarrow \mu_y = 0 \quad (15.12.16)$$

and variance is

$$\sigma_y^2 = (-1)^2 \left(\frac{1}{2}\right) + (1)^2 \left(\frac{1}{2}\right) \quad (15.12.17)$$

$$= 1 \quad (15.12.18)$$

c) Given

$$S_n = \sum_{i=1}^n \text{sign}(X_i) \quad (15.12.19)$$

$$S_n(\theta = 0) = \sum_{i=1}^n Y_i \quad (15.12.20)$$

From central limit theorem

$$Z = \lim_{n \rightarrow \infty} \sqrt{n} \left( \frac{\frac{S_n}{n} - \mu_y}{\sigma_y} \right) \quad (15.12.21)$$

$$= \lim_{n \rightarrow \infty} \sqrt{n} \left( \frac{S_n}{n} \right) \quad (15.12.22)$$

$$= \lim_{n \rightarrow \infty} \left( \frac{S_n}{\sqrt{n}} \right) \quad (15.12.23)$$

where  $Z$  is a standard normal variable  $N(0,1)$ .

i) Given

$$\alpha = P\{Z > z_\alpha\} \quad (15.12.24)$$

So from (15.12.23) and (15.12.24)

$$\lim_{n \rightarrow \infty} P\left\{ \frac{S_n}{\sqrt{n}} > z_\alpha \right\} = \alpha \quad (15.12.25)$$

$$\Rightarrow \lim_{n \rightarrow \infty} P\{S_n > \sqrt{n}z_\alpha\} = \alpha \quad (15.12.26)$$

$H_1 : \theta > 0$  is true

a) Given  $X$  is symmetric around  $\theta > 0$ . Let us

assume  $\theta = \theta_0 > 0$ .

$$f_X(\theta_0 - x) = f_X(\theta_0 + x) \quad (15.12.27)$$

$$\int_{\theta_0}^{\infty} f_X(\theta_0 - x) dx = \int_{\theta_0}^{\infty} f_X(\theta_0 + x) dx \quad (15.12.28)$$

i) Solving LHS of (15.12.28). Changing  $(\theta_0 - x) \rightarrow t$

$$\int_{\theta_0}^{\infty} f_X(\theta_0 - x) dx = \int_{-\infty}^0 f_X(t) dt \quad (15.12.29)$$

$$= \Pr(X \leq 0) \quad (15.12.30)$$

ii) Solving RHS of (15.12.28). Changing  $(\theta_0 + x) \rightarrow t$

$$\int_{\theta_0}^{\infty} f_X(\theta_0 + x) dx = \int_{2\theta_0}^{\infty} f_X(t) dt \quad (15.12.31)$$

$$= \int_0^{\infty} f_X(t) dt - \int_0^{2\theta_0} f_X(t) dt \quad (15.12.32)$$

$$= \Pr(X \geq 0) - k \quad (15.12.33)$$

where

$$k = \int_0^{2\theta_0} f_X(t) dt > 0 \quad (15.12.34)$$

From (15.12.28), (15.12.14) and (15.12.33)

$$\Pr(X \geq 0) > \Pr(X \leq 0) \quad (15.12.35)$$

b) So

$$\Pr(Y = 1) > \Pr(Y = -1) \quad (15.12.36)$$

Therefore, if we perform the experiment and find the value of  $\left(\frac{S_n}{\sqrt{n}}\right)$ , it is most likely to occur on the right side of the distribution of  $\left(\frac{S_n}{\sqrt{n}}\right)$ . In (15.12.23) it is shown that distribution of the random variable  $\left(\frac{S_n}{\sqrt{n}}\right)$  is  $N(0,1)$  when  $n$  is very large. So

$$\lim_{n \rightarrow \infty} P\left\{ \frac{S_n}{\sqrt{n}} > Z_\alpha \right\} = 1 \quad (15.12.37)$$