

Measuring charged current neutrino interactions in the Electromagnetic Calorimeters in near detector of T2K

Dominic Brailsford
TODO

A thesis submitted to Imperial College London
for the degree of Doctor of Philosophy

Abstract

To write

Declaration

This dissertation is the result of my own work, except where explicit reference is made to the work of others, and has not been submitted for another qualification to this or any other university. This dissertation does not exceed the word limit for the respective Degree Committee.

Dominic Brailsford

Acknowledgements

Something about my supervisor ...

Preface

This thesis describes my analysis of the ν_μ charged-current cross-section on lead using the T2K near detector electromagnetic calorimeters.

Contents

1	Introduction	1
1.1	The state of the field	2
1.2	The future	4
	Bibliography	9
	List of Figures	11
	List of Tables	13

“These chickens jackin’ my style. ”

— Fergie

Chapter 1

Introduction

The field of neutrino physics is currently undergoing a revolution. With its tenuous postulation [1] acting as a future omen, the neutrino's mark on history would not become apparent from its discovery [2–4], but rather from a spate of surprising discoveries at the end of the 20th century [5–7] which conclusively proved that the Standard Model, while very successful, was incomplete. This revelation was experimental proof of Maki, Nagakawa and Sakata's extension [8] to Pontecorvo's theory of neutrino oscillation [9] with the inclusion of the Mikheyev-Smirnov-Wolfenstein (MSW) effect [10, 11]. The findings were groundbreaking as the underlying theory requires massive neutrinos, which is in direct contradiction to the Standard Model. The, now, standard theory of neutrino oscillation defines three neutrino flavours and three neutrino masses. However, the map between flavour and mass is not one-to-one, but rather a rotation of mass space onto flavour space. The main consequence of this rotation is that the flavour eigenstates are a superposition of mass eigenstates, namely

$$|\nu_\alpha\rangle = \sum_{i=1}^3 U_{\alpha i}^* |\nu_i\rangle, \quad (1.1)$$

where $\alpha \in \{e, \mu, \tau\}$, ν_i are the neutrino mass eigenstates and $U_{\alpha i}^*$ is an element of a unitary rotation matrix which is known as the PMNS mixing matrix. As there are 3 mass and flavour eigenstates, the PMNS matrix is a 3×3 matrix and is often

parameterised as

$$U \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 3 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (1.2)$$

where $c_{ij} \equiv \cos \theta_{ij}$ and $s_{ij} \equiv \sin \theta_{ij}$. θ_{ij} are known as the mixing angles which parameterise how strong mixings between the flavour and mass eigenstates are and δ is a CP violating phase. The most surprising observable feature of this mechanism is the non-zero probability to detect a neutrino of specific flavour which was created at source in a different flavour state. By propagating the mass eigenstates through time, one can arrive at this probability which has the following form

$$P(\nu_\alpha \rightarrow \nu_\beta) = |\langle \nu_\beta | \nu(t) \rangle|^2 = |U_{\beta k} e^{-iE_k t} U_{\alpha k}^*|^2, \quad (1.3)$$

where $\nu(t)$ is the time-dependent neutrino mass eigenstate and E_k is the energy of the ν_k . For an accelerator-based neutrino oscillation experiment, the beam will be ν_μ dominated. So, the ν_μ survival probability, $P(\nu_\mu \rightarrow \nu_\mu)$, and ν_e appearance probability, $P(\nu_\mu \rightarrow \nu_e)$, have are typically of interest and can be approximated in the following forms

$$P(\nu_\mu \rightarrow \nu_\mu) \approx 1 - \cos^4 \theta_{13} \sin^2 2\theta_{23} \sin^2 \left(1.27 \frac{\Delta m_{23}^2}{(\text{eV}^2)} \frac{L}{(\text{km})} \frac{(\text{GeV})}{E} \right) \quad (1.4)$$

$$P(\nu_\mu \rightarrow \nu_\mu) \approx \sin^2 2\theta_{13} \sin^2 \theta_{23} \sin^2 \left(1.27 \frac{\Delta m_{23}^2}{(\text{eV}^2)} \frac{L}{(\text{km})} \frac{(\text{GeV})}{E} \right), \quad (1.5)$$

where $\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$, L is the distance the neutrino propagates and E is the energy of the neutrino.

1.1 The state of the field

Data provided from a wide range of experiments show excellent agreement with the theory of neutrino oscillation and with a 3 flavour neutrino picture. Global fits applied to the data provided by these experiments gives best fit values for the oscil-

Parameter	best-fit ($\pm 1\sigma$)
Δm_{12}^2 [10^{-5}eV^2]	$7.54^{+0.26}_{-0.22}$
$ \Delta m^2 $ [10^{-3}eV^2]	2.43 ± 0.06 (2.36 ± 0.06)
$\sin^2 \theta_{12}$	0.308 ± 0.017
$\sin^2 \theta_{23}, \Delta m^2 > 0$	$0.437^{+0.033}_{-0.023}$
$\sin^2 \theta_{23}, \Delta m^2 < 0$	$0.455^{+0.039}_{-0.031}$
$\sin^2 \theta_{13}, \Delta m^2 > 0$	$0.0234^{+0.0020}_{-0.0019}$
$\sin^2 \theta_{13}, \Delta m^2 < 0$	$0.0240^{+0.019}_{-0.022}$
$\sin^2 \theta_{13}, \Delta m^2 < 0$	$0.0240^{+0.019}_{-0.022}$
δ/π (2σ range quoted)	$1.39^{+0.38}_{-0.27}$ ($1.31^{+0.29}_{-0.33}$)

Table 1.1: The best-fit values of the 3-neutrino oscillation parameters. $\Delta m^2 \equiv m_3^2 - (m_2^2 - m_1^2)/2$. The values (values in brackets) correspond to $m_1 < m_2 < m_3$ ($m_3 < m_1 < m_2$) [12].

lation parameters, which are summarised in table 1.1 [12]. The experiments which provided the data inputs to the global fit generally fall into one of four categories, with each category sensitive to a different subset of the neutrino oscillation parameters.

Solar neutrino experiments detect neutrinos generated in the core of the Sun as a result of nuclear reaction chains. Such experiments are primarily sensitive to θ_{12} and Δm_{12}^2 which are often referred to as the solar mixing parameters. The final state neutrinos created in the Sun's core are MeV-scale ν_e but, because of propagation through the core's surrounding matter, the MSW effect results in a highly pure state of ν_2 at the Sun's surface. As ν_2 is a mass eigenstate, no oscillation occurs between the surface of the Sun and the Earth. Homestake [13], SAGE [14] and SNO [6] are examples of such experiments.

Reactor neutrino experiments measure $\bar{\nu}_e$ disappearance provided by inverse β decay in nuclear reactors with an average neutrino energy of 3 MeV. The baseline for oscillations varies between experiments, but a baseline of around 1 km provides excellent sensitivity to θ_{13} . Examples of reactor experiments are CHOOZ [15], Double CHOOZ [16], Daya Bay [17] and RENO [18].

Atmospheric neutrino experiments detect neutrinos which are produced from π and K mesons, created by cosmic rays interactions with the upper atmosphere of the Earth,

decay. The neutrinos produced are a mixture of ν_μ , $\bar{\nu}_\mu$, ν_e and $\bar{\nu}_e$. Because the cosmic ray flux is fairly uniform, atmospheric neutrino experiments are exposed to neutrinos from all directions, which results in a very wide range of oscillation baselines. The oscillation parameters that such experiments are sensitive to are θ_{23} and Δm_{13}^2 . Super-Kamiokande [5] is an example of an atmospheric neutrino experiment.

Accelerator neutrino experiments produce beams of high purity ν_μ (or $\bar{\nu}_\mu$) at GeV-scale energy with wide ranging baselines which are generally $\mathcal{O}(100 \text{ km})$. The highly man-made nature of such experiments allows almost complete control over L/E allowing careful tuning of parameter sensitivity. Accelerator neutrino experiments are generally sensitive to θ_{13} , θ_{23} , Δm_{13}^2 and δ . K2K [19], MINOS [20], T2K [21] and NO ν A [22] are examples of such experiments.

1.2 The future

It should be clear that an immense amount of progress has been made in the field, with remarkable contributions to the picture coming only in the last 20 years. However, there are several key questions which remain unanswered.

By far the most sought after answer is whether CP violation occurs in the lepton sector. The magnitude of CP violation is encapsulated in the CP violating phase δ and so it is this parameter which current and future long-baseline experiments are aiming towards. Currently, T2K and NO ν A can provide the strongest constraints on δ . The future long-baseline experiments, Hyper-Kamiokande [23] and DUNE (formerly LBNE) [24] are being designed with a possible measurement of δ as a primary goal.

The second question still to be answered is the ordering of the mass eigenstates. Specifically is $m_3 \gg m_2 > m_1$ (the normal mass hierarchy) or $m_2 > m_1 \gg m_3$ (the inverted mass hierarchy)? The matter effects introduced by the MSW effect are mass hierarchy dependent. So, for very long-baseline experiments, there is mass hierarchy sensitivity. Currently NO ν A is set to resolve the mass hierarchy problem. However, both Hyper-Kamiokande (via atmospheric measurements) and DUNE have measurement of the hierarchy as a primary goal.

Oscillation experiments only have the capability to measure the square of the mass splitting differences. This means that all oscillation experiments have no sensitivity to the absolute neutrino mass scale. This means an entirely different type of neutrino experiment is required. Neutrinos are one of the final states associated with β decay and the mass of the neutrino should appear as a cut off in the β spectrum. The visibility of the cut-off entirely depends on the mass scale. So, the KATRIN experiment [25] will

attempt to utilise the β decay feature, with a neutrino mass sensitivity of 0.2 eV.

It is not currently known whether neutrinos are their own anti-particle, otherwise known as Majorana neutrinos. A number of experiments are currently investigating this, all by searching for neutrinoless double β decay. A large neutrinoless double β decay experiment effort is ongoing, including EXO [26], SuperNEMO [27] and SNO+ [28].

While the long-baseline neutrino oscillation programme has been very successful, the short-baseline programme has seen several anomalies [29]. The LSND experiment found evidence of $\bar{\nu}_e$ in a $\bar{\nu}_\mu$ beam, which was consistent with neutrino oscillations [30]. However the data suggested a mass-squared splitting of 0.2-10 eV². This large splitting is consistent with a fourth species of neutrino. Because the data suggesting 3 flavours of weakly-interacting neutrino is strong, this postulated fourth species must be sterile. More recently, the Mini-BooNE experiment observed a similar short baseline excess of $\bar{\nu}_e$ in a $\bar{\nu}_\mu$ beam with a mass-squared splitting of 0.01-1.0 eV² [31], further suggesting the sterile hypothesis. New experiments are now under development which aim to test this hypothesis, which include MicroBooNE [32] and SBND (formally LAr1-ND) [33].

Bibliography

- [1] W. Pauli, Open letter to the Gauverein meeting in Tübingen, 1930.
- [2] C. L. Cowan, F. Reines, F. B. Harrison, H. W. Kruse, and A. D. McGuire, *Science* **124**, 103 (1956).
- [3] G. Danby *et al.*, *Phys. Rev. Lett.* **9**, 36 (1962).
- [4] K. Kodama *et al.*, *Physics Letters B* **504**, 218 (2001).
- [5] Y. Fukuda *et al.*, *Phys. Rev. Lett.* **81**, 1562 (1998).
- [6] Q. R. Ahmad *et al.*, *Phys. Rev. Lett.* **87**, 071301 (2001).
- [7] K. Eguchi *et al.*, *Phys. Rev. Lett.* **90**, 021802 (2003).
- [8] Z. Maki, M. Nakagawa, and S. Sakata, *Progress of Theoretical Physics* **28**, 870 (1962).
- [9] B. Pontecorvo, *JETP Lett.* **33**, 549 (1957).
- [10] L. Wolfenstein, *Phys. Rev. D* **17**, 2369 (1978).
- [11] S. Mikheev and A. Y. Smirnov, *Sov.J.Nucl.Phys.* **42**, 913 (1985).
- [12] Particle Data Group, K. Olive *et al.*, *Chin.Phys.* **C38**, 090001 (2014).
- [13] B. T. Cleveland *et al.*, *The Astrophysical Journal* **496**, 505 (1998).
- [14] J. N. Abdurashitov *et al.*, *Phys. Rev. C* **80**, 015807 (2009).
- [15] M. Apollonio *et al.*, *The European Physical Journal C - Particles and Fields* **27**, 331 (2003).
- [16] Y. Abe *et al.*, *Physics Letters B* **723**, 66 (2013).
- [17] F. P. An *et al.*, *Phys. Rev. Lett.* **108**, 171803 (2012).

- [18] J. K. Ahn *et al.*, Phys. Rev. Lett. **108**, 191802 (2012).
- [19] M. H. Ahn *et al.*, Phys. Rev. D **74**, 072003 (2006).
- [20] D. G. Michael *et al.*, Phys. Rev. Lett. **97**, 191801 (2006).
- [21] (T2K Collaboration), K. Abe *et al.*, Phys. Rev. Lett. **112**, 061802 (2014).
- [22] NOvA, D. Ayres *et al.*, (2004), hep-ex/0503053.
- [23] Hyper-Kamiokande Working Group, K. Abe *et al.*, (2014), 1412.4673.
- [24] LBNE, C. Adams *et al.*, (2013), 1307.7335.
- [25] C. Weinheimer, Progress in Particle and Nuclear Physics **48**, 141 (2002).
- [26] M. Auger *et al.*, Phys. Rev. Lett. **109**, 032505 (2012).
- [27] A. Barabash and the SuperNemo Collaboration, Journal of Physics: Conference Series **375**, 042012 (2012).
- [28] SNO+, M. C. Chen, (2008), 0810.3694.
- [29] J. Fan and P. Langacker, Journal of High Energy Physics **2012** (2012).
- [30] LSND, A. Aguilar-Arevalo *et al.*, Phys.Rev. **D64**, 112007 (2001), hep-ex/0104049.
- [31] MiniBooNE, A. Aguilar-Arevalo *et al.*, Phys.Rev.Lett. **110**, 161801 (2013), 1207.4809.
- [32] C. M. Ignarra, ArXiv e-prints (2011), 1110.1604.
- [33] LArTPC, C. Adams *et al.*, (2013), 1309.7987.

List of Figures

List of Tables

1.1	The best-fit values of the 3-neutrino oscillation parameters. $\Delta m^2 \equiv m_3^2 - (m_2^2 - m_1^2) / 2$. The values (values in brackets) correspond to $m_1 < m_2 < m_3$ ($m_3 < m_1 < m_2$) [12].	3
-----	---	---