

Quadratic Formula Derivation

Alex Glandon

May 2026

$$Ax^2 + Bx + C = 0$$

$$\begin{aligned}(Dx + E)^2 &= D^2x^2 + 2DEx + E^2 \\ &= Ax^2 + Bx + E^2\end{aligned}$$

$$A = D^2$$

$$\text{case 1: } D = +\sqrt{A}$$

$$B = DE = +2\sqrt{A}E \rightarrow E = \frac{+B}{2\sqrt{A}}$$

$$\left(+\sqrt{A}x + \frac{B}{2\sqrt{A}}\right)^2 = Ax^2 + Bx + \frac{B^2}{4A}$$

$$Ax^2 + Bx + C = 0$$

$$Ax^2 + Bx + \frac{B^2}{4A} - \frac{B^2}{4A} + C = 0$$

$$\left(+\sqrt{A}x + \frac{B}{2\sqrt{A}}\right)^2 - \frac{B^2}{4A} + C = 0$$

$$\left(+\sqrt{A}x + \frac{B}{2\sqrt{A}}\right)^2 = \frac{B^2}{4A} - C$$

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

or

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

case 2: $D = -\sqrt{A}$

$$B = DE = -2\sqrt{A}E \rightarrow E = \frac{-B}{2\sqrt{A}}$$

$$\left(-\sqrt{A}x - \frac{B}{2\sqrt{A}}\right)^2 = Ax^2 + Bx + \frac{B^2}{4A}$$

$$Ax^2 + Bx + C = 0$$

$$Ax^2 + Bx + \frac{B^2}{4A} - \frac{B^2}{4A} + C = 0$$

$$\left(-\sqrt{A}x - \frac{B}{2\sqrt{A}}\right)^2 = \frac{B^2}{4A} - C$$

$$-\sqrt{A}x - \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

or

$$-\sqrt{A}x - \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

from case 1 and case 2 we have four options:

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

or

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

or

$$-\sqrt{A}x - \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

or

$$-\sqrt{A}x - \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

or if we multiply the bottom two equations by -1

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

or

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

or

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

or

$$+\sqrt{A}x + \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

since the first and last are the same and the second and third are the same, we are left with the following two equations

$$+\sqrt{Ax} + \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

or

$$+\sqrt{Ax} + \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

simplifying:

$$+\sqrt{Ax} + \frac{B}{2\sqrt{A}} = +\sqrt{\frac{B^2}{4A} - C}$$

$$+Ax + \frac{B}{2} = +\sqrt{(\frac{B^2}{4A} - C)A}$$

$$+Ax + \frac{B}{2} = +\sqrt{\frac{B^2}{4} - AC}$$

$$+2Ax + B = +\sqrt{(\frac{B^2}{4} - AC)2^2}$$

$$+2Ax + B = +\sqrt{(\frac{B^2}{4} - AC)4}$$

$$+2Ax + B = +\sqrt{B^2 - 4AC}$$

$$+2Ax = -B + \sqrt{B^2 - 4AC}$$

$$x = \frac{-B + \sqrt{B^2 - 4AC}}{2A}$$

or

$$+\sqrt{Ax} + \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

simplifying:

$$+\sqrt{Ax} + \frac{B}{2\sqrt{A}} = -\sqrt{\frac{B^2}{4A} - C}$$

$$+Ax + \frac{B}{2} = -\sqrt{(\frac{B^2}{4A} - C)A}$$

$$+Ax + \frac{B}{2} = -\sqrt{\frac{B^2}{4} - AC}$$

$$+2Ax + B = -\sqrt{(\frac{B^2}{4} - AC)2^2}$$

$$+2Ax + B = -\sqrt{(\frac{B^2}{4} - AC)4}$$

$$+2Ax + B = -\sqrt{B^2 - 4AC}$$

$$+2Ax = -B - \sqrt{B^2 - 4AC}$$

$$x = \frac{-B - \sqrt{B^2 - 4AC}}{2A}$$