

Decision and Risk Analysis

Lecture 7: **A Simple Manual Example of MCMC**

Metropolis-Hastings for a Bernoulli Parameter

DAEN 427 / ISEN 427 / ISEN 627 - Fall 2026

Alexander Abuabara



We want to understand how MCMC can generate samples from a posterior distribution.

Suppose we observe four Bernoulli trials:

$$y = (1, 1, 1, 0)$$

That is:

3 successes, 1 failure

The unknown parameter is:

$$\theta = \Pr(y = 1)$$

Prior and Likelihood

Assume a uniform prior:

$$\theta \sim \text{Uniform}(0, 1)$$

The likelihood for 3 successes and 1 failure is:

$$p(y \mid \theta) \propto \theta^3(1 - \theta)^1$$

Since the prior is uniform, the posterior is proportional to:

$$p(\theta \mid y) \propto \theta^3(1 - \theta)$$

We define the unnormalized posterior density:

$$f(\theta) = \theta^3(1 - \theta)$$

Metropolis-Hastings Rule

Start at some current value:

$$\theta_t$$

Propose a new value:

$$\theta^*$$

Compute the acceptance ratio:

$$r = \frac{f(\theta^*)}{f(\theta_t)}$$

Then compute:

$$\alpha = \min(1, r)$$

Accept the proposed value if:

$$u < \alpha, \quad u \sim \text{Uniform}(0, 1)$$

Step 1: Move from 0.50 to 0.60

Start at:

$$\theta_0 = 0.50$$

Compute:

$$f(0.50) = 0.50^3(1 - 0.50) = 0.125 \times 0.50 = 0.0625$$

Propose:

$$\theta^* = 0.60$$

Compute:

$$f(0.60) = 0.60^3(1 - 0.60) = 0.216 \times 0.40 = 0.0864$$

Acceptance ratio:

$$r = \frac{0.0864}{0.0625} = 1.3824$$

So:

$$\alpha = 1$$

Accept:

$$0.50 \rightarrow 0.60$$

Step 2: Move from 0.60 to 0.80

Current value:

$$\theta_t = 0.60$$

Propose:

$$\theta^* = 0.80$$

Compute:

$$f(0.80) = 0.80^3(1 - 0.80) = 0.512 \times 0.20 = 0.1024$$

Acceptance ratio:

$$r = \frac{0.1024}{0.0864} = 1.185$$

Thus:

$$\alpha = 1$$

Accept:

$$0.60 \rightarrow 0.80$$

Step 3: A Rejected Proposal

Current value:

$$\theta_t = 0.80$$

Propose:

$$\theta^* = 0.95$$

Compute:

$$f(0.95) = 0.95^3(1 - 0.95) \approx 0.8574 \times 0.05 = 0.0429$$

Acceptance ratio:

$$r = \frac{0.0429}{0.1024} \approx 0.419$$

So:

$$\alpha = 0.419$$

Suppose we draw:

$$u = 0.70$$

Since:

$$0.70 > 0.419$$

reject the proposal and stay at:

$$\theta = 0.80$$

Step 4: Move from 0.80 to 0.70

Current value:

$$\theta_t = 0.80$$

Propose:

$$\theta^* = 0.70$$

Compute:

$$f(0.70) = 0.70^3(1 - 0.70) = 0.343 \times 0.30 = 0.1029$$

Acceptance ratio:

$$r = \frac{0.1029}{0.1024} \approx 1.005$$

Thus:

$$\alpha = 1$$

Accept:

$$0.80 \rightarrow 0.70$$

Manual Chain

The chain we recorded is:

0.50, 0.60, 0.80, 0.80, 0.70

Notice the repeated value:

0.80, 0.80

This happens because a rejected proposal keeps the chain at the current state.

After many iterations, the histogram of the chain approximates the posterior:

$$p(\theta \mid y)$$

Summary Table

Step	Current	Proposal	$f(\text{Current})$	$f(\text{Proposal})$	α	Result
1	0.50	0.60	0.0625	0.0864	1.000	Accept
2	0.60	0.80	0.0864	0.1024	1.000	Accept
3	0.80	0.95	0.1024	0.0429	0.419	Reject
4	0.80	0.70	0.1024	0.1029	1.000	Accept

Key Idea

MCMC does not only search for the best value.

It constructs a chain whose long-run behavior matches the posterior distribution.

Regions with higher posterior probability are visited more often.

Regions with lower posterior probability are visited less often, but not necessarily never.

That is why MCMC provides uncertainty, not just a point estimate.

In the simple example, the unknown was one parameter:

$$\theta$$

In Bayesian linear regression, the unknowns are usually:

$$(\beta_0, \beta_1, \sigma)$$

MCMC samples from:

$$p(\beta_0, \beta_1, \sigma \mid x, y)$$

Each draw gives one plausible regression model:

$$y = \beta_0 + \beta_1 x + \varepsilon$$

The collection of draws gives the full posterior uncertainty.