

Lecture 8 (section 1.5)

Section 1.5 — elementary matrices; method for finding A^{-1}

\Rightarrow Elementary matrices:

E is an elementary matrix if E is obtained from I_n (the identity matrix) by performing a single elementary row operation.

Is elementary?

•

$$\begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

yes $\Rightarrow r_1 = 3 \cdot r_1^I$

row multiplication;

•

$$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

yes $\Rightarrow r_1^I \leftrightarrow r_3^I$

row interchange;

•

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -4 & 0 & 1 \end{bmatrix}$$

yes $\Rightarrow r_3 = r_3^I + (-4) \cdot r_1^I$

linear combination of 2 rows

\Rightarrow Note: every elementary matrix is invertible:

E

E^{-1}

$$\begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} \frac{1}{3} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -4 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 4 & 0 & 1 \end{bmatrix}$$

Theorem:

Let A be $n \times n$ (a square) matrix. The following statements are equivalent:

- A is invertible
- the homogeneous system

$$A \cdot \overline{X} = \overline{0}$$

has only zero solution

- The RREF of A is I_n
- A is a product of elementary matrices,

$$A = E_1 \cdot E_2 \cdots E_k$$

Algorithm for finding A^{-1}

- Form $n \times (2n)$ matrix:

$$\left[A \mid I_n \right]$$

- perform elementary operations to bring the left half of this matrix to RREF
- repeat identical sequence of operations with the right half of the matrix
- if the left side is I_n , the right side is A^{-1}

\Rightarrow Example:

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

Follow the algorithm:

$$\left[A \mid I_3 \right] = \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \rightarrow r_3 \rightarrow r_3 - r_1 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 0 & 1 \end{array} \right]$$

$$\rightarrow r_3 \rightarrow r_3 - r_2 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & -2 & -1 & -1 & 1 \end{array} \right] \rightarrow r_3 \rightarrow -\frac{1}{2} \cdot r_3 \rightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \end{array} \right] \rightarrow \begin{array}{l} r_1 \rightarrow r_1 - r_3 \\ r_2 \rightarrow r_2 - r_3 \end{array} \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 1 & \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \end{array} \right] = \left[I_3 \mid A^{-1} \right]$$

\Rightarrow

$$A^{-1} = \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \end{bmatrix}$$

\Rightarrow Check that $A \cdot A^{-1} = A^{-1} \cdot A = I_3$