×

# Graded quiz on Cartesian Plane and Types of Function



**13/13** points earned (100%)

Quiz passed!

Back to Week 2



# Be Recognized for Your Achievements.

"Course Certificates give you the recognition you need to get the job, the material gives you the skills to do the job.

It makes you look more valuable because you are more valuable." - Peter B., USA, Software Developer

# Showcase Your Accomplishment! Earn Your Course Certificate! \$49 USD >



1/1 points

1.

Which of the following points in the Cartesian Plane have positive x-coordinate and negative y-coordinate?

- O(0,0)
- O (5,7)
- $\mathbf{O}$  (7, -1)

#### Correct

The x-coordinate, 7, is positive, and the y-coordinate, -1, is negative.

- O (-4,5)
- 1/1 points
- 2. Which of the following points is in the first quadrant of the Cartesian Plane?
  - O(-5,1)
  - O (5,-1)
  - O (-4, -7)
  - $\mathbf{O}$  (7,11)

## Correct

The first quadrant is defined to be all points in the Cartesian plane whose coordinates are both positive.

**/** 

1/1 points

3.

Let A,B,C,D be points in the Cartesian Plane, and let the set  $S=\{B,C,D\}$ 

Suppose that the distances from A to B,C,D are 5.3,2.1, and 11.75, respectively.

Which of the following points is the nearest neighbor to the point A in the set S?

- Ор
- O
- 0

### Correct

The distance from A to C is 2.1 and that is smaller than the distance from A to any other element of S.

Ов



1/1 points

4

Find the distance between the points  $A=\left( 2,2\right)$  and  $B=\left( -1,-2\right)$ .

- O -25
- O 5

#### Correct

Recall that the distance between points (a,b) and (c,d) is  $\sqrt{(c-a)^2+(d-b)^2}$ 

In this case we have:

$$\sqrt{\left(-1-2
ight)^2+\left(-2-2
ight)^2}=\sqrt{\left(-3
ight)^2+\left(-4
ight)^2}=\sqrt{25}=5$$

- 1
- 25



1/1 points

5.

Find the slope of the line segment between the points  $A=\left(0,1\right)$ and B = (1, 0).





The slope of this line segment is  $\frac{0-1}{1-0} = -1$ 



1/1 points

Find the point-slope form of the equation of the line with slope -2that goes through the point (5,4).

O 
$$y-5=-2(x-4)$$

O 
$$y-4=-2(x-5)$$

#### Correct

The point-slope form for the equation of a line with slope m that goes through the point  $(x_0,y_0)$  is

$$y - y_0 = m(x - x_0)$$
.

In this case, the slope m=-2 is given and the point  $(\mathbf{5},\mathbf{4})$  on the line is given.

O 
$$y-4=2(x-5)$$



1/1 points

7.

Which of the following equations is for a line with the same slope as y=-3x+2?

$$O \quad y = 5x$$

$$O \quad y = 8x - 3$$

$$0 \quad y = -3x - 8$$

#### Corroct

The slope-intercept formula for a line is y=mx+b, where m is the slope and b is the y-coordinate of the point where the line hits the y-axis.

This line has slope m=-3 which is the same slope as the given line.

$$O \quad y = 5x + 2$$



1/1 points

8.

Which of the following equations is for a line with the same y-intercept as y=-3x+2?

O 
$$y = -3x - 8$$



The the slope-intercept formula for a line is y=mx+b, where m is the slope and b is the y-coordinate of the point where the line hits the y-axis. This line has a y-intercept of 2 which is the same as the given line.

$$O \quad y = 8x - 3$$

O 
$$y=5x$$



1/1 points

9

How many lines contain both the point A=(1,1) and the point B=(2,2)?

- O None
- $O^{2}$
- O Infinity many
- O 1

#### Corroct

The line with equation y=x is the one and only line that meets the stated requirements.



1/1 points

10.

Suppose that we have two sets,  $A=\{a,b\}$  and  $Z=\{x,y\}$ . How many different functions  $F:A\to Z$  are possible?

- O There are none
- O

4

#### Correct

A function F:A o Z is a rule which assigns an element  $F(a)\in Z$  to each element  $a\in A$ .

There are two elements in A; namely, a and b. For each of these elements, there are two assignment choices we could make: x and y.

Here are the four possible functions:

$$F(a)=x, F(b)=y$$
, OR

$$F(a)=y, F(b)=x$$
, OR

$$F(a)=x, F(b)=x$$
, OR

$$F(a) = y, F(b) = y.$$

- O There are infinitely many
- O 1

1 / 1



points

11.

How many graphs contain both the point  $A=\left(0,0\right)$  and the point  $B=\left(1,1\right)$ 



#### Correct

The graphs of  $f(x)=x, g(x)=x^2, h(x)=x^3, s(x)=x^4, \ldots$  all contain both A and B

- O None
- O 1
- $\bigcirc$  2



1/1 points

12.

Suppose that  $g:\mathbb{R}\to\mathbb{R}$  is a continuous function whose graph intersects the x-axis more than once. Which of the following statements is true?

 $oldsymbol{\mathsf{Q}}$  is neither strictly increasing nor strictly decreasing.

# Correct

The function g fails the horizontal line test, so it can neither be strictly increasing nor strictly decreasing.

- $igcolon{d}{igcolon}g$  is strictly decreasing.
- O All of the above.
- $\bigcirc$  g is strictly increasing.