

Conductor-changing prime-degree isogenies of ECC2K-130

Report dated 2026-10-07. Study directory research/ecc2k130_conductor_isogenies_20261007/. Standalone study: no H-*, EV-*, DEC-* or RUN-* record was written. Run artifacts are the .out files in the directory. Labels: **DERIVED** proof or exact computation from stated facts; **MEASURED** observed in this study's runs; **INDEPENDENTLY VERIFIED** confirmed by a second, independent method; **PROPOSED** a hypothesis, not shown here.

1. Question and scope

Which prime-degree isogenies from ECC2K-130 change the conductor of the endomorphism ring, and which of them can be computed? The scope is the whole $a_2 = 0$ isogeny class over $\mathbb{F}_{2^{131}}$, all prime degrees.

2. Object

- $E : y^2 + xy = x^3 + 1$ over $\mathbb{F}_2[x]/(x^{131} + x^8 + x^3 + x^2 + 1)$. A curve E_b is named by the integer of the bits of b . **DERIVED**
- $\#E = 4\ell$ with $\ell = 680564733841876926932320129493409985129$ prime; the trace is -22283658519494248867 . **DERIVED**
- $\text{End}(E) = \mathcal{O}_K$ with $K = \mathbb{Q}(\sqrt{-7})$ and $h = 1$. The Frobenius conductor is $f_\pi = [\mathcal{O}_K : \mathbb{Z}[\pi]] = 263 \cdot 146505763881528721$; the prime 263 splits and the other is inert. **DERIVED**




Figure 1: The isogeny class by level. Solid: computed here, with endpoints verified. Dashed: exists by theory but cannot be computed. Source: volcano_ecc2k130.dot.

3. Which prime degrees can change the conductor

An ℓ -isogeny between ordinary curves changes the conductor by a factor of at most ℓ , and can change it only if $\ell \mid f_\pi$ (Kohel). Every other prime degree gives a horizontal isogeny, with conductor ratio 1. Here that leaves exactly two degrees: 263 and 146505763881528721. **DERIVED**

4. The 263-isogenies (computed)

- $\pi \equiv -1 \pmod{263}$, so $E[263]$ is $\mathbb{F}_q$ -rational on the quadratic twist $E_t : y^2 + xy = x^3 + x^2 + 1$, with the same x-coordinates. No extension field is needed. **DERIVED**
- $\#E_t = 2 \cdot 263^2 \cdot 19678316408850118605767852657510239$ and $E_t(\mathbb{F}_q) \cong \mathbb{Z}/(\#E_t/263) \times \mathbb{Z}/263$, so $E_t[263]$ is all rational. **MEASURED**
- All 264 kernels, via char-2 Velu ($b' = 1 + v + v^2$, checked 6/6 against PARI ellisogeny at $m = 37$):
 - 2 map back to E . These are the horizontal isogenies; since 263 splits and $h = 1$, they are endomorphisms. **MEASURED**
 - 262 give 262 distinct curves, the whole 263-floor ($h(\mathcal{O}_{263}) = 262$). Every one has $\#E = N$ by ellcard. **MEASURED**

- **Level checks.** Each image's twist has a cyclic $\mathbb{Z}/263^2$ 263-part (262/262), against $(\mathbb{Z}/263)^2$ on the crater. The horizontal 11-isogeny cycle has length 131 (262/262), equal to $\text{ord}[l_{11}]$ in $\text{Cl}(-7 \cdot 263^2)$; on the crater it is 1. **INDEPENDENTLY VERIFIED**
- The 262 curves form 2 Galois orbits of 131, with representatives $b' = 0x3415808f755dad3868860b0a0fb8103$ and $0xa05d3c82ee7321154f4f4fd13cea070$ ($a_2 = 0$). **MEASURED**

5. The 146505763881528721-isogenies: why they cannot be computed

All $\ell + 1 = 146505763881528722$ of them exist and descend. **DERIVED** Every route to one is out of reach:

Route	Barrier
Kernel points (Vélu, $\sqrt{\ell}$ u)	$\pi \equiv c \pmod{\ell}$, with $\text{ord}(c) = 1.22 \cdot 10^{16}$ and $-1 \in \langle c \rangle$. The kernel x-coordinates first appear in $\mathbb{F}_{2^{131k}}$ with $k = 6104406828397030$, where one element takes $8.0 \cdot 10^{17} \approx 2^{59.5}$ bits. DERIVED
Kernel polynomial over $\mathbb{F}_{2^{131}}$	Degree $(\ell - 1)/2 = 7.3 \cdot 10^{16}$; about 1.2 EB just to store. DERIVED
Modular polynomial $\Phi_\ell(1, Y)$	Degree $\ell + 1 \approx 1.5 \cdot 10^{17}$, with integer coefficients of order $\ell \log \ell$ bits. DERIVED
CM / Hilbert class polynomial	Discriminant $-7\ell^2$, degree $h = 1.5 \cdot 10^{17}$. DERIVED
Random search for an image	A random b lands on that floor with probability 2^{-74} . DERIVED
$\sqrt{\ell}$ u	About $\sqrt{\ell} \approx 2^{28.5}$ operations, but each is in the $2^{59.5}$ -bit field. DERIVED

Every route costs at least about 2^{56} . Pollard rho on ECC2K-130 with Frobenius costs about $2^{60.8}$, so just writing one of these isogenies down costs about as much as solving the discrete logarithm. The same barrier holds for the ℓ -isogenies from the 263-floor and for every curve at the bottom: a random b lies in the class at all with probability 2^{-66} , and every route from a known curve passes through an ℓ -isogeny. **DERIVED**

6. ECDLP consequence (scoped) and limitations

- The 263-isogenies make transferring a discrete logarithm to the 263-floor essentially free. **DERIVED**
- A 263-floor curve has no Frobenius endomorphism; its cheapest non-integer endomorphism has degree $\geq 7 \cdot 263^2/4 \approx 121045$. **DERIVED**
- The volcano studies at $m = 37$ and $m = 41$ measured that rho on such curves runs with negation only, $\approx \sqrt{m}$ times more steps than the crater with Frobenius. At $m = 131$ that predicts about $11.4 \times$ more steps. That is an extrapolation, not measured here. **PROPOSED**

7. Next checks

- Run the audit-curve skill on one representative from each 263-floor orbit: twist, MOV/FR, anomalous and GHS descent checks. **PROPOSED**