Graph-wide discovery capacity and admission gap

Planning assessment, 2026-10-22. Derived inventory and archived observations only; no new scientific trial.

Fixed inventory and open quota

The prospective P29/P37 schedule has 2,049 and 257 vertices. Three native bases on every vertex give **6,918 mandatory discovery arms** and **51,935,232 target probes**: P29 has 6,147 arms at 8,192 targets each; P37 has 771 arms at 2,048 each. Check indices add 257 P29 and 65 P37 vertices. Three full-quota controls per check vertex give a conditional **7,884 arms** and **58,650,624 probes**. A separate full-quota `u=2` transported-base oracle would raise that to **8,206 arms** and **60,889,088 probes**. The additive v2 inventory distinguishes the graph-transport control from that coordinate oracle. The Coordinator must fix its execution mode and all check-arm quotas before admission.

A machine-readable plan now binds the 6,918 main rows to exact archived models and generators, base rules and quotas. Independent point arithmetic checked all 2,306 generator identities and group-order multiplications. Main-arm SHA-256: `bf436f6a3df707b799cb738ed0522aa2a0fa68e921afd6d8099bf929d94105ef`. This plan has no seed or target points and is unapproved.

Three archived 32,768-target observations

P29 E2 native (`RUN-ECTD-c769f1`) recorded 68.37 s wall; P29 E3 native (`RUN-ECTD-f395a9`) 77.68 s; P37 floor `u=2` native (`RUN-ECTD-26d06d`) 952.04 s. These are different curves and bases in a separate 20-arm study. The graph shows their measured collector, count, replay, and interval phase times in seconds. Pair-table setup and target probing were not separated, so no full-scan runtime is inferred.



Figure 1: Archived phase wall times on a logarithmic seconds axis. Three output-validated records at 32,768 targets each; run IDs are shown under the marks.

Next gate

Freeze the check-arm quotas and implementation/seed commitments, then obtain a runnable Coordinator-approved collector contract. The configured Coordinator backend remains unavailable in the current adapter probe. The graph's native yield, useful rank, and full same-point ECDLP costs remain unmeasured. Source paths, exact derived counts, receipt values, and limits are in the accompanying Markdown report and data file.