

Norm-one weak curves at larger field sizes and a prime-degree orbit theorem

ISO-1 research evidence

9 October 2026

Abstract

We verify norm-one weak-curve isogenies across 17 finite fields, including fields with $\log_2 Q \simeq 252$ and extension degrees 10 and 14. All 49 explicit geometric controls and all 17 independently counted source/target pairs pass. We also prove an exact absolute-Frobenius orbit formula for every odd prime extension degree. It specializes to the cubic formula and prices the complete parameter census at degrees 5 and 7. The completed $p = 59$ census adds 100,408 weak ordinary rows and 540 higher-depth zero rows; the finite local worker continues at $p = 61$.

1 Measured larger-field controls

For $q = p^2$ and $K = \mathbf{F}_{q^n}$, generate a nonidentity norm-one parameter λ from a uniformly drawn nonzero field element raised to $q-1$. The fixed seed is 20261009+ i for curve number i . A fourth root $\mu \in \ker N_{K/\mathbf{F}_q}$ gives $s = \mu^2$ and the models

$$E_\lambda : y^2 = x(x-1)(x-\lambda), \quad E'_\lambda : y^2 = x[x^2 + 2(1+\lambda)x + (1-\lambda)^2].$$

With $a = -(1+\lambda)$ and $b = \lambda$, the rational map $(x, y) \mapsto (x+a+b/x, y(1-b/x^2))$ has kernel $\{O, (0, 0)\}$. The retained polynomial certificate IDC1he02ab9a5cf019a56 checks its cleared equation. For each field, the geometric controls verify the norm, fourth root, an explicit source point and its image, and two independent rational halves of target 2-torsion points. Both curves are independently point-counted with PARI/GP 2.17.3. Every pair has equal cardinality divisible by 16 and passes Hasse. All 17 counted curves are ordinary.

p	odd degree n	$\log_2 Q$	paired counts
257, 509, 1009	3	48.03–59.87	3
2003, 8191, 65537	3	65.81–96.00	3
1048583	3	120.00	1
4294967311	3	192.00	1
4398046511119	3	252.00	1
13, 257, 1009, 65537	5	37.00–160.00	4
13, 257, 1009, 65537	7	51.81–224.00	4

These are field-size measurements; a prime-subgroup order is not assigned. The independent counts took 7,822 and 8,630 ms in the largest-field cell; these are stage resource times on the ordinary host. Every cell had a 120-second count cap, and every count completed. The geometry phase had 49 controls and a 30-second cap per field. Source moduli, coefficient vectors, full ICV1 identifiers, seeds, raw outputs, and hashes are retained in [the curve records](#) and [the raw receipts](#).

2 An exact orbit theorem for larger degrees

Let p be an odd prime and $n \geq 3$ an odd prime. Put

$$N = \frac{p^{2n} - 1}{p^2 - 1}, \quad A = \frac{p^n - 1}{p - 1}, \quad B = \frac{p^n + 1}{p + 1}, \quad e = \begin{cases} 1, & p^2 \equiv 1 \pmod{n}, \\ 0, & \text{otherwise.} \end{cases}$$

Theorem 1. *On the nonidentity norm-one subgroup of $\mathbf{F}_{p^{2n}}^\times$ over $\mathbf{F}_{p^2}$, absolute p -Frobenius and inversion preserve the trace pair. The orbit populations are*

size	number of orbits
2	$e(n - 1)/2$
$2n$	$[A + B - 2 - e(n - 1)]/(2n)$
$4n$	$[N + 1 - A - B]/(4n)$.

Therefore one point count per orbit needs exactly

$$C_{p,n} = \frac{N + A + B - 3 + 2e(n - 1)^2}{4n}$$

calls. Adding each orbit's weight to both trace signs recovers the $2(N - 1)$ normalized representatives.

Proof. Hilbert 90 identifies the projective non-base-field parameters with the nonidentity norm-one subgroup T . Its order $N = AB$ is odd. Frobenius preserves cardinality, and inversion gives an isomorphic Legendre curve because every $\lambda \in T$ is a square. The generated action has the $4n$ exponents $\pm p^j$, $0 \leq j < 2n$.

The integers A, B are odd and coprime: a common divisor divides both $p^n - 1$ and $p^n + 1$, hence divides 2. On T_A , p^n acts as the identity; on T_B , it acts as inversion. Their nonidentity union has $A + B - 2$ elements. Since $A > p - 1$, the p -action on T_A has order n ; on T_B it has order $2n$, since p^n acts as inversion and $B > p + 1$ excludes order 2. Inversion is independent of the odd-order action on T_A , so the $4n$ transformations are distinct.

For a further nonidentity stabilizer with $j \neq 0, n$, an element's order must divide both $p^{2j} - 1$ and $p^{2n} - 1$. Since n is prime and $j \neq n$, their common-divisor condition reduces to $p^2 - 1$. Now $\gcd(N, p^2 - 1) = \gcd(n, p^2 - 1)$, equal to n precisely when $e = 1$. If present, the $n - 1$ nontrivial n th roots each have orbit size 2, since $p \equiv \pm 1 \pmod{n}$. Every other element of $T_A \cup T_B$ has stabilizer of size 2 and orbit size $2n$. Outside their union the stabilizer is trivial and the orbit size is $4n$. Summing gives the formula. Both scaling square classes remain available in an odd extension and supply the two trace signs. $\square$

For $n = 3$ and $p > 3$, this is $(p^4 + 3p^2 + 8)/12$. At $n = 5, 7$, the exceptional term depends on $p^2 \pmod{n}$.

The cleared orbit sum has record `IDC1h90d58cc0e0c48fe3`, with 32 recorded plus 64 additional evaluations and mutation rejection. Exact GP expansion independently returns zero. Eight exhaustive cyclic group audits cover 6,928,970 nonidentity parameters and agree with all three orbit populations. Forty-five stabilizer cases also pass. The record checks the polynomial sum; the stabilizer classification is the deductive argument above.


3 Complete census work and continuation

At $p = 257, 509, 1009$ in the cubic family, the exact call counts are 363,555,713; 5,593,645,151; and 86,374,331,401. At $p = 13$, moving from $n = 3$ to 5 to 7 changes that count from 2,423 to 41,032,186 to 837,027,639,705. These are algebraic work counts, not measured durations. The larger-field controls

verify individual positive weak models; complete class labels still require exhaustive enumeration or a sufficient class-existence criterion.

The earlier temporary worker and directory were absent at this round's start. The original every-prime request through 199 is preserved. The $p = 59$ census has completed and passed its audit. Of 201,898 ordinary rows, 100,408 are weak, with zero depth-1 weak rows and 540 higher-depth zero rows. All 5,000 independent GP controls are positive. The worker has advanced to $p = 61$ using four workers and an internal-volume runtime copy, checked against the frozen sources and executables. A finite `launchd` service owns the process independently of this chat; `KeepAlive=false` preserves a failure or completion without restarting it. The retained status and launch configuration identify the actual process and persistent output location.

Verified rational 2-isogeny over $F_{13^{10}}$



Full ICV1 IDs name the short models; coordinate changes and map are recorded.


Independent point counts agree; explicit point and target-halving replays pass.

Figure 1: A counted degree-10 example. Full source and target ICV1 IDs name the short models; the retained records specify their coordinate changes and the original-model map.


4 Sources and validation

The [PARI reference](#) documents `ellcard` on finite extensions and the available counting algorithms. The existing [norm-one theorem](#) supplies the conductor obstruction at every size. The dated Markdown report links all code, exact field and curve records, the cyclic audits, native tests, failed preliminary attempts, and the persistent-service receipts.


Larger fields verified; complete-census work remains explicit Verified norm-one curves over $F_{(p^{(2n)})}$



Exact orbit work for a complete parameter census



Complete observed census through $p = 59$



Top: 17 completed point-count pairs; every plotted marker is a measured field case. Middle: exact orbit-call requirements from the proved formula, displayed on a logarithmic scale. Bottom: complete observed ordinary trace strata through $p = 59$. Orange is depth 1, green is observed weak, and yellow is higher-depth zero; those census labels retain their randomized point-count uncertainty.