An explicit genus-3 plane quartic over F_q for an elliptic curve over F_{q³}

2026-10-12. Conductor T-2267. **Structural stage diagnostic on toy fields.** A curve is built and its Frobenius checked. No discrete logarithm is solved, no index calculus is run, and no speedup is claimed. Protocol frozen before the first run: PROTOCOL.md. Run `20261013T0320Z`: records, environment, run report. Addendum run `20261013T0347Z` (the order-332 class at q = 7, see below): records, environment, run report. This report as PDF: REPORT.pdf. Follows `research/ghs_cubic_cover_census_20261012/` (PR #1759), which counted these covers by genus without building them.

Question


The census found that every isogeny class with `4 | #E(K)` over `K = F_{q³}` has a genus-3 cover by branch data. Does that cover exist as a curve over `F_q` whose Jacobian has `#E(K)` rational points, and is the transfer of points an explicit `F_q`-rational object?

Result

Yes on all eleven cases built (`q ∈ {5, 7, 11}`, `#E0(K) = 4ℓ`; nine in the first run, the two of the order-332 class at q = 7 in the addendum). For each:

- the curve is a **smooth plane quartic with coefficients in `F_q`**, so it is not hyperelliptic;
- its point counts over `F_q`, `F_{q²}`, `F_{q³}` give L-polynomial `1 − t_E T³ + q³T⁶`, hence **`#Jac(C)(F_q) = #E0(K)`**;
- every transfer divisor tested is an `F_q`-rational divisor on it.

So, in the eleven built cases, the covers that the census of PR #1759 counted by branch data exist as curves over `F_q` with the predicted Jacobian order. That is a statement about these eleven cases only, not about every cover the census counted. What is still not shown by execution is the transfer as a group homomorphism, because no class arithmetic on a plane quartic is implemented (problem record).



Construction (derived; two steps in Lean)

`u ∈ K[t]` monic quadratic, `u' = u^σ`, `u'' = u^{σ²}`, six distinct roots, `u, u', u''` linearly independent.

1. `E0 : y0² = u·u''`, `E1 = E0^σ : y1² = u·u'`, `E2 : y2² = u'·u''` with `y2 = y0·y1/u` (`DescentCycle.third_cover`).
2. `σ~ : y0 ↦ y1 ↦ y2 ↦ y0`, `t ↦ t`, σ on K, has order 3 (`DescentCycle.cycle_closes`), so the compositum descends to `F_q`.
3. With `X = 1/y0, Y = 1/y1, Z = 1/y2`: `(u : u' : u'') = (Z² : X² : Y²)`. If `(1, t, t²)ᵀ = R·(u, u', u'')ᵀ` then `t·t = 1·t²` reads `G = (R₁·w)² − (R₀·w)(R₂·w) = 0, w = (Z², X², Y²)`: a quartic in `X², Y², Z²`. The three differentials `dt/y_i` are holomorphic and `(X : Y : Z)` is the canonical map, so a smooth `G` means `C` is not hyperelliptic.
4. σ~ permutes `X → Y → Z`. In `W_j = ω_j X + ω_j^σ Y + ω_j^{σ²} Z, ω = (1, g, g²)`, the quartic has `F_q` coefficients up to a scalar.
5. Transfer `φ = (1 + Fr + Fr²)∘π0^*`, with `Fr(t, y0, y1, y2) = (t^q, y2^q, y0^q, y1^q)`. `π_j_* ∘ π_0^* = 0` for `j ≠ 0` (it factors through the `t`-line), so `π0_* ∘ φ = [2]` and `ker φ ⊂ E0[2]`. With `#Jac(C)(F_q) = #E0(K) = 4ℓ`, φ maps `E0(K)[ℓ]` isomorphically onto `Jac(C)(F_q)[ℓ]`.

Report statement	Lean declaration	Status
`(u·y2)² = u²·u'·u''`	`third_cover`	proved, any commutative multiplication
`u·(y1·y2) = (u·u')·y0` (cycle closes)	`cycle_closes`	proved
cancelling u; the function field; descent; step 5	none	**not formalized**

Lean: lean/DescentCycle.lean, toolchain v4.34.1, no Mathlib, no axioms.

Identity certificates (`identity.certificate/v1`, see docs/identity-certificates/README.md). The two Lean statements are conditional on the curve equations, so they are not themselves polynomial identities. The polynomial identities underneath them are the cofactor expansions of the conclusion differences in the relations `y0² − u·u''`, `y1² − u·u'`, `u·y2 − y0·y1`, and those are certified, with the Lean file as the `proof` pointer (recorded, not run by the checker):

Identity	Certificate	File
`(u·y2)² − u²·u'·u'' = (u·y2 − y0·y1)(u·y2 + y0·y1) + y1²(y0² − u·u'') + u·u''(y1² − u·u')`	`IDC1h4304bc232b34fa40`	third_cover.json
`u·y1·y2 − u·u'·y0 = y1(u·y2 − y0·y1) + y0(y1² − u·u')`	`IDC1h14765be340922acc`	cycle_closes.json

Issued and self-checked by `examples/identity_certificate_ghs_cubic_genus3.rs` (32 points each). Not certified: the step from the cofactor identity to `y2² = u'·u''` (cancelling `u`), the function field, the descent, and the quartic equation `G` of step 3, which is a linear-algebra consequence over `K` of the matrix `R` and is instead checked by the point counts below.

Measurements (run 20261013T0320Z, exact counts)

Tool: genus3_build.rs. The records of both runs were regenerated after review (see "Review changes" below); every pre-existing field is unchanged. `u = t² + bt + c` in lexicographic order of field index; the first cases with `#E0(K) = 4ℓ`. `N_k` are projective point counts of the `F_q` quartic over `F_{q^k}` by exhaustion; `N3` was also counted on the `K` model and agreed in all nine cases.

q	u (index of b, c)	roots of u in K	#E0(K)	t_E	in F_q	smooth	N1, N2, N3	predicted	#Jac(C)(F_q)
5	5, 30	0	148 = 4·37	−22	yes	yes	6, 26, 192	6, 26, 192	148
5	5, 51	2	148 = 4·37	−22	yes	yes	6, 26, 192	6, 26, 192	148
7	7, 52	0	356 = 4·89	−12	yes	yes	8, 50, 380	8, 50, 380	356
7	7, 54	0	356 = 4·89	−12	yes	yes	8, 50, 380	8, 50, 380	356
7	7, 110	2	356 = 4·89	−12	yes	yes	8, 50, 380	8, 50, 380	356
7	7, 118	2	356 = 4·89	−12	yes	yes	8, 50, 380	8, 50, 380	356
7	7, 129	2	356 = 4·89	−12	yes	yes	8, 50, 380	8, 50, 380	356
11	11, 130	0	1348 = 4·337	−16	yes	yes	12, 122, 1380	12, 122, 1380	1348
11	11, 172	2	1324 = 4·331	8	yes	yes	12, 122, 1308	12, 122, 1308	1324
7†	49, 8	0	332 = 4·83	12	yes	yes	8, 50, 308	8, 50, 308	332
7†	49, 9	0	332 = 4·83	12	yes	yes	8, 50, 308	8, 50, 308	332

† Addendum run `20261013T0347Z`: the second 4ℓ isogeny class at `q = 7` (`#E0(K) = 332`, trace +12, the sign opposite to the 356 class's −12). The frozen first-in-index rule of the first run reached only the 356 class. Exhausting the parametrization `u = t² + bt + c`, this class has exactly two irreducible-u representatives satisfying the genus-3 conditions and **no split-u representative** (65432 candidates examined). The order-356 run was not exhaustive, so this is not a multiplicity comparison between the two classes.

Example, `q = 7`, u index (7, 110), over `F_7`:

```
W2⁴ + W1²W2² + 6W1³W2 + 6W1⁴ + 4W0W2³ + W0W1W2² + 4W0W1²W2 + 4W0W1³
  + W0²W2² + 4W0²W1W2 + 5W0²W1² + 6W0³W2 + 6W0⁴ = 0
```

All eleven quartics are in the records (`quartic_Fq`, entries `[i, j, k, coefficient]` for `W0^i W1^j W2^k`; the two order-332 quartics are in the addendum run's records).


Transfer divisors (P4). For each affine `P = (t0, y0) ∈ E0(K)` with `y0 ≠ 0` and `u·u'(t0)` a non-zero square, the two points of `C` over `P` and their two Frobenius images each. `#E0(K)` is accounted exactly: `#E0(K) = tested + nonsquare + ramification + 2`, where the 2 are the points at infinity of the quartic model (never tested), `ramification` are the points with `y0 = 0` (never tested; 0 when u is irreducible, 4 when u splits), and `nonsquare` are the affine points with `y0 ≠ 0` for which `u·u'(t0)` is zero or a non-square (their preimages are not K-rational or are ramified, and need a field the tool does not build). The records carry all four counts (`e0_points_tested`, `e0_affine_points_nonsquare_uu1`, `e0_ramification_points_y0_zero`, `e0_points_at_infinity`) and the tool asserts the sum.

q	u	#E0(K)	E0 points tested	tests (6 per point)	on the F_q quartic	q-power = model Fr	projected point satisfies the E0 equation	nonsquare u·u'	y0 = 0	infinity
5	5, 30	148	94	564	564	564	564	52	0	2
5	5, 51	148	94	564	564	564	564	48	4	2

q	u	#E0(K)	E0 points tested	tests (6 per point)	on the F_q quartic	q-power = model Fr	projected point satisfies the E0 equation	nonsquare u·u'	y0 = 0	infinity
7	7, 52 / 7, 54	356	188	1128	1128	1128	1128	166	0	2
7	7, 110 / 118 / 129	356	182	1092	1092	1092	1092	168	4	2
11	11, 130	1348	688	4128	4128	4128	4128	658	0	2
11	11, 172	1324	646	3876	3876	3876	3876	672	4	2
7†	49, 8 / 49, 9	332	152	912	912	912	912	178	0	2

† Addendum run `20261013T0347Z` (order-332 class).

"Projected point satisfies the E0 equation" (record field `projected_points_on_E0`) is only the check that the `(t, y0)` coordinates of each tested point satisfy `y0² = u·u''`. It is **not** the pushforward `π0_*` of a divisor and does not verify `π0_*φ(P) = [2]P`; that check is not executed (see Scope and limits).



All eleven cases (nine from the first run, two order-332 from the addendum); the run-20261013T0320Z report keeps the nine-case graph. P1, P2, P3 and P4 hold on every case; no falsifier fired.

Scope and limits

- **Not executed:** the pushforward `π0_*` of a divisor and the identity `π0_*φ(P) = [2]P`; `φ` as a homomorphism, its injectivity on `E0[ℓ]`, any discrete logarithm, any relation collection. These need class arithmetic on the quartic; see the problem record.
- **Selection.** The frozen rule takes the first cases in index order. All five `q = 7` cases of the first run landed in one isogeny class (`#E0(K) = 356`); the other 4ℓ class at `q = 7` (332) was built in the addendum run after a target-order argument was added to the tool (the argument defaults to off, so the first run reproduces byte for byte). Most candidates were skipped as linearly dependent (2394 of 2443 at `q = 7`) because `b ∈ F_q` comes first in index order.
- **One construction.** Only the `m = 2, r = 6` type. The hyperelliptic `m = 3, r = 5` type is not built.
- **Cost (published asymptotics, not measured):** index calculus on a smooth plane quartic over `F_q` is reported at about `q` operations (Diem), against about `q^{3/2}` for rho on `E(K)`. These toy fields are far too small to compare anything.
- `n = 3`, `q ∈ {5, 7, 11}`, odd characteristic above 3.

Next discriminating step

Implement linear equivalence on a plane quartic and execute the transfer on the `q = 7` case (7, 110); then relation collection by lines through pairs of `F_q`-points, counted against rho. The acceptance conditions are in the problem record.

Curve identity (ICV1)

Each source `E0` is recorded as the short Weierstrass model `Y² = X³ − 27·I·X − 27·J` of the Jacobian of the quartic `y0² = u·u''` (`I, J` the quartic's invariants; the quartic has `K`-rational points at infinity, so the Jacobian is `E0`), over `F_{q³} = F_q[x]/`

$(x^3 + c_2x^2 + c_1x + c_0)$ with the modulus the tool built. The tool emits the field modulus and `(a, b)` as coefficient lists (`E0_weierstrass_ext`) and asserts that the Weierstrass model has the same order as the quartic. reports/e0_icv1.py recounts that order independently with `scripts/curve_id.py` and computes each model's ICV1 identity, written to results/e0_icv1.json (field modulus, `a`, `b`, order, ICV1 string, model JSON, for all 11 cases). The order agrees with `order_E0` in all eleven.

Not registered yet. `docs/curves/registry.json` has not been updated: registering needs `scripts/build_curve_registry.py` and a harvester for this record shape, under `docs/curves/`, which is reserved by Conductor task T-2240. Until the registration PR lands, this README writes no slug (an unregistered slug fails `scripts/check_curve_names.py`); cases are named by `(q, u index)` as in the tables, and the exact binding of each case to its field and model is `results/e0_icv1.json`.

Review changes

- `points_skipped` counted only the affine points with non-square or zero `u·u'`; it omitted the 2 points at infinity and, when `u` splits, the 4 points with `y0 = 0`. Replaced by `e0_affine_points_nonsquare_uu1` plus the other explicit counts above; the original values are unchanged.
- `pushforward_on_E0` renamed `projected_points_on_E0` (values unchanged).
- Runs `20261013T0320Z` and `20261013T0347Z` regenerated with the updated tool (new `source sha256` in each `.env.txt`); `all11.jsonl` rebuilt.

Graphs checked and left unchanged

No canonical graph changes (no ECDLP rung, no timing; no `benchmarks/` entry).

Reproduce

```
rustc --edition 2021 -O genus3_build.rs && ./genus3_build 5 1 1 && ./genus3_build 7 3 2 && ./genus3_build
11 1 1
./genus3_build 7 3 2 332                # addendum: the order-332 class at q = 7
python3 reports/plot.py results/run-20261013T0320Z.jsonl reports/run-20261013T0320Z   # layout only
cat results/run-20261013T0320Z.jsonl results/run-20261013T0347Z.jsonl > results/all11.jsonl
python3 reports/plot.py results/all11.jsonl reports/run-20261013T0347Z   # then orders.* renamed to
orders_all11.*
cd lean && lake build
python3 reports/e0_icv1.py                # from the repository root: results/e0_icv1.json
cargo run --example identity_certificate_ghs_cubic_genus3 -- research/ghs_cubic_genus3_build_20261012/
certificates
```

The fourth argument to `genus3_build` is an optional target order (0 or absent keeps the original first-in-index rule); `./genus3_build 7 3 2` reproduces the q = 7 records of run `20261013T0320Z` byte for byte.