

feynsage

Feynman Integrals in SageMath, from Graphs to Numbers

Kazi Abu Rousan

Built on the notes of Prof. B. Ananthanarayan's lecture "Feynman integrals at one loop", NISER, 29 September 2026

github.com/aburousan/feynsage · MIT licence · every feynsage result in this talk was computed by the package

Why another package?

Today a one-loop or multi-loop calculation usually means several programs:

- Package-X, FeynCalc and LiteRed need Mathematica; FIRE uses it for set-up around a C++ reducer
- LoopTools gives numbers, not formulas
- Kira and FIRE reduce, but do not draw or explain

We wanted **one** free tool where every step is exact and can be checked.

FEYNSAGE

- written natively for SageMath (free, Python)
- exact algebra in $\mathbb{Q}(D, s, t, m, \dots)$ and graph theory come built in
- one loop like Package-X, many loops like LiteRed / Kira
- FORM for the Dirac algebra
- every function explains its output: `explain=True`

What is inside

Step	feynsage	Plays the role of
diagram	<code>graph("A-B, A-B:m", legs, kin), diagram("kite"), pictures</code>	pen and paper
$\mathcal{U}, \mathcal{F}$	spanning trees, 2-forests, Kirchhoff, $\det M$	the speaker's notebook, FeynCalc
family, sectors	<code>family([...])</code> , zero sectors, symmetries	LiteRed
IBP	Laporta; finite fields + rational reconstruction	FIRE, Kira
one loop	<code>loop(...)</code> $\rightarrow A_0, B_0, C_0, D_0$	Package-X, FeynCalc
values	closed forms of C_0, D_0 ; IR poles	Package-X, LoopTools
Dirac traces	<code>form.dirac_trace(...)</code> , γ_5 included	FORM, FeynCalc

Three lines to start

```
from feynsage import *
loop("l^mu l^nu", ["l", "m"], ["l + p", "m"], kin={"p^2": "s"})      # one loop, exact
ibp_reduce(family(["l1", "l1 + q", "l1 + l2", "l1 + l2 + q", "l2"], kin={"q^2": 1}),
            ["F(2,2,1,2,2)"])      # two loops, IBP
diagram("kite").plot()      # a picture
```

Install: `git clone ... && ./install.sh` (macOS, Linux, WSL; it finds or installs Sage and FORM).

NOTATION, AS IN THE LECTURE NOTE

$D = 4 - 2\varepsilon$, loop momentum ℓ , Feynman parameters x_i , graph polynomials $\mathcal{U}$ and $\mathcal{F} = \mathcal{F}_0 + \mathcal{U} \sum_i x_i m_i^2$. One-loop functions in the Package-X normalisation $\mu^{2\varepsilon} e^{\varepsilon\gamma_E} \int d^D \ell / (i\pi^{D/2})$.

$\mathcal{U}$ and $\mathcal{F}$: the idea of the lecture

The speaker's notebook (uf-new-short.nb) completes the square, loop by loop:

$$\sum_i x_i D_i = \sum_{r,s} \ell_r \cdot \ell_s M_{rs} - 2 \sum_r \ell_r \cdot Q_r + J$$


$$\mathcal{U} = \det M, \quad \mathcal{F} = \det M (J - Q^T M^{-1} Q)$$

and gives four examples. feynsage reproduces all four, **two ways**:

diagram	from propagators	from the graph
massless triangle	✓	✓
bubble, two masses	✓	✓
massless box	✓	✓
two-loop sunset	✓	✓

READ OFF THE GRAPH

- $\mathcal{U}$: cut L lines so that a spanning tree is left, multiply their x_i
- $\mathcal{F}_0$: cut $L + 1$ lines so that two trees are left, times the momentum² flowing between them
- $\mathcal{U}$ again: det of the Kirchhoff matrix (matrix-tree theorem)



the 8 spanning trees of the kite, drawn by feynsage

$\mathcal{U}$ and $\mathcal{F}$: the same idea, pushed further

The notebook stops at two loops. Here are bigger diagrams, each done **three independent ways** inside feynsage, and then by FeynCalc's FCFeynmanPrepare:

diagram	L	N	trees	terms of $\mathcal{F}$	trees = $\det M$	Kirchhoff = trees	= FeynCalc	time (trees)
kite	2	5	8	8	✓	✓	✓	< 10 ms
double box	2	7	15	11	✓	✓	✓	< 10 ms
banana (3 loops)	3	4	4	13	✓	✓	✓	< 10 ms
triple box (3 loops)	3	10	56	65	✓	✓	✓	10 ms

The “tennis court” we first tried had a 2-valent vertex (two lines with the same momentum): FeynCalc merges such lines into one doubled propagator, feynsage keeps two parameters. Both are right; the triple box above is the clean test.

kite: $\mathcal{U} =$

$$x_1x_3 + x_2x_3 + x_1x_4 + x_2x_4 + x_1x_5 + x_2x_5 + x_3x_5 + x_4x_5$$


$\mathcal{F} =$

$$-sx_1x_2x_3 - sx_1x_2x_4 - sx_1x_3x_4 - sx_2x_3x_4 - sx_1x_2x_5 - sx_2x_3x_5 - sx_1x_4x_5 - sx_3x_4x_5$$

Two loops: IBP reduction of the kite

```
kite = family(["l1", "l1 + q", "l1 + l2", "l1 + l2 + q", "l2"],
kin={"q^2": 1})
ibp_reduce(kite, ["F(2,2,1,2,2)"])
```

$$F(2,2,1,2,2) = \frac{(-4) \cdot (d-9) \cdot (d-6) \cdot (d-5) \cdot (d-3)}{(d-8)} F(1,1,1,1,0) \\ + \frac{(-729) \cdot (d-7) \cdot (d-\frac{16}{3}) \cdot (d-\frac{14}{3}) \cdot (d-\frac{10}{3}) \cdot (d-\frac{8}{3}) \cdot (d^2-11d+20)}{(d-10) \cdot (d-4) \cdot (d-8)^2} F(0,1,1,0,1)$$



Two master integrals: the product of two bubbles $F(1,1,1,1,0)$ and the sunset $F(0,1,1,0,1)$. The coefficients are **identical** to LiteRed and Kira 3.1.

kite, seeds up to	equations	exact Laporta	finite fields	Kira 3.1
4 dots, 2 numerators	9 368	7.3 s	0.2 s	
9 dots, 4 numerators	157 843	too slow	2.7 s	6.7 s*

*Kira on a server with one thread, feynsage on a laptop: a rough guide only.

One loop, the Package-X way

The rank-2 bubble with all symbols, 0.03 s:

```
loop("l^mu l^nu", ["l", "m"], ["l + p", "m"], kin={"p^2": "s"})
```

$$g^{\mu\nu}: \quad \frac{1}{6}m^2 \left(\frac{1}{\epsilon} + \log\left(\frac{\mu^2}{m^2}\right) + 1 \right) + \frac{1}{3}m^2 + \frac{1}{12}(4m^2 - s) \left(\frac{1}{\epsilon} + \Lambda(s; m, m) + \log\left(\frac{\mu^2}{m^2}\right) + 2 \right) - \frac{1}{18}s$$

$$p^\mu p^\nu: \quad \frac{m^2 \left(\frac{1}{\epsilon} + \log\left(\frac{\mu^2}{m^2}\right) + 1 \right)}{3s} - \frac{m^2}{9s} - \frac{(m^2 - s) \left(\frac{1}{\epsilon} + \Lambda(s; m, m) + \log\left(\frac{\mu^2}{m^2}\right) + 2 \right)}{3s} - \frac{4m^2 - s}{18s}$$

$\Lambda(s; m, m)$ is Package-X's DiscB.

- projection on a symmetric basis, Gram matrix inverted **exactly** (adjugate)
- rational terms from $D = 4 - 2\epsilon$, kept to $\mathcal{O}(\epsilon^2)$ when IR poles appear
- DiscB, LogM carry the $+i\epsilon$ branch, as in Package-X
- PVB, PVC, PVD: Package-X's coefficient functions
- Euclidean or Minkowski: euclidean=True
- numerators from FORM traces go straight in

C_0 and D_0 in closed form

`explicit(C0(0, 0, 5, 1, 1, 1))`, exact, with the branches fixed by the $+i\epsilon$:

$$\begin{aligned} & -\frac{2}{15}\pi^2 + \frac{1}{5}i\pi \log\left(\frac{10}{\sqrt{5}+5}\right) + \frac{1}{10} \log\left(\frac{10}{\sqrt{5}+5}\right)^2 \\ & -\frac{1}{5}i\pi \log\left(-\frac{10}{\sqrt{5}-5}\right) + \frac{1}{10} \log\left(-\frac{10}{\sqrt{5}-5}\right)^2 + \frac{1}{5}\text{Li}_2\left(\frac{1}{10}\sqrt{5} + \frac{1}{2}\right) \\ & + \frac{1}{5}\text{Li}_2\left(-\frac{1}{10}\sqrt{5} + \frac{1}{2}\right) \end{aligned}$$

`.n(digits=30)` gives `-0.8943345118780581048846775 - 0.6047086137711147833763413 i`

D_0 : Denner's 16 dilogarithms (no massless line), Denner–Dittmaier for one to four massless lines. `explicit(D0(1,2,3,4,-5,-6,1,1,1,1))` is exact, 8852 characters long, value `0.27279190436796813983`.



HOW EVERY BRANCH IS FIXED

Each quantity is carried twice: as an 80-digit number with a tiny $+i\epsilon$ and as an exact expression. Where an argument lies on a cut, the number decides the side. At the end the two are compared at two precisions.

Through the thresholds

```
quick_plot([B0(s, 1, 1), C0(0, 0, s, 1, 1, 1)], (s, -4, 12), parts="both")
```

finite part (ϵ^0) at $\mu=1$



Soft and collinear poles

All 6 triangles and 16 boxes of Ellis and Zanderighi, any order of the lines, in the Package-X normalisation:

$C_0(0, 5, 0, 0, 0, 0)$ (massless, $s = 5$, $\mu = 1$):

$$\frac{1}{5\epsilon^2} + \frac{1}{\epsilon} \left(\frac{1}{5}i\pi - \frac{1}{5}\log(5) \right) - \frac{7}{60}\pi^2 - \frac{1}{5}i\pi\log(5) + \frac{1}{10}\log(5)^2$$

$D_0(0,0,0,0, -2,-3, 0,0,0,0)$ (massless box, $\mu = 1$):

$$\frac{2}{3\epsilon^2} + \frac{1}{\epsilon} \left(-\frac{1}{3}\log(3) - \frac{1}{3}\log(2) \right) - \frac{2}{9}\pi^2 + \frac{1}{3}\log(3)\log(2)$$

QED vertex $C_0(m^2, q^2, m^2; 0, m, m)$, photon massless, electrons on shell ($m = 1$, $q^2 = -3$): $1/\epsilon$ coefficient 0.3419036239, finite part -0.1252442693 .

CHECKED

118 points against Package-X (all three orders in ϵ , double precision). Boxes 15 and 16, whose finite parts Package-X cannot evaluate, against a photon mass λ : $\log \lambda^2 \rightarrow 1/\epsilon + \log \mu^2$.

When the Gram determinant vanishes

The $g - 2$ vertex at $q^2 = 0$: $p_1 = p_2$, so the usual reduction divides by zero. feynsage takes the coefficients straight from Feynman parameters (lines with identical kinematics are grouped and integrated out exactly):

```
PVC(0, 1, 0, m^2, 0, m^2, 0, m, m)
PVC(1, 0, 0, m^2, 0, m^2, 0, m, m)
```

$$C_1 = \frac{1}{2m^2}$$

$$C_{00} = \frac{1}{4\epsilon} - \frac{1}{2} \log(m) + \frac{1}{2} \log(\mu) + \frac{1}{4}$$

PACKAGE-X

$$C_1 = 1/(2m^2), \quad C_{00} = \frac{1}{4} + \frac{1}{4} \left(\frac{1}{\epsilon} + \log \frac{\mu^2}{m^2} \right) \quad \checkmark$$

Also all invariants zero (divided differences, any masses) and a massive photon: 11 more points agree with Package-X.

Real physics, computed

QED, PESKIN AND SCHROEDER

- vacuum polarisation: Ward identity $C_g + q^2 C_{qq} = 0$ **exactly**
- Peskin (7.91) reproduced to 10^{-15}
- running of α from the leptons at one loop: $\Delta\alpha(M_Z^2) = 0.031419$
- electron $g - 2$: $\mathcal{U}, \mathcal{F}$ from the vertex graph, Dirac algebra in FORM, $F_2(0) = \alpha/(2\pi)$ **exactly**

$\pi^0 \rightarrow \gamma\gamma$ FROM THE QUARK TRIANGLE

- trace with γ_5 in FORM:
 $4im \varepsilon(k_1, k_2, \mu, \nu)$
- triangle from its graph polynomials
- width 7.79 eV vs PrimEx-II 7.80 ± 0.12 eV
- lifetime 8.35×10^{-17} s vs PDG 8.43×10^{-17} s

Both are worked out in the notebooks `examples/peskin_examples.ipynb` and `examples/feynsage_walkthrough.ipynb`.

Comparison with other packages

	feynsage	Package-X	LoopTools	FeynCalc	LiteRed	FIRE	Kira
needs Mathematica	no	yes	no	yes	yes	yes ^a	no
one-loop tensor reduction	✓	✓	numbers	✓			
C_0 as a formula	✓	✓		via Package-X			
D_0 as a formula	✓	IR cases	numbers	via Package-X			
IR poles of C_0 , D_0	✓	✓	✓	via Package-X			
multi-loop IBP	✓				✓	✓	✓
finite fields	✓					✓	✓
$\mathcal{U}$, $\mathcal{F}$ from the graph	✓			✓			
Dirac algebra	FORM	Spur		✓			

An empty cell: we did not find this feature in that program. ^aFIRE has a C++ core and a Mathematica front end.

Head to head: the same job, measured

Speed (same laptop, symbols everywhere)

task	feynsage	Package-X
C_{001}	0.42 s	0.95 s
C_{112}	0.39 s	1.59 s
D_{00}	6.0 s	0.08 s
10 numerical D_0	0.23 s	0.25 s
digits of a numerical D_0	≈ 40	16

Package-X is much faster for symbolic box tensors: honest numbers.

Agreement

against	result
Package-X, 190 points	every digit of its values
Package-X, IR, 118 points	double precision
LoopTools 2.16 (Denner), hard D_0 points	10^{-15}
Kira 3.1, LiteRed	identical coefficients
FeynCalc, $\mathcal{U}$ and $\mathcal{F}$	identical polynomials
D_0 relabellings	all 24 at 15 points; 6 random ones at 2000 random points

A point where Package-X and LoopTools fail

$C_0(-4, -3, -1; 0, 0, 1)$: every invariant is spacelike, so $C_0 = -\int d^3x \delta(1 - \sum x_i)/\mathcal{F}$ must be **real and negative**.

program	value
feynsage (closed form)	-0.585976809672364723
Mathematica NIntegrate (no Package-X)	-0.585976809672364723
FeynCalc FCFeynmanParametrize, integrated	-0.58597680967236472
mpmath, 45 digits	-0.585976809672364722650390572218069267
Package-X 2.1.1, two labellings	+0.671253, -0.721045
LoopTools 2.16, version a (FF)	+0.671253, -0.721045, -1.843207
LoopTools 2.16, version b (Denner)	four different complex numbers

At nearby points Package-X and LoopTools' FF version agree with NIntegrate to 16 digits (the Denner version is still wrong there). Package-X's wrong values are LoopTools' FF values digit for digit.

How we know it is right

- 12 test files and a random stress test, run on a laptop (Sage 10.9) and on a CentOS 7 server (Sage 10.7)
- every formula checked against another program **or** a direct integral
- D_0 is symmetric under the 24 relabellings of its lines: all 24 agree at 15 points, 6 random ones at each of 2000 random points
- exact expressions are evaluated at two precisions before they are returned
- two independent code audits; every finding fixed and turned into a test

LIMITS, HONESTLY

- finite-field IBP: D and at most one invariant
- closed forms of C_0 , D_0 need numerical kinematics; with symbols they stay symbols
- vanishing Gram determinant: one group of identical lines, or two groups with equal masses
- symbolic box tensors are slower than Package-X

Summary

FEYNSAGE

- graphs $\rightarrow \mathcal{U}, \mathcal{F} \rightarrow$ families $\rightarrow$ IBP $\rightarrow$ masters
- one loop: exact tensor reduction, C_0 and D_0 in closed form, IR poles, zero Gram determinants
- free, in SageMath, checked against Package-X, LoopTools, FeynCalc, LiteRed and Kira

```
git clone https://github.com/aburousan/feynsage
cd feynsage && ./install.sh --test
```

Start with `examples/feynsage_walkthrough.ipynb`.

Thanks to Prof. B. Ananthanarayan for the lecture and the `uf-new-short.nb` notebook that started the $\mathcal{U}, \mathcal{F}$ part.